

#### Home

2. Zarah

## Zarah

Generic name: drospirenone and ethinyl estradiol [ dro-SPY-re-nown, ETH-in-il, ESS-tra-dy-ol ]

Brand names: Jasmiel, Loryna, Nikki, Ocella, Syeda, ... show all 9 brands

**Drug class: Contraceptives** 



Medically reviewed by Drugs.com on Apr 4, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

## What is Zarah?

Zarah is a combination birth control pill containing female hormones that prevent ovulation (the release of an egg from an ovary). Zarah also causes changes in your cervical mucus and uterine lining, making it harder for sperm to reach the uterus and harder for a fertilized egg to attach to the uterus.

Zarah is used as contraception to prevent pregnancy. This medicine is also used to treat moderate acne in women who are at least 14 years old and have started having menstrual periods, and who wish to use birth control pills.

Yaz is also used to treat the symptoms of premenstrual dysphoric disorder (PMDD), such as anxiety, depression, irritability, trouble concentrating, lack of energy, sleep or appetite changes, breast tenderness, joint or muscle pain, headache, and weight gain.

Zarah may also be used for purposes not listed in this medication guide.

## Zarah side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have:

- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance;
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood, pain or warmth in one or both legs;
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;
- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing
  of the skin or eyes);

- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears;
- swelling in your hands, ankles, or feet;
- a change in the pattern or severity of migraine headaches; or
- symptoms of depression--sleep problems, weakness, tired feeling, mood changes.

#### Common side effects of Zarah may include:

- · nausea, vomiting;
- breast tenderness;
- headache, mood changes, feeling tired or irritable;
- · weight gain; or
- changes in your menstrual periods, decreased sex drive.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

<u>Zarah side effects</u> (more detail)

# Related/similar drugs

#### **Zoloft**

Zoloft is an antidepressant used to treat major depression, obsessivecompulsive disorder, panic ...

#### **Reviews & ratings**

7.3 / 10

1,972 Reviews

#### View more

#### **Fluoxetine**

Fluoxetine is used to treat depression or obsessive-compulsive disorder in adults. Learn about side ...

## Reviews & ratings

7.3 / 10

2,232 Reviews

#### View more

#### FEATURED

#### **Ozempic**

Learn about Ozempic (semaglutide)

for type 2 diabetes treatment, weight management, cardiovascular ...

## **Reviews & ratings**

6.6 / 10

1,504 Reviews

#### View more

## **Prozac**

Prozac (fluoxetine) is an SSRI antidepressant used to treat depression, OCD, panic disorder ...

## Reviews & ratings

7.2 / 10

1,239 Reviews

#### View more

## **Paxil**

Paxil is used to treat depression, obsessive-compulsive disorder and anxiety. Learn about side ...

## Reviews & ratings

7.1 / 10

960 Reviews

#### View more

## Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

## **Reviews & ratings**

6.1 / 10

2,624 Reviews

#### View more

## **Depo Provera**

Depo-Provera is used as contraception to prevent pregnancy. It is also used to reduce pain cause by

- - -

## Reviews & ratings

4.2 / 10

1,967 Reviews

View more

## Medroxyprogesterone

Medroxyprogesterone (Provera) treats conditions such as absent or irregular menstrual periods, or ...

#### **Reviews & ratings**

4.5 / 10

2,469 Reviews

View more

#### **Accutane**

Accutane (isotretinoin) is a form of vitamin A and is used to treat severe nodular acne. Includes ...

## **Reviews & ratings**

7.9 / 10

657 Reviews

View more

## **Nexplanon**

Nexplanon is a long-term, reversible birth control implant that lasts for three years. Learn about ...

#### **Reviews & ratings**

5.0 / 10

4,864 Reviews

View more

# **Warnings**

Do not use if you are pregnant or if you recently had a baby.

You should not take Zarah if you have: an adrenal gland disorder, kidney disease, uncontrolled high blood pressure, heart disease, coronary artery disease, circulation problems (especially with diabetes), undiagnosed vaginal bleeding,

liver disease or liver cancer, severe migraine headaches, if you also take certain hepatitis C medication, if you will have major surgery, if you smoke and are over 35, or if you have ever had a heart attack, a stroke, a blood clot, jaundice caused by pregnancy or birth control pills, or cancer of the breast, uterus/cervix, or vagina.

Taking this medicine can increase your risk of blood clots, stroke, or heart attack.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. You should not take this medicine if you smoke and are over 35 years old.

## Before taking this medicine

Taking this medicine can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have high blood pressure, diabetes, high cholesterol, or if you are overweight. Your risk of stroke or blood clot is highest during your first year of taking birth control pills. Your risk is also high when you restart birth control pills after not taking them for 4 weeks or longer.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. Your risk increases the older you are and the more you smoke. You should not take this medicine if you smoke and are over 35 years old.

**Do not use if you are pregnant.** Stop using this medicine and tell your doctor right away if you become pregnant, or if you miss 2 menstrual periods in a row. If you have recently had a baby, wait at least 4 weeks before taking this medicine.

You should not take this medicine if you have:

- · an adrenal gland disorder;
- · kidney disease;
- untreated or uncontrolled high blood pressure;
- heart disease (coronary artery disease, history of heart attack, stroke, or blood clot);
- an increased risk of having blood clots due to a heart problem or a hereditary blood disorder;
- circulation problems (especially if caused by diabetes);
- a history of hormone-related cancer, or cancer of the breast, uterus/cervix, or vagina;
- unusual vaginal bleeding that has not been checked by a doctor;
- · liver disease or liver cancer:
- severe migraine headaches (with aura, numbness, weakness, or vision changes), especially if you are older than 35;
- a history of jaundice caused by pregnancy or birth control pills;
- if you smoke and are over 35 years old; or
- if you take any hepatitis C medication containing ombitasvir/paritaprevir/ritonavir (Technivie).

Tell your doctor if you have ever had:

- heart problems, high blood pressure, or if you are prone to having blood clots;
- high levels of potassium in your blood;
- high cholesterol or triglycerides, or if you are overweight;

- · depression;
- liver or kidney disease;
- · underactive thyroid, diabetes, gallbladder disease; or
- a migraine headache.

The hormones in this medicine can pass into breast milk and may harm a nursing baby. This medicine may also slow breast milk production. Do not use if you are breast feeding.

## How should I take Zarah?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Take your first pill on the first day of your period or on the first Sunday after your period begins. You may need to use back-up birth control, such as condoms with spermicide, when you first start using this medication.

Take one pill every day, no more than 24 hours apart. When the pills run out, start a new pack the following day. **You could get pregnant if you do not take one pill daily**.

You may have breakthrough bleeding, especially during the first 3 months. Tell your doctor if this bleeding continues or is very heavy.

If you need major surgery or will be on long-term bed rest, you may need to stop using Zarah for a short time. Any doctor or surgeon who treats you should know that you are using this medicine.

Store at room temperature away from moisture and heat.

Detailed Zarah dosage information

# What happens if I miss a dose?

Follow the patient instructions provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions. Missing a pill increases your risk of becoming pregnant.

If you miss 1 active pill, take 2 pills on the day that you remember. Then take 1 pill per day for the rest of the pack.

If you miss 2 active pills in a row in Week 1 or 2, take 2 pills per day for 2 days in a row. Then take 1 pill per day for the rest of the pack. Use back-up birth control for at least 7 days following the missed pills.

If you miss 2 active pills in a row in Week 3, throw out the rest of the pack and start a new pack the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack that day.

If you miss 3 active pills in a row in Week 1, 2, or 3, throw out the rest of the pack and start a new pack on the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack that day.

If you miss 2 or more pills, you may not have a period during the month. If you miss a period for 2 months in a row, call your doctor because you might be pregnant.

If you miss a reminder pill, throw it away and keep taking 1 reminder pill per day until the pack is empty.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Overdose may cause nausea or vaginal bleeding.

# What should I avoid while taking Zarah?

Do not smoke while taking Zarah, especially if you are older than 35 years of age.

This medicine will not protect you from sexually transmitted diseases, including HIV and AIDS. Using a condom is the only way to protect yourself from these diseases.

# What other drugs will affect Zarah?

Other drugs may interact with drospirenone and ethinyl estradiol, including prescription and over-the-counter medicines, vitamins, and herbal products. Some drugs can make this medicine less effective, which may result in pregnancy. Tell your doctor about all your current medicines and any medicine you start or stop using.

<u>Zarah drug interactions</u> (more detail)

## Does Zarah interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Zarah	
+	
Enter a drug name	Add

# More about Zarah (drospirenone / ethinyl estradiol)

- Check interactions
- Compare alternatives
- Reviews (84)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: contraceptives
- En español

## **Patient resources**

· Zarah advanced reading

#### Other brands

Yaz, Yasmin, Ocella, Loryna, ... +6 more

## **Professional resources**

· Zarah prescribing information

#### Other brands

Yaz, Yasmin, Ocella, Loryna, ... +7 more

## Related treatment guides

- Acne
- Birth Control
- Premenstrual Dysphoric Disorder

## **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

#### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 10.03.

#### **DRUG STATUS**

#### **Availability**

Rx Prescription only

**Pregnancy & Lactation** 

ം Risk data available

CSA Schedule\*

N/A Not a controlled drug

## **User Reviews & Ratings**

6.2 / 10

84 Reviews

## **Images**

#### Zarah inert (WATSON 983)



# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com Advertising policy

Attribution & citations

Terms & privacy

Terms of use

**Editorial policy** 

Privacy policy

## Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







