


QDCS : Digrammatic Calculus and Error Correction

Renaud Vilmart


TD 3

1 Simplification and Decomposition in Clifford+T


Question 1. Reduce the following diagram so that the only inner spiders are “T-spiders” or spiders with 0 or π phases only connected to T-spiders:



Answer:



Question 2. Consider the following diagram:



1. Apply the “2 T to 2 terms” decomposition from the course
2. Put each of the 2 terms in Clifford normal form.

2 Consecutive Measurements

Question 1. Define the “upside-down” trace as: $\overline{\top} := \bigcap \overline{\perp}$. This is hence a quantum state. Compute its interpretation. Explain why it can be understood as complete noise.


Question 2. Compute the interpretation of the diagram: , and show it is equivalent to tracing out the

input state and outputting complete noise.

Question 3. Show the equality between the diagrams of the previous question, but graphically this time, in two different ways:

1. directly using the axioms of the partial trace
2. by going through the CPM construction

Answer: Everything was done in class, except Q.3.2. Using the CPM construction, we get




The last diagram being the image of the partial trace followed by noise, we can get the equality that way.

3 Quantum Teleportation

The problem is the following: Two parties, Alice and Bob, can communicate, but only classically, i.e. they can exchange classical bits, but not qubits. Alice has a qubit in an unknown state $|\psi\rangle$ which she wishes to send to Bob. If no further assumption is made, the problem can be shown to be impossible to solve: Alice cannot send her qubit to Bob through a classical channel. Measurement would destroy the state and only give him (very) partial information.


The problem is however shown to be solvable if Alice and Bob initially share an entangled pair of qubits (Alice has the first one, and Bob the second one). In the following, we can imagine that this state is prepared by a third party called Eve, who splits the pair between Alice and Bob before the protocol really starts. Then, when Alice wants to send her qubit to Bob, she entangles it with her half of the pair (using a CNot), then measures her two qubits in the appropriate basis. Doing this disturbs the system but the disturbance depends on the measurement results. In this case, the errors introduced by Alice’s measurements can be corrected by Bob, provided he has access to the measurement results (which are classical). Alice hence sends them to Bob. After correction, the qubit is flawlessly transmitted.

The precise protocol (where measurements use the observable Z) is given in circuit form as follows:



Question 1. Translate the quantum teleportation protocol in ZX using variables for the measurements. Show that the whole diagram is equivalent to the identity (or more exactly to a direct 1-qubit communication from Alice to Bob).

Answer:



Question 2. Translate the quantum teleportation protocol in ZX using partial traces for the measurements. Show that the whole diagram is equivalent to the identity.

Answer:

