

DMRG Study of the $S \geq 1$ quantum Heisenberg Antiferromagnet on a Kagome-like lattice without loops

R. Zach Lamberty, Hitesh Changlani, and C. L. Henley
Cornell University, Ithaca, New York, 14853
(Dated: November 7, 2012)

The Kagome quantum Heisenberg antiferromagnet, not only for spin $S = 1/2$ but for $S = 1$ and perhaps $S = 3/2$, is a prime candidate to realize a quantum spin liquid or valence bond crystal state. However, theoretical and computational studies for $S > 1/2$ are difficult and few. We consider the quantum Heisenberg antiferromagnet for $S \geq 1$ on the Husimi cactus, a graph of corner sharing triangles each of whose centers is a vertex of a Bethe lattice, using a DMRG procedure tailored for tree graphs [1]. Since the geometry is like the Kagome lattice locally, properties dominated by nearest-neighbor spin correlations should be captured by the same interactions on the Husimi cactus; on the other hand, since the cactus lacks loops, properties dependent on them cannot be captured. The cactus antiferromagnet is known to have a disordered valence bond state at $S = 1/2$ [2,3] but a three-sublattice coplanar ordered state in the large S limit [4]. Thus, our focus is the possible transition(s) that must occur with increasing S . Using this approach, we also investigate the phase diagram of the $S = 1$ quantum XXZ model with on-site anisotropy, which we expect to have a variety of phases similar to the kagome case [5]. This work is supported by the National Science Foundation through a Graduate Research Fellowship to R. Zach Lamberty, as well as grant DMR-0552461.

- [1] H. J. Changlani, S. Ghosh, C. L. Henley, A. M. Läuchli, arXiv:1208.1773v1 (2012).
- [2] V. Elser and C. Zeng, Phys. Rev. B **48**, 13647 ((1993).
- [3] P. Chandra and B. Doucot, J. Phys. A **27**, 1541 (1994).
- [4] B. Doucot and P. Simon, J. Phys. A **31**, 5855 (1998).
- [5] S. V. Isakov and Y. B. Kim, Phys. Rev. B **79**, 094408 (2009).