

# Вычислительные схемы для Higher Order модели

Фареник А.

14 марта 2023 г.

# Дискретная сетка

В пространстве  $\{x, y, z\} \in \mathbb{R}^3$  строится сетка

Перед введением схемы по  $z$  производится координатное отображение.

$$\xi|_{x,y} = \frac{z|_{x,y}}{s(x,y) - b(x,y)}$$

- равномерная сетка по  $x$ :  $\{x_i\} : x_i = i \cdot \Delta x$
- равномерная сетка по  $y$ :  $\{y_j\} : y_j = j \cdot \Delta y$
- неравномерная сетка по  $\xi : \{\xi_k\}$
- некоторая сетка по времени  $t$ :  $\{t_l\}$

Указанные индексные обозначения закрепим за координатами в пространстве и времени, т.е.  $i$  будет нумеровать узлы по координате  $x$ ,  $j$  – по координате  $y$ ,  $k$  – по координате  $z$  и  $l$  – по времени  $t$ .

Также для получения схем, включающих  $z$  нужны следующие соотношения.

$$\begin{aligned}\Delta \xi_{k-\frac{1}{2}} &= \xi_k - \xi_{k-1} \\ \Delta \xi_{k+\frac{1}{2}} &= \xi_{k+1} - \xi_k \\ \Delta \xi_k &= \frac{\Delta \xi_{k+\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}}{2} = \frac{\xi_{k+1} - \xi_{k-1}}{2} \\ f_{k-\frac{1}{2}} &= \frac{f_k + f_{k-1}}{2} \\ f_{k+\frac{1}{2}} &= \frac{f_{k+1} + f_k}{2} \\ f_k &= \frac{\Delta \xi_{k-\frac{1}{2}}}{2\Delta \xi_k} f_{k+\frac{1}{2}} + \frac{\Delta \xi_{k+\frac{1}{2}}}{2\Delta \xi_k} f_{k-\frac{1}{2}} \\ \left( \frac{\partial f}{\partial \xi} \right)_k &= \frac{\Delta \xi_{k-\frac{1}{2}}}{\Delta \xi_k \Delta \xi_{k+\frac{1}{2}}} f_{k+\frac{1}{2}} - \frac{\Delta \xi_{k+\frac{1}{2}}}{\Delta \xi_k \Delta \xi_{k-\frac{1}{2}}} f_{k-\frac{1}{2}} + 2 \frac{\Delta \xi_{k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}}{\Delta \xi_{k+\frac{1}{2}} \Delta \xi_{k-\frac{1}{2}}} f_k\end{aligned}$$

# Преобразование координат

Для моделирования ледников обычно производят преобразование по вертикальной координате, скалирую область ледника на единичную высоту. Для этого используется координатное отображение

$$(x, y, z, t) \mapsto (x', y', \xi, t)$$

Примером такого отображения может быть

$$\xi = \frac{z - b}{s - b} = \frac{z - b}{H}$$

Тогда нижняя граница ледника будет всегда находится на плоскости  $\xi = 0$ , а верхняя граница на плоскости  $\xi = 1$ .

Рассмотрим, как происходит преобразование некоторой достаточно гладкой функции  $f = f(x, y, z, t)$  при таком координатном отображении.

С небольшой погрешностью можно утверждать, что

$$\begin{cases} x \approx x' \\ y \approx y' \\ t \approx t' \end{cases}$$

Перед тем, как выводить вид функций в новых координатах приведем вспомогательные производные.

$$\begin{aligned} a_x = \frac{\partial \xi}{\partial x} &= \frac{\partial}{\partial x} \left( \frac{z - b}{H} \right) = \frac{1}{H} \frac{\partial}{\partial x} (z - b) + (z - b) \frac{\partial}{\partial x} \left( \frac{1}{H} \right) = -\frac{1}{H} \frac{\partial b}{\partial x} + (z - b) \cdot \frac{-1}{H} \frac{\partial H}{\partial x} = \\ &= \frac{-1}{H} \left( \frac{\partial b}{\partial x} + \frac{z - b}{H} \frac{\partial H}{\partial x} \right) = -\frac{1}{H} \left( \frac{\partial b}{\partial x} + \xi \frac{\partial H}{\partial x} \right) \approx -\frac{1}{H} \left( \frac{\partial b}{\partial x'} + \xi \frac{\partial H}{\partial x'} \right) \end{aligned}$$

Аналогично

$$a_y = \frac{\partial \xi}{\partial y} = -\frac{1}{H} \left( \frac{\partial b}{\partial y} + \xi \frac{\partial H}{\partial y} \right) \approx -\frac{1}{H} \left( \frac{\partial b}{\partial y'} + \xi \frac{\partial H}{\partial y'} \right)$$

$$a_z = \frac{\partial \xi}{\partial z} = \frac{\partial}{\partial z} \left( \frac{z - b}{H} \right) = \frac{1}{H}$$

$$\begin{aligned} b_x = \frac{\partial a_x}{\partial x} &= \frac{\partial}{\partial x} \left[ -\frac{1}{H} \left( \frac{\partial b}{\partial x} + \xi \frac{\partial H}{\partial x} \right) \right] = \frac{\partial}{\partial x} \left( -\frac{1}{H} \right) \left( \frac{\partial b}{\partial x} + \xi \frac{\partial H}{\partial x} \right) + \left( -\frac{1}{H} \right) \frac{\partial}{\partial x} \left( \frac{\partial b}{\partial x} + \xi \frac{\partial H}{\partial x} \right) = \\ &= \frac{1}{H^2} \frac{\partial H}{\partial x} \left( \frac{\partial b}{\partial x} + \xi \frac{\partial H}{\partial x} \right) - \frac{1}{H} \left( \frac{\partial^2 b}{\partial x^2} + \frac{\partial \xi}{\partial x} \frac{\partial H}{\partial x} + \xi \frac{\partial^2 H}{\partial x^2} \right) = \\ &= \frac{1}{H^2} \frac{\partial H}{\partial x} \left( \frac{\partial b}{\partial x} + \xi \frac{\partial H}{\partial x} \right) - \frac{1}{H} \left( \frac{\partial^2 b}{\partial x^2} + a_x \frac{\partial b}{\partial x} + \xi \frac{\partial^2 H}{\partial x^2} \right) = \\ &= -\frac{1}{H} \frac{\partial H}{\partial x} a_x - \frac{1}{H} a_x \frac{1}{H} - \frac{1}{H} \left( \frac{\partial^2 b}{\partial x^2} + \xi \frac{\partial^2 H}{\partial x^2} \right) = -\frac{1}{H} \left( \frac{\partial^2 b}{\partial x^2} + \xi \frac{\partial^2 H}{\partial x^2} + 2a_x \frac{\partial H}{\partial x} \right) \approx \\ &\approx -\frac{1}{H} \left( \frac{\partial^2 b}{\partial x'^2} + \xi \frac{\partial^2 H}{\partial x'^2} + 2a_x \frac{\partial H}{\partial x'} \right) \end{aligned}$$

$$b_y = \frac{\partial a_y}{\partial y} = -\frac{1}{H} \left( \frac{\partial^2 b}{\partial y^2} + \xi \frac{\partial^2 H}{\partial y^2} + 2a_y \frac{\partial H}{\partial y} \right) \approx -\frac{1}{H} \left( \frac{\partial^2 b}{\partial y'^2} + \xi \frac{\partial^2 H}{\partial y'^2} + 2a_y \frac{\partial H}{\partial y'} \right)$$

$$b_z = \frac{\partial a_z}{\partial z} = \frac{\partial}{\partial z} \left( \frac{1}{H} \right) = 0$$

$$\begin{aligned}
c_{xy} &= \frac{\partial a_y}{\partial x} = c_{yx} = \frac{\partial a_x}{\partial y} = \frac{\partial}{\partial x} \left[ -\frac{1}{H} \left( \frac{\partial b}{\partial y} + \xi \frac{\partial H}{\partial y} \right) \right] = \\
&= -\frac{\partial}{\partial x} \left( \frac{1}{H} \right) \left( \frac{\partial b}{\partial y} + \xi \frac{\partial H}{\partial y} \right) - \frac{1}{H} \frac{\partial}{\partial x} \left( \frac{\partial b}{\partial y} + \xi \frac{\partial H}{\partial y} \right) = \\
&= \frac{1}{H^2} \frac{\partial H}{\partial x} \left( \frac{\partial b}{\partial y} + \xi \frac{\partial H}{\partial y} \right) - \frac{1}{H} \left( \frac{\partial^2 b}{\partial x \partial y} + \frac{\partial \xi}{\partial x} \frac{\partial H}{\partial y} + \xi \frac{\partial^2 H}{\partial x \partial y} \right) = \\
&= -\frac{1}{H} a_y \frac{\partial H}{\partial x} - \frac{1}{H} \left( \frac{\partial^2 b}{\partial x \partial y} + a_x \frac{\partial H}{\partial y} + \xi \frac{\partial^2 H}{\partial x \partial y} \right) = -\frac{1}{H} \left( \frac{\partial^2 b}{\partial x \partial y} + a_x \frac{\partial H}{\partial y} + a_y \frac{\partial H}{\partial x} + \xi \frac{\partial^2 H}{\partial x \partial y} \right) \approx \\
&\approx -\frac{1}{H} \left( \frac{\partial^2 b}{\partial x' \partial y'} + a_x \frac{\partial H}{\partial y'} + a_y \frac{\partial H}{\partial x'} + \xi \frac{\partial^2 H}{\partial x' \partial y'} \right) \\
c_{xz} &= \frac{\partial a_z}{\partial x} = c_{zx} = \frac{\partial a_x}{\partial z} = \frac{\partial}{\partial x} \left( \frac{1}{H} \right) = -\frac{1}{H^2} \frac{\partial H}{\partial x} \approx -\frac{1}{H^2} \frac{\partial H}{\partial x'} \\
c_{yz} &= \frac{\partial a_z}{\partial y} = c_{zy} = \frac{\partial a_y}{\partial z} = -\frac{1}{H^2} \frac{\partial H}{\partial y} \approx -\frac{1}{H^2} \frac{\partial H}{\partial y'}
\end{aligned}$$

Перейдем к первым производным функции  $f$

$$\begin{aligned}
\frac{\partial f}{\partial x} &= \frac{\partial f}{\partial x'} \frac{\partial x'}{\partial x} + \frac{\partial f}{\partial y'} \frac{\partial y'}{\partial x} + \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial f}{\partial t'} \frac{\partial t'}{\partial x} \approx \frac{\partial f}{\partial x'} + a_x \frac{\partial f}{\partial \xi} \\
\frac{\partial f}{\partial y} &\approx \frac{\partial f}{\partial y'} + a_y \frac{\partial f}{\partial \xi} \\
\frac{\partial f}{\partial z} &= \frac{\partial f}{\partial x'} \frac{\partial x'}{\partial z} + \frac{\partial f}{\partial y'} \frac{\partial y'}{\partial z} + \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial z} + \frac{\partial f}{\partial t'} \frac{\partial t'}{\partial z} \approx a_z \frac{\partial f}{\partial \xi}
\end{aligned}$$

Далее нужно рассмотреть вторые производные

$$\begin{aligned}
\frac{\partial^2 f}{\partial x^2} &= \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x'} + a_x \frac{\partial f}{\partial \xi} \right) = \frac{\partial}{\partial x} \frac{\partial f}{\partial x'} + \frac{\partial}{\partial \xi} \frac{\partial f}{\partial x'} \frac{\partial \xi}{\partial x} + \frac{\partial a_x}{\partial x} \frac{\partial f}{\partial \xi} + a_x \frac{\partial}{\partial x} \frac{\partial f}{\partial \xi} + a_x \frac{\partial}{\partial \xi} \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial x} \approx \\
&\approx \frac{\partial^2 f}{\partial x'^2} + 2a_x \frac{\partial^2 f}{\partial x' \partial \xi} + a_x^2 \frac{\partial^2 f}{\partial \xi^2} + b_x \frac{\partial f}{\partial \xi} \\
\frac{\partial^2 f}{\partial y^2} &\approx \frac{\partial^2 f}{\partial y'^2} + 2a_y \frac{\partial^2 f}{\partial y' \partial \xi} + a_y^2 \frac{\partial^2 f}{\partial \xi^2} + b_y \frac{\partial f}{\partial \xi} \\
\frac{\partial^2 f}{\partial z^2} &= \frac{\partial}{\partial z} \left( a_z \frac{\partial f}{\partial \xi} \right) = a_z \frac{\partial}{\partial z} \frac{\partial f}{\partial \xi} \approx a_z^2 \frac{\partial^2 f}{\partial \xi^2}
\end{aligned}$$

Включая смешанные частные производные

$$\begin{aligned}
\frac{\partial^2 f}{\partial x \partial y} &= \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y'} + a_y \frac{\partial f}{\partial \xi} \right) = \frac{\partial}{\partial x} \frac{\partial f}{\partial y'} + \frac{\partial}{\partial \xi} \frac{\partial f}{\partial y'} \frac{\partial \xi}{\partial x} + \frac{\partial a_y}{\partial x} \frac{\partial f}{\partial \xi} + a_y \frac{\partial}{\partial x} \frac{\partial f}{\partial \xi} = \\
&= \frac{\partial}{\partial x'} \frac{\partial f}{\partial y'} \frac{\partial x'}{\partial x} + a_x \frac{\partial^2 f}{\partial y' \partial \xi} + c_{xy} \frac{\partial f}{\partial \xi} + a_y \frac{\partial}{\partial x'} \frac{\partial f}{\partial \xi} \frac{\partial x'}{\partial x} + a_y \frac{\partial}{\partial \xi} \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial x} \approx \\
&\approx \frac{\partial^2 f}{\partial x' \partial y'} + a_x \frac{\partial^2 f}{\partial y' \partial \xi} + a_y \frac{\partial^2 f}{\partial x' \partial \xi} + a_x a_y \frac{\partial^2 f}{\partial \xi^2} + c_{xy} \frac{\partial f}{\partial \xi} \\
\frac{\partial^2 f}{\partial x \partial z} &= \frac{\partial^2 f}{\partial z \partial x} = \frac{\partial}{\partial x} \frac{\partial f}{\partial z} = \frac{\partial}{\partial x} \left( a_z \frac{\partial f}{\partial \xi} \right) = \frac{\partial a_z}{\partial x} \frac{\partial f}{\partial \xi} + a_z \frac{\partial}{\partial x} \frac{\partial f}{\partial \xi} = \\
&= c_{xz} \frac{\partial f}{\partial \xi} + a_z \left( \frac{\partial}{\partial x'} \frac{\partial f}{\partial \xi} \frac{\partial x'}{\partial x} + \frac{\partial}{\partial \xi} \frac{\partial f}{\partial \xi} \frac{\partial \xi}{\partial x} \right) \approx a_z \frac{\partial^2 f}{\partial x' \partial \xi} + a_x a_z \frac{\partial^2 f}{\partial \xi^2} + c_{xz} \frac{\partial f}{\partial \xi}
\end{aligned}$$

$$\frac{\partial^2 f}{\partial y \partial z} = \frac{\partial^2 f}{\partial z \partial y} \approx a_z \frac{\partial^2 f}{\partial y' \partial \xi} + a_y a_z \frac{\partial^2 f}{\partial \xi^2} + c_{yz} \frac{\partial f}{\partial \xi}$$

# Схемы основных операторов

В данном разделе приведены дискретные схемы основных дифференциальных операторов.

## 2D операторы

$$\frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial x} \right)_{i,j} = \frac{1}{\Delta x^2} \left\{ f_{i-1,j} \left[ g_{i-\frac{1}{2},j} \right] + f_{i,j} \left[ -g_{i-\frac{1}{2},j} - g_{i+\frac{1}{2},j} \right] + f_{i+1,j} \left[ g_{i+\frac{1}{2},j} \right] \right\}$$

$$\begin{aligned} \frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial y} \right)_{i,j} = & \frac{1}{4\Delta x \Delta y} \left\{ f_{i-1,j-1} \left[ g_{i-\frac{1}{2},j-\frac{1}{2}} \right] + f_{i-1,j} \left[ -g_{i-\frac{1}{2},j-\frac{1}{2}} + g_{i-\frac{1}{2},j+\frac{1}{2}} \right] + \right. \\ & + f_{i-1,j+1} \left[ -g_{i-\frac{1}{2},j+\frac{1}{2}} \right] + f_{i,j-1} \left[ g_{i-\frac{1}{2},j-\frac{1}{2}} - g_{i+\frac{1}{2},j-\frac{1}{2}} \right] + \\ & + f_{i,j} \left[ -g_{i-\frac{1}{2},j-\frac{1}{2}} + g_{i-\frac{1}{2},j+\frac{1}{2}} + g_{i+\frac{1}{2},j-\frac{1}{2}} - g_{i+\frac{1}{2},j+\frac{1}{2}} \right] + \\ & + f_{i,j+1} \left[ -g_{i-\frac{1}{2},j+\frac{1}{2}} + g_{i+\frac{1}{2},j+\frac{1}{2}} \right] + f_{i+1,j-1} \left[ -g_{i+\frac{1}{2},j-\frac{1}{2}} \right] + \\ & \left. + f_{i+1,j} \left[ g_{i+\frac{1}{2},j-\frac{1}{2}} - g_{i+\frac{1}{2},j+\frac{1}{2}} \right] + f_{i+1,j+1} \left[ g_{i+\frac{1}{2},j+\frac{1}{2}} \right] \right\} \end{aligned}$$

$$\begin{aligned} \frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial x} \right)_{i,j} = & \frac{1}{4\Delta x \Delta y} \left\{ f_{i-1,j-1} \left[ g_{i-\frac{1}{2},j-\frac{1}{2}} \right] + f_{i-1,j} \left[ g_{i-\frac{1}{2},j-\frac{1}{2}} - g_{i-\frac{1}{2},j+\frac{1}{2}} \right] + \right. \\ & + f_{i-1,j+1} \left[ -g_{i-\frac{1}{2},j+\frac{1}{2}} \right] + f_{i,j-1} \left[ -g_{i-\frac{1}{2},j-\frac{1}{2}} + g_{i+\frac{1}{2},j-\frac{1}{2}} \right] + \\ & + f_{i,j} \left[ -g_{i-\frac{1}{2},j-\frac{1}{2}} + g_{i-\frac{1}{2},j+\frac{1}{2}} + g_{i+\frac{1}{2},j-\frac{1}{2}} - g_{i+\frac{1}{2},j+\frac{1}{2}} \right] + \\ & + f_{i,j+1} \left[ g_{i-\frac{1}{2},j+\frac{1}{2}} - g_{i+\frac{1}{2},j+\frac{1}{2}} \right] + f_{i+1,j-1} \left[ -g_{i+\frac{1}{2},j-\frac{1}{2}} \right] + \\ & \left. + f_{i+1,j} \left[ -g_{i+\frac{1}{2},j-\frac{1}{2}} + g_{i+\frac{1}{2},j+\frac{1}{2}} \right] + f_{i+1,j+1} \left[ g_{i+\frac{1}{2},j+\frac{1}{2}} \right] \right\} \end{aligned}$$

$$\frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial y} \right)_{i,j} = \frac{1}{\Delta y^2} \left\{ f_{i,j-1} \left[ g_{i,j-\frac{1}{2}} \right] + f_{i,j} \left[ -g_{i,j-\frac{1}{2}} - g_{i,j+\frac{1}{2}} \right] + f_{i,j+1} \left[ g_{i,j+\frac{1}{2}} \right] \right\}$$

## 3D операторы

$$\frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial x} \right)_{i,j,k} = \frac{1}{\Delta x^2} \left\{ f_{i-1,j,k} \left[ g_{i-\frac{1}{2},j,k} \right] + f_{i,j,k} \left[ -g_{i-\frac{1}{2},j,k} - g_{i+\frac{1}{2},j,k} \right] + f_{i+1,j,k} \left[ g_{i+\frac{1}{2},j,k} \right] \right\}$$

$$\begin{aligned} \frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial y} \right)_{i,j,k} = & \frac{1}{4\Delta x \Delta y} \left\{ f_{i-1,j-1,k} \left[ g_{i-\frac{1}{2},j-\frac{1}{2},k} \right] + f_{i-1,j,k} \left[ -g_{i-\frac{1}{2},j-\frac{1}{2},k} + g_{i-\frac{1}{2},j+\frac{1}{2},k} \right] + \right. \\ & + f_{i-1,j+1,k} \left[ -g_{i-\frac{1}{2},j+\frac{1}{2},k} \right] + f_{i,j-1,k} \left[ g_{i-\frac{1}{2},j-\frac{1}{2},k} - g_{i+\frac{1}{2},j-\frac{1}{2},k} \right] + \\ & + f_{i,j,k} \left\{ -g_{i-\frac{1}{2},j-\frac{1}{2},k} + g_{i-\frac{1}{2},j+\frac{1}{2},k} + g_{i+\frac{1}{2},j-\frac{1}{2},k} - g_{i+\frac{1}{2},j+\frac{1}{2},k} \right\} + \\ & + f_{i,j+1,k} \left[ -g_{i-\frac{1}{2},j+\frac{1}{2},k} + g_{i+\frac{1}{2},j+\frac{1}{2},k} \right] + f_{i+1,j-1,k} \left[ -g_{i+\frac{1}{2},j-\frac{1}{2},k} \right] + \\ & \left. + f_{i+1,j,k} \left[ g_{i+\frac{1}{2},j-\frac{1}{2},k} - g_{i+\frac{1}{2},j+\frac{1}{2},k} \right] + f_{i+1,j+1,k} \left[ g_{i+\frac{1}{2},j+\frac{1}{2},k} \right] \right\} \end{aligned}$$

$$\begin{aligned} \frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial \xi} \right)_{i,j,k} &= \frac{1}{4\Delta x \Delta \xi_k \Delta \xi_{k-\frac{1}{2}} \Delta \xi_{k+\frac{1}{2}}} \times \\ &\times \left\{ f_{i-1,j,k-1} \left[ \Delta \xi_{k+\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k-\frac{1}{2}} \right] + f_{i-1,j,k} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k-\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k+\frac{1}{2}} \right] + \right. \\ &+ f_{i-1,j,k+1} \left[ -\Delta \xi_{k-\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k+\frac{1}{2}} \right] + f_{i,j,k-1} \left[ \Delta \xi_{k+\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k-\frac{1}{2}} - \Delta \xi_{k+\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k-\frac{1}{2}} \right] + \\ &+ f_{i,j,k} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k-\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k+\frac{1}{2}} + \Delta \xi_{k+\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k-\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k+\frac{1}{2}} \right] + \\ &+ f_{i,j,k+1} \left[ -\Delta \xi_{k-\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k+\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k+\frac{1}{2}} \right] + f_{i+1,j,k-1} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k-\frac{1}{2}} \right] + \\ &\left. + f_{i+1,j,k} \left[ \Delta \xi_{k+\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k-\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k+\frac{1}{2}} \right] + f_{i+1,j,k+1} \left[ \Delta \xi_{k-\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k+\frac{1}{2}} \right] \right\} \end{aligned}$$

$$\begin{aligned} \frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial x} \right)_{i,j,k} &= \frac{1}{4\Delta x \Delta y} \left\{ f_{i-1,j-1,k} \left[ g_{i-\frac{1}{2},j-\frac{1}{2},k} \right] + f_{i-1,j,k} \left[ g_{i-\frac{1}{2},j-\frac{1}{2},k} - g_{i-\frac{1}{2},j+\frac{1}{2},k} \right] + \right. \\ &+ f_{i-1,j+1,k} \left[ -g_{i-\frac{1}{2},j+\frac{1}{2},k} \right] + f_{i,j-1,k} \left[ -g_{i-\frac{1}{2},j-\frac{1}{2},k} + g_{i+\frac{1}{2},j-\frac{1}{2},k} \right] + \\ &+ f_{i,j,k} \left\{ -g_{i-\frac{1}{2},j-\frac{1}{2},k} + g_{i-\frac{1}{2},j+\frac{1}{2},k} + g_{i+\frac{1}{2},j-\frac{1}{2},k} - g_{i+\frac{1}{2},j+\frac{1}{2},k} \right\} + \\ &+ f_{i,j+1,k} \left[ g_{i-\frac{1}{2},j+\frac{1}{2},k} - g_{i+\frac{1}{2},j+\frac{1}{2},k} \right] + f_{i+1,j-1,k} \left[ -g_{i+\frac{1}{2},j-\frac{1}{2},k} \right] + \\ &\left. + f_{i+1,j,k} \left[ -g_{i+\frac{1}{2},j-\frac{1}{2},k} + g_{i+\frac{1}{2},j+\frac{1}{2},k} \right] + f_{i+1,j+1,k} \left[ g_{i+\frac{1}{2},j+\frac{1}{2},k} \right] \right\} \end{aligned}$$

$$\frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial y} \right)_{i,j,k} = \frac{1}{\Delta y^2} \left\{ f_{i,j-1,k} \left[ g_{i,j-\frac{1}{2},k} \right] + f_{i,j,k} \left[ -g_{i,j-\frac{1}{2},k} - g_{i,j+\frac{1}{2},k} \right] + f_{i,j+1,k} \left[ g_{i,j+\frac{1}{2},k} \right] \right\}$$

$$\begin{aligned} \frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial \xi} \right)_{i,j,k} &= \frac{1}{4\Delta y \Delta \xi_k \Delta \xi_{k-\frac{1}{2}} \Delta \xi_{k+\frac{1}{2}}} \times \\ &\times \left\{ f_{i,j-1,k-1} \left[ \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k-\frac{1}{2}} \right] + f_{i,j-1,k} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k-\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k+\frac{1}{2}} \right] + \right. \\ &+ f_{i,j-1,k+1} \left[ -\Delta \xi_{k-\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k+\frac{1}{2}} \right] + f_{i,j,k-1} \left[ \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k-\frac{1}{2}} - \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k-\frac{1}{2}} \right] + \\ &+ f_{i,j,k} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k-\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k+\frac{1}{2}} + \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k-\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k+\frac{1}{2}} \right] + \\ &+ f_{i,j,k+1} \left[ -\Delta \xi_{k-\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k+\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k+\frac{1}{2}} \right] + f_{i,j+1,k-1} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k-\frac{1}{2}} \right] + \\ &\left. + f_{i,j+1,k} \left[ \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k-\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k+\frac{1}{2}} \right] + f_{i,j+1,k+1} \left[ \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k+\frac{1}{2}} \right] \right\} \end{aligned}$$

$$\begin{aligned} \frac{\partial}{\partial \xi} \left( g \frac{\partial f}{\partial x} \right)_{i,j,k} &= \frac{1}{4\Delta x \Delta \xi_k \Delta \xi_{k-\frac{1}{2}} \Delta \xi_{k+\frac{1}{2}}} \times \left\{ f_{i-1,j,k-1} \left[ \Delta \xi_{k+\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k-\frac{1}{2}} \right] + \right. \\ &+ f_{i-1,j,k} \left[ -4\Delta \xi_k \left( \Delta \xi_{k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}} \right) g_{i-\frac{1}{2},j,k} + \Delta \xi_{k+\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k-\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k+\frac{1}{2}} \right] + \\ &+ f_{i-1,j,k+1} \left[ -\Delta \xi_{k-\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k+\frac{1}{2}} \right] + f_{i,j,k-1} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k-\frac{1}{2}} + \Delta \xi_{k+\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k-\frac{1}{2}} \right] + \\ &+ f_{i,j,k} \left[ 4\Delta \xi_k \left( \Delta \xi_{k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}} \right) \left( g_{i-\frac{1}{2},j,k} - g_{i+\frac{1}{2},j,k} \right) - \right. \\ &\quad \left. - \Delta \xi_{k+\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k-\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k+\frac{1}{2}} + \Delta \xi_{k+\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k-\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k+\frac{1}{2}} \right] + \\ &+ f_{i,j,k+1} \left[ \Delta \xi_{k-\frac{1}{2}}^2 g_{i-\frac{1}{2},j,k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k+\frac{1}{2}} \right] + f_{i+1,j,k-1} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k-\frac{1}{2}} \right] + \\ &+ f_{i+1,j,k} \left[ 4\Delta \xi_k \left( \Delta \xi_{k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}} \right) g_{i+\frac{1}{2},j,k} - \Delta \xi_{k+\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k-\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k+\frac{1}{2}} \right] + \\ &\left. + f_{i+1,j,k+1} \left[ \Delta \xi_{k-\frac{1}{2}}^2 g_{i+\frac{1}{2},j,k+\frac{1}{2}} \right] \right\} \end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial \xi} \left( g \frac{\partial f}{\partial y} \right)_{i,j,k} = & \frac{1}{4\Delta y \Delta \xi_k \Delta \xi_{k-\frac{1}{2}} \Delta \xi_{k+\frac{1}{2}}} \times \left\{ f_{i,j-1,k-1} \left[ \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k-\frac{1}{2}} \right] + \right. \\
& + f_{i,j-1,k} \left[ -4\Delta \xi_k \left( \Delta \xi_{k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}} \right) g_{i,j-\frac{1}{2},k} + \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k-\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k+\frac{1}{2}} \right] + \\
& + f_{i,j-1,k+1} \left[ -\Delta \xi_{k-\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k+\frac{1}{2}} \right] + f_{i,j,k-1} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k-\frac{1}{2}} + \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k-\frac{1}{2}} \right] + \\
& + f_{i,j,k} \left[ 4\Delta \xi_k \left( \Delta \xi_{k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}} \right) \left( g_{i,j-\frac{1}{2},k} - g_{i,j+\frac{1}{2},k} \right) - \right. \\
& \quad \left. - \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k-\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k+\frac{1}{2}} + \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k-\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k+\frac{1}{2}} \right] + \\
& + f_{i,j,k+1} \left[ \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j-\frac{1}{2},k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k+\frac{1}{2}} \right] + f_{i,j+1,k-1} \left[ -\Delta \xi_{k+\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k-\frac{1}{2}} \right] + \\
& + f_{i,j+1,k} \left[ 4\Delta \xi_k \left( \Delta \xi_{k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}} \right) g_{i,j+\frac{1}{2},k} - \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k-\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k+\frac{1}{2}} \right] + \\
& \left. + f_{i,j+1,k+1} \left[ \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j+\frac{1}{2},k+\frac{1}{2}} \right] \right\}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial \xi} \left( g \frac{\partial f}{\partial \xi} \right)_{i,j,k} = & \frac{1}{\Delta \xi_k^2 \Delta \xi_{k-\frac{1}{2}} \Delta \xi_{k+\frac{1}{2}}} \left\{ f_{i,j,k-1} \left[ \Delta \xi_{k+\frac{1}{2}}^2 g_{i,j,k-\frac{1}{2}} + \frac{\Delta \xi_{k+\frac{1}{2}} \left( \Delta \xi_{k-\frac{1}{2}} - \Delta \xi_{k+\frac{1}{2}} \right)}{2} g_{i,j,k+\frac{1}{2}} \right] + \right. \\
& + f_{i,j,k} \left[ \Delta \xi_k \left( \left( -2\Delta \xi_{k-\frac{1}{2}} + \Delta \xi_{k+\frac{1}{2}} \right) g_{i,j,k-\frac{1}{2}} + \left( -2\Delta \xi_{k+\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}} \right) g_{i,j,k+\frac{1}{2}} \right) \right] + \\
& \left. + f_{i,j,k+1} \left[ \frac{\Delta \xi_{k-\frac{1}{2}} \left( \Delta \xi_{k+\frac{1}{2}} - \Delta \xi_{k-\frac{1}{2}} \right)}{2} g_{i,j,k-\frac{1}{2}} + \Delta \xi_{k-\frac{1}{2}}^2 g_{i,j,k+\frac{1}{2}} \right] \right\}
\end{aligned}$$

Для удобства вывода вычислительных схем и повышения читаемости переобозначим коэффициенты при узловых значениях функции  $f$ .

2D:

$$\frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial x} \right) = f_{i-1,j} [\alpha_x(-1)] + f_{i,j} [\alpha_x(0)] + f_{i+1,j} [\alpha_x(1)]$$

$$\begin{aligned}
\frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial y} \right) = & f_{i-1,j-1} [\beta_{x,y}(-1, -1)] + f_{i-1,j} [\beta_{x,y}(-1, 0)] + f_{i-1,j+1} [\beta_{x,y}(-1, 1)] + \\
& + f_{i,j-1} [\beta_{x,y}(0, -1)] + f_{i,j} [\beta_{x,y}(0, 0)] + f_{i,j+1} [\beta_{x,y}(0, 1)] + \\
& + f_{i+1,j-1} [\beta_{x,y}(1, -1)] + f_{i+1,j} [\beta_{x,y}(1, 0)] + f_{i+1,j+1} [\beta_{x,y}(1, 1)]
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial x} \right) = & f_{i-1,j-1} [\beta_{y,x}(-1, -1)] + f_{i-1,j} [\beta_{y,x}(-1, 0)] + f_{i-1,j+1} [\beta_{y,x}(-1, 1)] + \\
& + f_{i,j-1} [\beta_{y,x}(0, -1)] + f_{i,j} [\beta_{y,x}(0, 0)] + f_{i,j+1} [\beta_{y,x}(0, 1)] + \\
& + f_{i+1,j-1} [\beta_{y,x}(1, -1)] + f_{i+1,j} [\beta_{y,x}(1, 0)] + f_{i+1,j+1} [\beta_{y,x}(1, 1)]
\end{aligned}$$

$$\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right) = f_{i,j-1} [\alpha_y(-1)] + f_{i,j} [\alpha_y(0)] + f_{i,j+1} [\alpha_y(1)]$$

Для введенных коэффициентов можно заметить некоторые свойства

$$\begin{cases} \beta_{x,y}(p, q) = -\beta_{y,x}(p, q), & |p| \neq |q| \\ \beta_{x,y}(p, q) = \beta_{y,x}(p, q), & |p| = |q| \end{cases}$$



Таким образом можно кратко записать схему для суммы.

$$\left[ \frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial y} \right) + \frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial x} \right) \right] = f_{i-1,j-1} [2\beta_{x,y}(-1, -1)] + f_{i-1,j+1} [2\beta_{x,y}(-1, 1)] + \\ + f_{i,f} [2\beta_{x,y}(0, 0)] + f_{i+1,j-1} [2\beta_{x,y}(1, -1)] + f_{i+1,j+1} [2\beta_{x,y}(1, 1)]$$

3D:

$$\frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial x} \right) = f_{i-1,j,k} [\alpha_x(-1)] + f_{i,j,k} [\alpha_x(0)] + f_{i+1,j,k} [\alpha_x(1)]$$

$$\frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial y} \right) = f_{i-1,j-1,k} [\beta_{x,y}(-1, -1)] + f_{i-1,j,k} [\beta_{x,y}(-1, 0)] + f_{i-1,j+1,k} [\beta_{x,y}(-1, 1)] + \\ + f_{i,j-1,k} [\beta_{x,y}(0, -1)] + f_{i,j,k} [\beta_{x,y}(0, 0)] + f_{i,j+1,k} [\beta_{x,y}(0, 1)] + \\ + f_{i+1,j-1,k} [\beta_{x,y}(1, -1)] + f_{i+1,j,k} [\beta_{x,y}(1, 0)] + f_{i+1,j+1,k} [\beta_{x,y}(1, 1)]$$

$$\frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial \xi} \right) = f_{i-1,j,k-1} [\beta_{x,\xi}(-1, -1)] + f_{i-1,j,k} [\beta_{x,\xi}(-1, 0)] + f_{i-1,j,k+1} [\beta_{x,\xi}(-1, 1)] + \\ + f_{i,j,k-1} [\beta_{x,\xi}(0, -1)] + f_{i,j,k} [\beta_{x,\xi}(0, 0)] + f_{i,j,k+1} [\beta_{x,\xi}(0, 1)] + \\ + f_{i+1,j,k-1} [\beta_{x,\xi}(1, -1)] + f_{i+1,j,k} [\beta_{x,\xi}(1, 0)] + f_{i+1,j,k+1} [\beta_{x,\xi}(1, 1)]$$

$$\frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial x} \right) = f_{i-1,j-1,k} [\beta_{y,x}(-1, -1)] + f_{i-1,j,k} [\beta_{y,x}(-1, 0)] + f_{i-1,j+1,k} [\beta_{y,x}(-1, 1)] + \\ + f_{i,j-1,k} [\beta_{y,x}(0, -1)] + f_{i,j,k} [\beta_{y,x}(0, 0)] + f_{i,j+1,k} [\beta_{y,x}(0, 1)] + \\ + f_{i+1,j-1,k} [\beta_{y,x}(1, -1)] + f_{i+1,j,k} [\beta_{y,x}(1, 0)] + f_{i+1,j+1,k} [\beta_{y,x}(1, 1)]$$

$$\frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial y} \right) = f_{i,j-1,k} [\alpha_y(-1)] + f_{i,j,k} [\alpha_y(0)] + f_{i,j+1,k} [\alpha_y(1)]$$

$$\frac{\partial}{\partial \xi} \left( g \frac{\partial f}{\partial \xi} \right) = f_{i,j-1,k-1} [\beta_{y,\xi}(-1, -1)] + f_{i,j-1,k} [\beta_{y,\xi}(-1, 0)] + f_{i,j-1,k+1} [\beta_{y,\xi}(-1, 1)] + \\ + f_{i,j,k-1} [\beta_{y,\xi}(0, -1)] + f_{i,j,k} [\beta_{y,\xi}(0, 0)] + f_{i,j,k+1} [\beta_{y,\xi}(0, 1)] + \\ + f_{i,j+1,k-1} [\beta_{y,\xi}(1, -1)] + f_{i,j+1,k} [\beta_{y,\xi}(1, 0)] + f_{i,j+1,k+1} [\beta_{y,\xi}(1, 1)]$$

$$\frac{\partial}{\partial \xi} \left( g \frac{\partial f}{\partial x} \right) = f_{i-1,j,k-1} [\beta_{\xi,x}(-1, -1)] + f_{i-1,j,k} [\gamma_{\xi,x}(-1) \beta_{\xi,x}(-1, 0)] + f_{i-1,j,k+1} [\beta_{\xi,x}(-1, 1)] + \\ + f_{i,j,k-1} [\beta_{\xi,x}(0, -1)] + f_{i,j,k} [\gamma_{\xi,x}(0) \beta_{\xi,x}(0, 0)] + f_{i,j,k+1} [\beta_{\xi,x}(0, 1)] + \\ + f_{i+1,j,k-1} [\beta_{\xi,x}(1, -1)] + f_{i+1,j,k} [\gamma_{\xi,x}(1) \beta_{\xi,x}(1, 0)] + f_{i+1,j,k+1} [\beta_{\xi,x}(1, 1)]$$

$$\frac{\partial}{\partial \xi} \left( g \frac{\partial f}{\partial y} \right) = f_{i,j-1,k-1} [\beta_{\xi,y}(-1, -1)] + f_{i,j-1,k} [\gamma_{\xi,y}(-1) \beta_{\xi,y}(-1, 0)] + f_{i,j-1,k+1} [\beta_{\xi,y}(-1, 1)] + \\ + f_{i,j,k-1} [\beta_{\xi,y}(0, -1)] + f_{i,j,k} [\gamma_{\xi,y}(0) \beta_{\xi,y}(0, 0)] + f_{i,j,k+1} [\beta_{\xi,y}(0, 1)] + \\ + f_{i,j+1,k-1} [\beta_{\xi,y}(1, -1)] + f_{i,j+1,k} [\gamma_{\xi,y}(1) \beta_{\xi,y}(1, 0)] + f_{i,j+1,k+1} [\beta_{\xi,y}(1, 1)]$$

$$\frac{\partial}{\partial \xi} \left( g \frac{\partial f}{\partial \xi} \right) = f_{i,j,k-1} [\alpha_\xi (-1)] + f_{i,j,k} [\alpha_\xi (0)] + f_{i,j,k+1} [\alpha_\xi (1)]$$

Аналогично предыдущим выводам, приведем схемы для сумм операторов.

$$\begin{aligned} \left[ \frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial y} \right) + \frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial x} \right) \right] &= f_{i-1,j-1,k} [2\beta_{x,y} (-1, -1)] + f_{i-1,j+1,k} [2\beta_{x,y} (-1, 1)] + \\ &+ f_{i,f,k} [2\beta_{x,y} (0, 0)] + f_{i+1,j-1,k} [2\beta_{x,y} (1, -1)] + f_{i+1,j+1,k} [2\beta_{x,y} (1, 1)] \end{aligned}$$

$$\begin{aligned} \left[ \frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial \xi} \right) + \frac{\partial}{\partial \xi} \left( g \frac{\partial f}{\partial x} \right) \right] &= f_{i-1,j,k-1} [2\beta_{x,\xi} (-1, -1)] + f_{i-1,j,k} [\gamma_{\xi,x} (-1)] + \\ &+ f_{i-1,j,k+1} [2\beta_{x,\xi} (-1, 1)] + f_{i,f,k} [\gamma_{\xi,x} (0) + 2\beta_{x,\xi} (0, 0)] + f_{i+1,j,k-1} [2\beta_{x,\xi} (1, -1)] + \\ &+ f_{i+1,j,k} [\gamma_{\xi,x} (1)] + f_{i+1,j,k+1} [2\beta_{x,\xi} (1, 1)] \end{aligned}$$

$$\begin{aligned} \left[ \frac{\partial}{\partial y} \left( g \frac{\partial f}{\partial \xi} \right) + \frac{\partial}{\partial \xi} \left( g \frac{\partial f}{\partial y} \right) \right] &= f_{i,j-1,k-1} [2\beta_{y,\xi} (-1, -1)] + f_{i,j-1,k} [\gamma_{\xi,y} (-1)] + \\ &+ f_{i,j-1,k+1} [2\beta_{y,\xi} (-1, 1)] + f_{i,f,k} [\gamma_{\xi,y} (0) + 2\beta_{y,\xi} (0, 0)] + f_{i,j+1,k-1} [2\beta_{y,\xi} (1, -1)] + \\ &+ f_{i,j+1,k} [\gamma_{\xi,y} (1)] + f_{i,j+1,k+1} [2\beta_{y,\xi} (1, 1)] \end{aligned}$$

## Дискретизация уравнений компонентов скорости

Проведем дискретизацию уравнений для компонентов  $u$  и  $v$  скорости среды. Вывод проведем только с уравнением для компоненты  $u$ , так как уравнения идентичны с учетом замен  $u \leftrightarrow v$  и  $x \leftrightarrow y$ , и получить второе можно заменой указанных переменных и индексов дискретизации.

Исходное уравнение

$$\begin{aligned} & 4 \frac{\partial}{\partial x'} \left( \eta \frac{\partial u}{\partial x'} \right) + 4a_x \frac{\partial}{\partial x'} \left( \eta \frac{\partial u}{\partial \xi} \right) + \frac{\partial}{\partial y'} \left( \eta \frac{\partial u}{\partial y'} \right) + a_y \frac{\partial}{\partial y'} \left( \eta \frac{\partial u}{\partial \xi} \right) + 4a_x \frac{\partial}{\partial \xi} \left( \eta \frac{\partial u}{\partial x'} \right) + a_y \frac{\partial}{\partial \xi} \left( \eta \frac{\partial u}{\partial y'} \right) + \\ & + (4a_x^2 + a_y^2 + a_z^2) \frac{\partial}{\partial \xi} \left( \eta \frac{\partial u}{\partial \xi} \right) + (4b_x + b_y) \left( \eta \frac{\partial u}{\partial \xi} \right) = \rho g \frac{\partial s}{\partial x'} - \\ & - 2 \frac{\partial}{\partial x'} \left( \eta \frac{\partial v}{\partial y'} \right) - 2a_y \frac{\partial}{\partial x'} \left( \eta \frac{\partial v}{\partial \xi} \right) - \frac{\partial}{\partial y'} \left( \eta \frac{\partial v}{\partial x'} \right) - a_x \frac{\partial}{\partial y'} \left( \eta \frac{\partial v}{\partial \xi} \right) - \\ & - a_y \frac{\partial}{\partial \xi} \left( \eta \frac{\partial v}{\partial x'} \right) - 2a_x \frac{\partial}{\partial \xi} \left( \eta \frac{\partial v}{\partial y'} \right) - 3a_x a_y \frac{\partial}{\partial \xi} \left( \eta \frac{\partial v}{\partial \xi} \right) - 3c_{xy} \left( \eta \frac{\partial v}{\partial \xi} \right) \end{aligned}$$

## Дискретизация уравнения поверхности

При дискретизации уравнения поверхности можно использовать полученные выше дискретизации операторов. Например, дискретизацию для оператора  $\frac{\partial}{\partial x} \left( g \frac{\partial f}{\partial x} \right)$ .

Запишем исходное уравнение и сразу перейдем к дискретной форме

$$\frac{\partial s}{\partial t} = -\frac{\partial}{\partial x} \left( D_x \frac{\partial s}{\partial x} \right) - \frac{\partial}{\partial y} \left( D_y \frac{\partial s}{\partial y} \right) + \frac{\partial b}{\partial t} + \dot{M}_s - \dot{M}_b$$

$$\begin{aligned} \frac{s_{ijl} - s_{ijl-1}}{\Delta t_l} = & -\frac{1}{\Delta x^2} \left\{ s_{i-1jl} D_{x_{i-\frac{1}{2}jl}} + s_{ijl} \left[ -D_{x_{i-\frac{1}{2}jl}} - D_{x_{i+\frac{1}{2}jl}} \right] + s_{i+1jl} D_{x_{i+\frac{1}{2}jl}} \right\} - \\ & -\frac{1}{\Delta y^2} \left\{ s_{ij-1l} D_{y_{ij-\frac{1}{2}l}} + s_{ijl} \left[ -D_{y_{ij-\frac{1}{2}l}} - D_{y_{ij+\frac{1}{2}l}} \right] + s_{ij+1l} D_{y_{ij+\frac{1}{2}l}} \right\} + \\ & + \frac{b_{ijl} - b_{ijl-1}}{\Delta t_l} + \dot{M}_{s_{ijl}} - \dot{M}_{b_{ijl}} \end{aligned}$$

$$\begin{aligned} s_{i-1jl} \left[ \frac{D_{x_{i-\frac{1}{2}jl}}}{\Delta x^2} \right] + s_{ij-1l} \left[ \frac{D_{y_{ij-\frac{1}{2}l}}}{\Delta y^2} \right] + s_{ijl} \left[ \frac{1}{\Delta t_l} - \frac{D_{x_{i-\frac{1}{2}jl}} + D_{x_{i+\frac{1}{2}jl}}}{\Delta x^2} - \frac{D_{y_{ij-\frac{1}{2}l}} + D_{y_{ij+\frac{1}{2}l}}}{\Delta y^2} \right] + \\ s_{ij+1l} \left[ \frac{D_{y_{ij+\frac{1}{2}l}}}{\Delta y^2} \right] + s_{i+1jl} \left[ \frac{D_{x_{i+\frac{1}{2}jl}}}{\Delta x^2} \right] = s_{ijl-1} \left[ \frac{1}{\Delta t_l} \right] + \frac{b_{ijl} - b_{ijl-1}}{\Delta t_l} + \dot{M}_{s_{ijl}} - \dot{M}_{b_{ijl}} \end{aligned}$$