









SPEEDIER

Integrated Bite Case Management Field Operations Manual

Version 1

Authors: Rosemarie Arroyo, Nai Rui Chng, Franz Egipto, Jobin Maestro, Katie Hampson,

Daria Manalo, Mary Elizabeth Miranda, Betty Quiambao,

Date: 12 June 2019

Table of Contents

Contact Information	2
Introduction and Purpose Roles	3
IBCM	4
Current Procedures	4
Risk Assessments by Public Health Workers at ABTCs	6
Animal investigations by Animal Health Workers	8
Communications	12
Rabies Hotline Error! Bookmark not d	
Tools and training	13
Appendices	13
IBCM Overview	14
2. IBCM Risk Assessments	15
2a. Algorithm for Public Health Workers	15
2b. Paper-based risk assessment form	16
2c. IEC for bite victims/ animal owners to be included with PEP certificates and provided on bite consultations	d 18
3. Animal Investigations	19
3a. Algorithm for designated Animal Health Workers	19
3b. Paper-based Animal Investigation form	19
3c. Checklists for Animal Investigations:	23
Investigation Equipment Checklist	23
Checklist for Investigation Procedures	24
3d. IEC for anyone exposed to a suspect rabid animal	25
3e. Procedures to follow when an animal has been in contact with an identified suspect rabid animal	d 26
3f. Quarantine Notice for non-suspect animals that should be observed	27
English Quarantine Notice	27
Tagalog Quarantine Notice	28

Contact Information

The SPEEDIER project has been developed by researchers from the Field Epidemiology Training Programme Alumni Foundation Inc (FETPAFI), the Research Institute for Tropical Medicine (RITM) and the University of Glasgow, UK.

If you require any further information about the project, please contact the Principal Investigator:

Mary Elizabeth Miranda, DVM, DVPH, PHSAE

Email: megmiranda@fetpafi.org

Phone: (02) 281 2465

For further information about activities in the Province contact the local investigator:

Jobin Maestro, MD, PHSAE

Email: maestromd79@gmail.com

Phone: (+63) 9171525821

In case of any difficulties during IBCM work contact the Disease Surveillance Officer:

Franz Jonathan Egipto, RN, MSN Email: fegipto.fetpafi@gmail.com

Phone: (+63) 9157613967

For peer support and discussion we encourage active participation on the SPEEDIER messenger group:

Facebook Messenger Group: SPEEDIER - San Jose Peer Support Group

General information about SPEEDIER and regular updates on progress and events are posted on the SPEEDIER website:

https://rabiesresearch.github.io/SPEEDIER/

Introduction and Purpose

Integrated Bite Case Management (IBCM) is a strategy that formally engages the medical and veterinary sectors (a 'One Health' approach) to assess the risk of genuine exposure to rabies and the subsequent need for **post-exposure prophylaxis (PEP)**. IBCM is advocated by the World Health Organisation (WHO), to ensure appropriate management of animal bite patients and reduce the costs of PEP once rabies has been controlled. IBCM has also been identified as a potential strategy that can sufficiently enhance surveillance to enable verification of rabies freedom.

The key components of IBCM are, a) **risk assessments** of exposure events by public health workers based on the clinical history of the animal involved and circumstances of the exposure and, b) linking exposure from animals deemed high-risk to **investigations** of the animal. Clear communication between human and animal health sectors on the outcomes of these two components are critical to ensuring exposed victims are treated appropriately, necessary control measures are undertaken, and accurate surveillance information is reported.

Many aspects of IBCM are already operational as part of the Philippines health and veterinary system and surveillance within the National Rabies Prevention and Control Programme (NRPCP). The SPEEDIER project aims to support the implementation of IBCM to augment current practice.

In this Field Operations Manual we provide SOPs for undertaking IBCM, that are in line with the latest WHO technical guidance and adapted for the Philippines context to augment current practice. This manual contains the SOPs and copies of materials for implementing IBCM, including algorithms to guide investigations and risk assessments.

Roles

There are 4 key roles in undertaking IBCM:

- Public Health Workers (PHWs) responsible for patient consultations and risk assessment which lead to prescription and administration of PEP, as well as reporting of information on high risk rabies exposures
- 2. Animal Health Workers (AHWs) responsible for investigating high risk biting animals, which can lead to different management options such as quarantining, euthansia by veterinary officers, sample collection and rapid diagnostic testing as well as reporting of investigation outcomes
- 3. **Provincial Officers** responsible for public health workers or animal health workers in their jurisdiction and the health information systems and control and prevention activities for the NRPCP e.g. Provincial Health Officers (PHOs) and Provincial Veterinary Officers (PVOs), as well as surveillance officers and other personnel.
- **4. Laboratory Personnel** responsible for laboratory diagnosis and reporting in their jurisdiction (regional or national level).

For a given geographic area different personnel might be assigned to these roles depending upon human resource availability and infrastructure. For example, in some areas a Barangay Health Worker may be *designated* as an AHW responsible for investigations, because there is no official AHW assigned to the Municipal Agriculture Office (MAO). Many other people may participate in the process of IBCM or may be interested in or use information from IBCM, and this involvement is valuable but these SOPs focus on the aforementioned roles.

IBCM


Current Procedures

Procedures for managing bite victims and for epidemiological investigations are outlined in Figure 1 and described as follows:

Bite victims typically present to a Rural Health Unit (RHU) where a nurse interviews them, assesses the wound (according to WHO categories I-III), provides wound cleaning and, if necessary, administers tetanus toxoid injection. Patients with either category II or III wounds are referred to an Animal Bite Treatment Centre (ABTC) for post-exposure prophylaxis (PEP); patients with category I wounds are not offered PEP. At the ABTC the rabies nurse (RN) registers the patient before the Medical Doctor (MD) prescribes PEP and refers the patient back to the RN who administers PEP, completes the National Rabies Information System (NaRIS) form and rabies registry and provides the patient with health education. The patient is required to return to the ABTC according to their PEP regimen to complete their PEP course. The updated Thai Red Cross Intradermal Regimen was formerly the recommended PEP regimen in the Philippines, but a recent administrative order updated this to the one-week 2-site ID regimen according to the latest WHO position.


If a patient presents to an RHU, an ABTC or a hospital with signs and symptoms of rabies, palliative care is provided and an investigation is conducted by the Provincial Rabies Coordinator, typically involving staff from both the Provincial Health Office (PHO) and the Provincial Veterinary Office (PVO), and recording the case in the Philippines Integrated Disease Surveillance and Response (PIDSR). If the dog/animal has been killed the veterinarian is expected to collect a sample and submit it to the regional laboratory for diagnostic services.

Figure 1. Procedures for patient management and epidemiological investigations



IBCM procedures to augment existing surveillance and patient management are shown below (Figure 2, italicized red font and arrows indicating communication channels):

Figure 2. Procedures for IBCM. Augmentation of current procedures are shown in red.



Each Task requires reporting & sharing through defined channels

Risk Assessments by Public Health Workers at ABTCs

When a new bite patient presents to an ABTC a risk assessment should be completed using the tailored **mobile phone-based application (app)** to ensure rapid, accurate and standardized recording. The risk assessment comprises questions to assess the biting animal's vaccination history, if known; the animal's outcome (alive or dead or disappeared) following the bite and its health status, as well as the category and severity of the bite (*Appendices - Risk Assessment form*). On submission of risk assessment forms alerts are automatically generated for 'high risk' bites, which should trigger an investigation by the animal health worker. 'High-risk' bites involve animals that die, are killed, disappear or show signs of illness after the bite and therefore are considered suspect for rabies. The subsequent investigation should in many cases be able to evaluate if the responsible animal was rabid and should inform rabies control and prevention activities. Risk assessments should also be undertaken for patients presenting with clinical signs of rabies, also using the IBCM app, and with other procedures for the patient carried out as per current practice. All other aspects of the patient consultation remain the same.

Steps to be undertaken when a bite patient presents are:

- 1. Carry out **risk assessment (Figure 3)** using the app (usually done by a nurse).
- 2. Review risk assessment to confirm the biting animal status and prescribe PEP (usually done by a doctor).
- 3. Administer PEP (usually done by a nurse).
- 4. Submit the risk assessment form using the app so that for "high-risk" bites an alert is generated to trigger an investigation
- 5. Take necessary actions for high- versus low-risk bites:
 - a. Immediately follow up each 'high-risk' bite with a direct phone call to the animal health worker to the designated AHW (Livestock Technician at the MAO or Barangay Health Worker) to confirm that they will investigate and have all the necessary details to do so.
 - b. For animals available for observation that are considered no/low risk, request the patient to observe the animal for 10 days and to immediately report back via the ABTC hotline number if any changes in health are observed.
- 6. Issue PEP certificate sto patients indicating the ABTC hotline number that they should use in the event of observing any signs of illness in the dog, or for other advice on first aid and PEP (Figure 4).

Figure 3. Clinical risk assessment algorithm to determine the status of the biting animal and recommended action.

Clinical Risk Assessment Algorithm




Figure 4. Advice for bite patients and animal owners following a bite patient consultation. Ideally this information should be printed as part of the vaccination certificate.

BITE PATIENT & OWNER ADVICE

- If the biting animal was not vaccinated, observe the animal to confirm it is alive and healthy 10 days after the bite
- If the animal becomes ill during these 10 days, call the ABTC hotline and immediately report to your MAO
- If the animal remains healthy during these days, it currently does not have rabies and no further action is required
- If in any doubt, call the ABTC hotline: XXX-XXX-XXXX

PAYO PARA SA NAKAGAT NA PASYENTE AT MAY-ARI NG HAYOP


- Kung hindi bakunado ang kumagat na hayop, obserbahan ito upang malaman kung ito ay buhay pa at malusog 10 araw mula ng ito'y nakakagat
- Kung ito'y magkasakit sa loob ng 10 araw na ito, tumawag sa ABTC hotline, at makipag-ugnayan sa inyong MAO
- Kung ito'y manatiling malusog, sa loob ng 10 araw na ito, nangangahulugang ito'y walang rabis, wala nang kailangang gawin pa.
- Kung mayroong pag-aalinlangan, tumawag sa ABTC hotline: XXX-XXX-XXXX

Animal investigations by Animal Health Workers

When a *designated* AHW receives an alert of a high-risk bite, they should investigate immediately and always within 24 hours following the **investigation algorithm** (Figure 5). Two checklists are to be followed: one of the investigation procedure and one of the equipment required that should be compiled before departure for the investigation (*Appendices - Investigation checklists*). Wherever possible, the AHW should conduct a field investigation, involving a visual inspection of the animal, recording details via the app, before deciding on a course of action. At the end of the investigation the completed form should be submitted via the app and the PHW called to report the investigation result to guide patient management. The PVO and PHO should always be called directly if the investigation suggests the animal is rabid, so as to guide subsequent control activities.

If an animal is suspected to be rabid a sample should be collected and tested immediately using a **Rapid Diagnostic Test (RDT)**. If the animal is not already dead, the assistance of the PVO should be sought to euthanize the animal. After testing on site the sample should be sent directly to the nearest laboratory (regional laboratory or RITM) for confirmatory testing and sequencing of positive cases. Further interviews should be conducted with the bite victim (or witnesses) and other persons or animals that were bitten, even if the biting animal cannot be found. For all bitten persons/ animals, contact details should be recorded and advice provided on PEP. This may identify other bite victims who have not sought care and is therefore of critical importance. Alternatively, if the biting animal is alive and not vaccinated and not suspect for rabies, a quarantine notice should be issued and the owner should ensure the dog is home quarantined for 10 days (*Figure 8, Appendices - Quarantine Notice*) to confirm the animal's condition. If at any point during the quarantine the dog shows signs of illness the owner should immediately contact the AHW. The AHW is *not* required to check the animal every day, but should send a message every 3 days to check, and if no response is received, a visit may be required.

Figure 5. Animal Investigation Algorithm to evaluate the animal status and determine recommended actions.



Steps to be undertaken for an investigation are:

- 1. Review the checklist and prepare equipment
- 2. Follow up with the patient by phone if possible to initiate the investigation following the questions on the app, and to see if a field investigation is necessary
 - a. If the patient cannot be reached by phone follow up directly with a site visit to track down the patient and animal
- 3. Visit patient/ animal owner to complete the investigation
 - a. undertake a visual check of the biting animal
 - b. record the circumstances of the bite and details of the animal behavior and health according to criteria following the investigation form (*Appendices Animal Investigation Form*), administered via the app.
- 4. Decide on a course of action:
 - a. If the animal is suspected to have rabies and displays clinical signs compatible with rabies, notify the PVO
 - i. Request support to immediately euthanize the animal if it is alive.
 - ii. If the animal is dead or following euthanasia, collect a sample using the RITM recommended technique and carry out the rapid test:
 - 1. Following the manufacturers' instructions, insert swab of brain homogenate into specimen tube containing assay diluent.
 - 2. Mix swab sample with the assay diluent well.
 - 3. Remove device from package and place on flat, dry surface.
 - 4. Using the disposable dropper to add 4 drops of the mixed assay diluent/sample from specimen tube into the sample well. If migration does not appear after 1 minute, add 1 more drop.
 - 5. Read test result 5-10 minutes afterwards and record via app.
 - iii. Conduct further interviews with bite victim(s), witnesses, and any other persons or animals (owners) who were bitten and record on investigation form, even if the biting animal cannot be found. For all bitten persons/ animals, record contact details and provide advice (Figure 6) and follow procedures for in-contact animals (Figure 7).
 - b. If animal is alive and not vaccinated, request the owner to home quarantine the dog for 10 days. Issue a quarantine notice (*Figure 8, Appendices Quarantine Notice*) to confirm the animal's condition and ensure adherence.
 - i. The owner should contact the AHW if at any point during home quarantine the dog shows signs of illness and procedures should be followed for a rabid animal (4a).
- 5. On completion of the investigation, submit the investigation form via the App. This will create and send an automated summary to the RHU and ABTC
- 6. Call the corresponding public health worker to notify them of the investigation result and call the PVO and PHO immediately if the animal is suspect for rabies

Figure 6. Advice for persons identified who have been bitten by an identified suspect rabid animal.

SUSPECT RABIES EXPOSURE ADVICE

- If you, your family or anyone in your community was bitten by a suspect rabid animal, report immediately to your ABTC
- If any of your animals, or other animals in your community were bitten by a suspect animal, contact your Animal Health Worker/ MAO
- If in any doubt, call the ABTC hotline:

XXX-XXX-XXXX

PAYO SA PINAGHIHINALAANG NA-EXPOSE SA RABIES

- Kung ikaw, iyong pamilya, o kung sino man sa iyong kumunidad ay nakagat ng isang pinaghihinalaang may rabies na hayop, ireport agad sa inyong ABTC
- Kung meron sa inyong mga alagang hayop o iba pang hayop sa inyong kumunidad na nakagat ng pinaghihinalaang hayop, kontakin ang inyong Animal Health Worker/ MAO
- Kung mayroong pag-aalinlangan, tumawag sa ABTC hotline: XXX-XXX-XXXX

Figure 7. Procedures to follow for animals identified to have been in contact with a suspect animal.

Animal Exposed to a Suspect Rabid Animal




Figure 8. Quarantine Notice to be administered to dog owners for dogs that have not been vaccinated and require observation. See Appendices.

Protect your community from Rabies! Keep this dog ALIVE and AWAY from people and other animals for 10 days

Based on an assessment it was decided this dog might have rabies and could pose a danger to your community. Please keep the dog alive and away from people for 10 days (quarantine). This gives us time to learn if people in your community were exposed to rabies. As the dog's caretaker, you must follow quarantine regulations. Do not kill the dog yourself because that could expose you and others to rabies. If the dog becomes il or aggressive, a vet will safely kill it. If the dog is healthy after 10 days, it currently does not have rabies and can be safely released.

The investigating Animal Health Worker will put a collar on the dog. Do not remove the collar.

Keep the dog on your property in a place away from people and animals! rabies and could pose a danger to your community. Please

- and animals
 Give the dog food and water during the 10-day quarantine
 period. Keeping the dog alive is the ONLY way to know for
 sure if it exposed others to rabies
- If the dog shows any of these signs during quarantine, contact your Animal Health Worker right away:
 - Lack of appetite Lack of energy
- Lack of appetite
 Lack of energy
 Change in behaviour
 Aggressive behaviour
 Increased salivation
 Difficulty walking
 Change in bark
 If the dog shows signs of illness during quarantine, contact your Animal Health Worker/ MAO
 A vet will then safely capture the dog using drugs that cause a painless death. Any people and animals that had contact with the animal will require a medical assessment with the animal will require a medical assessment
 If you have any doubt call the hotline: XXX-XXX-XXXX

Protektahan ang inyong kumunidad mula sa Rabies! Panatilihing BUHAY ang aso at MALAYO mula sa mga tao at iba pang hayop sa loob ng 10 araw

Base sa pagsusuri napag-desisyunan na itong aso ay maaaring may rabies at pwedeng magdulot ng panganib sa inyong kumunidad. Maari lamang panalilihing buhay ang aso at malayo sa mga tao sa loob ng 10 araw (kuwarentenas). Ito ay nagbibigay ng oras para malaman kung ang mga tao sa iyong kumunidad ay na-expose sa rabies. Bilang taga-pangalaga ng aso, dapati mong sundini ang regulasyon ng kuwarentenas. Huwag mong patayin ang aso dahil maaari mong ma-expose ang iyong sarili at iba pa sa rabies. Kung ang aso ay majksaskit or naging agresibo, ang beterinaryo ang papatay dito sa ligtas na paraan. Kung ang aso ay malusog sa loob ng 10 araw, ito ay walang rabies sa kasalukuran at maaari no padkawalian

- walang rabies sa kasalukuyan at maaari ng pakawalan.
 Ano ang iyong kaliangang gawin:
 Ang nag limbestigang Animal Health Worker ay mag lalagay ng panliig sa aso. Huwag tanggalin ang panliig sa aso. Huwag tanggalin ang panliig Panatilihing ang aso ay nasa iyong pangangalaga sa isang lugar na malayo sa mga tao at hayop.
 Bigyan ang aso ng pagkalin at tubig sa loob ng 10-araw na kuwarentenas. Ang pag papanatiling buhay ang aso lamang ang tanging parana para malaman ng lubos kung ilo'y nakapag-expose ng iba pa sa rabies.
 Kung ang aso ay napanakita ng alim man sa mga sinyales na ilih bahang.

- Norter:

 Kawalan ng ganang kumain

 Kawalan ng lakas

 Pagbabago sa pakiktungo

 Pagiging agresibo

 Nadagdagang paglalaway

 Nahihirapang maglakad

 Pagbabago sa pagtahol

 Kung ang aso ay nagpakita ng sintomas ng sakit habang naka-kuwarentenas, kontakin ang inyong Animal Health Worker/ MAO

 Ang beterinanyo ang siyang bibihag sa aso gamit ang gamot na siyang magdudulot ng walang sakit ha pagkamatay. Sino mang tao o hayop na rikalapit sa nasabing hayop ay mangangallangan ng pagsusuring medikal
- ... Kung ikaw ay mayroong pag-aalinlangan, tumawag sa ho XXX-XXX-XXXX

QUARANTINE NOTICE

Bite date:

Quarantine release date – If the animal is healthy after 10 days the animal can be released on:

- Caretaker:
 1. I understand that I am responsible for monitoring I understand uter am responsible to intollioning the dog's health during quarantine

 I understand that my dog must be kept on my property until the quarantine release data

 understand that my dog must be kept away from
- people and animals I understand that I am responsible for ensuring that
- I understand that I am responsible for ensuring that the dog is provided with food, water and shelter
 I understand that the dog cannot be released until
- approved by the investigating officer
 6.____ I understand that the health and medical treatment
- of others may depend on the results of this quarantine period I understand that I must call the investigating officer if the animal becomes ill or dies

By the authority of the provincial government, your animal has been placed under quarantine to monitor it for signs of rables. If you have further questions contact:

Investigating Animal Health Worker:

Phone number:

PAUNAWA NG KUWARENTENAS

Petsa ng kagat:

Petsa ng pagpapalaya sa Kuwarentenas – Kung ang hayop ay malusog pagkatapos ng 10 araw, ang hayop ay maaari ng pakawalan sa:

Tagapag-alaga:

1. Nauunawaan ko na ako ang responsable sa pag subaybay ng kundisyon ng aso habang naka-kuwarentenas 2. Nauunawaan ko na ang aking aso ay kallangang manabili sa aking pangangalaga hanggang sa petsa ng

pagpapalaya sa kuwarentenas 3._____ Nauunawaan ko na ang aking aso ay kailangang

panatilihing malayo sa mga tao at hayop

panatilihing malayo sa mga tao at hayon

Mauunawaan ko na ako ang responsable na tiyaking
ang aso ay nabibigyan ng pagkain, tubig, at tirahan

Nauunawaan ko na ang aso ay hindi papakawalan
hangga't di pa inaaprobahan ng nag-limestigang opisyal

Mauunawaan ko na ang kalusugan at medikal na
pag-gamot ng iba ay nakadepende sa resulta ng
kuwarentenas

kuwarentenas 7.____Nauunawaan ko na kinakailangan kong tawagan ang nag-iimbestigang opisyal kung ang hayop ay biglang nagkasakit o namatay

Sa pamamagitan ng awtoridad ng pamahalaang panialawigan, ang iyong hayop ay inilagay sa ilalim ng kuwarentenas upang subaybayan ito para sa mga sintomas ng rabies. Kung mayroon kayong karagdagang katanungan makipag-ugnayan lamang sa:

Nag-iimbestigang Manggagawa sa Kalusugan ng Hayop: Numero ng Telepono:

Communications

Communications between PHWs, *designated* AHWs, Provincial Officers and Laboratory personnel is crucial to the success of IBCM. The IBCM app aims to facilitate communications between personnel, but direct phone call or face-to-face communications are essential for effective IBCM. The different communication channels that should be followed are outlined in Figure 9.

During implementation of IBCM, if any difficulties are encountered, your **Disease Surveillance Officer** should be contacted for advice (see page 2).

Task IBCM communication channels Phone call Automated alert/report **Event-based reporting** Risk assessment of bite patient Investigation of Follow up phone call Follow up phone call biting animal Automated high-Automated ABTC/ RHU ABTC/ RHU Investigation report risk bite alert PHW PHW **AHW** DSO DSO Phone call for patients Phone call to confire presenting with clinical signs rabies detection PAO/Pro-vet PAO/Pro-vet PHO & PESU **PHO & PESU** Routine reporting **AHW** MAO BAI Automated monthly IBCM **IBCM** database DoH summary Research team **PHW** ABTC/ RHU **PHO & PESU** BoE **DSOs** Regional Lab/ RITM

Figure 9. Flow of information and communication channels for IBCM implementation

Rabies Hotline

Every ABTC should have an emergency hotline, which members of the public can call should they have concerns about animal bites or rabies. This number should be recorded on patient consultation/ vaccination cards. The hotline can serve a dual purpose as a number that PHWs or AHWs in the community can call if they have any concerns about rabies.

A register or counter book should be used to record all calls to the hotline and serve as a source of information about common questions or concerns. Subsequent review or analysis of this log should inform rabies IEC materials for use locally or for guiding establishment of IBCM in other localities.

The register should include the following variables: the date and time, the caller's name and telephone number (in case they need to be called back and directed to a source of PEP for example), the problem the caller rang about, the suggested solution or response and the name of the call receiver. This information is outlined in the spreadsheet: HotlineRegister.xls.

Tools and training

The app will be used to guide and record risk assessments and animal investigations. The application is installed onto mobile phones for use in the field, or where, available laptops, for example at Municipal Offices and ABTCs. The training programme contains instructions on how to use the app on your device.

Vaccination cards will also be issued to patients indicating the hotline number for the ABTC that patients should use in the event of observing any signs of illness in the dog, or for advice on first aid and PEP (Figure 4).

The peer support group (messenger) is for practitioners to share their experiences, including challenges and to solicit peer support to address them. We encourage participation in the peer support group and updates on progress will also be posted regularly.

Feedback to practitioners will be provided through regular reports summarizing the surveillance data generated by the IBCM (including categorization of exposures and cases according to WHO case definitions), and through regular stakeholder meetings to discuss the interpretation of surveillance data. These data are directly accessible via the IBCM dashboard, which users login to with their IBCM app credentials (username and password). The website to login to the IBCM dashboard is: http://rabies.esurveillance.or.tz/login

Updates will also be available via the SPEEDIER website: https://rabiesresearch.github.io/SPEEDIER/


The training programme for IBCM also includes useful information including:

- 1. The WHO position on rabies PEP updated in 2018 and a visual summary
- 2. Lectures and description of how to use the IBCM app and dashboard
- 3. Lectures on sample collection and RITM SOPS and forms for recording sample collection

Appendices

1. IBCM Overview

Integrated Bite Case Management




Each Task requires reporting & sharing through defined channels

2. IBCM Risk Assessments

2a. Algorithm for Public Health Workers

Clinical Risk Assessment Algorithm



2b. Paper-based risk assessment form

Risk Assessment form

Nurses will complete risk assessments during patient consultations that Medical Doctors will confirm. The following data will be captured

Metadata ABTC Date of consultation Type of consultation (Responsible nurse		lose 3)
Patient characteristics Name Phone number Barangay	Age Pati	•	and 3 consultations) Sex
Bite history (only requ	ired for PEP initiation	on consultations)	
Date bitten			
Biting animal: Select of	dog, cat, livestock (d	cow/shoat /pig), v	vildlife (specify), human
Excessive sali	ggression (incl. bitin vation lullness/lethargy / r paralysis alization out reason g	• ,	e/ animals/ objects)
Is the animal known in If yes, name of owner	•	Barangay	
Is the animal still alive Rabies Assessment d	,	hy/ suspicious fo	r rabies/ sick - not rabies/ unknown
Bite site (tick those the ☐ Head/neck ☐ Tree		ınds □ Legs/	Feet

Bite details (tick those that app	oly)		
□ Scratch □ minor wound	s 🛘 🗀 large wour	nds 🗆 sev	ere (hospitalized)
□ single bite □ multiple bites			
Category of bite			
Treatment (tick those that appl	lv)		
□ None □ wound washi	• /	□ Antibiotics	☐ Immunoglobulin
PEP recommendation:			
 PEP advised & availab 	le		
 PEP advised & referred 	d because none ava	ilable	
 PEP not advised 			

Animal follow up (only required for dose 2 and 3 consultations) Is the animal still alive? Yes/No (select)

Comment (if any)

A bite is considered to require an investigation if, the animal either:

- died,
- was killed or disappeared after the bite,
- if the animal was alive but showed signs of illness.

Upon submission of the risk assessment an alert is sent automatically to the designated animal health worker to trigger the investigation according to the above criteria.

The generated message contains the patient ID, barangay and phone number and the dog owner name if available.

The patient is also given their vaccination card detailing the vaccination schedule and the free hotline number for the patient to call in the event of any concern, including if the biting animal becomes ill.

2c. IEC for bite victims/ animal owners to be included with PEP certificates and provided on bite consultations

BITE PATIENT & OWNER ADVICE

- If the biting animal was not vaccinated, observe the animal to confirm it is alive and healthy 10 days after the bite
- If the animal becomes ill during these 10 days, call the ABTC hotline and immediately report to your MAO
- If the animal remains healthy during these days, it currently does not have rabies and no further action is required
- If in any doubt, call the ABTC hotline: XXX-XXX-XXXX


PAYO PARA SA NAKAGAT NA PASYENTE AT MAY-ARI NG HAYOP

- Kung hindi bakunado ang kumagat na hayop, obserbahan ito upang malaman kung ito ay buhay pa at malusog 10 araw mula ng ito'y nakakagat
- Kung ito'y magkasakit sa loob ng 10 araw na ito, tumawag sa ABTC hotline, at makipag-ugnayan sa inyong MAO
- Kung ito'y manatiling malusog, sa loob ng 10 araw na ito, nangangahulugang ito'y walang rabis, wala nang kailangang gawin pa.
- Kung mayroong pag-aalinlangan, tumawag sa ABTC hotline:
 XXX-XXXX

3. Animal Investigations

3a. Algorithm for designated Animal Health Workers

Animal Investigation Algorithm



PVO - XXX-XXX-XXXX

3b. Paper-based Animal Investigation form

Animal Investigation Form

Designated animal health workers complete investigations if alerted by a high-risk bite trigger from the ABTC or on request from the public due to a suspect rabid animal. During investigations the following data will be captured:

Metadata Paravet	Municipality	Date	of investigation	time
Trigger for investigat	ion (multichoice:	ABTC, PVO	community, other):	
suspect rabidhit by carfound deadsick animal	ure (bite/scratch animal numan rabies ca			
Biting Animal Animal ID	Barangay			
Select □ dog □ cat	□ livestock (co	w/shoat /pig)	□ wildlife (specify)	□ human
Was this animal:	□ Unowned	Owned	□ Unknown	
If owned, name of ov	vner			
Animal age (Tick)	□ Pup <3 mths	s 🗆 juvenile	e <1y 🗆 Adult >1y	□ Unknown
Animal sex (Tick)	□ Male	□ Female	□ Unknown	
Animal vaccinated fo Yes, specify y No Unknown Not applicable	/ear(s):			
Animal signs/ circum Unprovoked a	-		t apply) g people/ animals/ ob	ojects)

- Excessive salivation
- Unexplained dullness/lethargy
- Hypersexuality
- Paresis and/or paralysis
- **Tremors**

- Appetite loss
- Abnormal vocalization
- Restlessness
- Running without reason
- Feeding young
- Eating
- Normal behavior
- None of the above

Was the animal found?	□ Yes	□ No		
Animal outcome (Tick):	□ Alive	□ Dead	□ Disappeared	
If dead, cause of death (Tick): • killed by owner/ by community • killed by animals • hit by car • illness • unknown Other bitten people and animals				
How many people were bitten/scratched by the animal?				
For each person bitten recor Name Whether sough	•	eat as required within t Whether advised to s	• • •	
Patient ID (if already sought	PEP)	Reason for no PEP (if not sought/ initiated)	
For each animal bitten record (records repeat as required within the application): Type: dog cat livestock (cow/shoat /pig) wildlife (specify)				
Owner of animal(s):				

Rabies Assessment (tick):

- healthy
- suspect for rabies
- sick not rabies
- unknown

If the animal is alive but suspect for rabies was the animal euthanized? If the animal is alive and sick, is the animal confined at home and under observation? If the animal was suspect for rabies was a sample collected?

If no, why not:

- Animal disappeared
- Animal decomposed
- Carcass disposed of in unknown location
- Carcass burned

•	Carcass consumed	
•	Other:	
	nple was collected, was a rapid test performed? Results: □ Positive □ Negative □ Inconclusive	☐ Unsatisfactory for testing
Wher	e was the specimen sent:	
•	MAO	
•	PVO	
•	Regional Laboratory:	

Comments if any

Other (Specify):

On completion of the investigation, the designated animal health worker provides advice to the animal owner(s) and the general community on rabies prevention, including the need for vaccination, and to seek and complete PEP in the event of a bite by a suspect rabid dog.

If the investigation results in the identification of a suspect rabid dog, the designated animal health worker will also inform the barangay captain and inform all at-risk persons and animal owners of appropriate action, including urgent health seeking for those who were exposed but had not initiated PEP.

In the event of other people or animals being bitten by a suspect rabid animal, these people/animals will also be visited to provide advice and to complete additional investigations (see IBCM protocol).

Before departure, the designated animal health worker will confirm with the animal owner the need to contact the designated animal health worker urgently if any behavior changes are observed in an animal under observation.

Selected information from the investigation will automatically be sent back to the ABTC where the alert was triggered to provide the investigation result/animal status and forewarning of exposures likely to attend the ABTC as well as verifying those who have already sought care. If the investigation was not initiated by an alert from the ABTC but instead by a community member or direct from the PVO, an alert will also be sent to the nearest ABTC and RHU to inform them of any high risk bites identified and persons who are required to seek PEP.

The designated animal health worker will be expected either to:

a) submit samples collected to the PVO within 24 hours or to the regional laboratory if closer; or b) to alert the PVO that an animal shows clinical signs consistent with rabies and their support is needed the euthanise the animal. In this case the PVO (or representative) would collect and submit the sample.

If the PVO does the investigation themselves, they will arrange for the sample to be sent to the regional laboratory for confirmation and sequencing.

3c. Checklists for Animal Investigations:

Investigation Equipment Checklist

Investigation Equipment Checklist

Reporting Equipment Charged mobile phone Investigation form Quarantine notices Quarantine collars Rabies IEC Pen	Sampling Kit Scalpel handle Scalpel blades Sample tubes (with preservative) Rapid Diagnostic Tests Secondary containers Sealable plastic bags		
Personal Protective Equipment (PPE) Gloves Face mask Safety glasses Laboratory coat First Aid Kit	☐ Straws / pipettes ☐ Tweezers ☐ Scissors ☐ Disinfectant ☐ Sharps container ☐ Garbage bag ☐ Permanent marker		
Specimen Disposal			

- 1. Bury deep in ground or burn and bury remains
- 2. Place sharps in Container and remove PPE
- 3. Put PPE in biohazard bag for disposal at MAO/RHU
- 4. Wash hands

DO NOT LEAVE BODY ON STREET/ OR SHALLOW BURIAL

Investigation Checklist

Before Departure □ Call owner/ contact & conduct verbal investigation □ Check inventory & phone (charged & logged in) □ Report investigation destination to colleague
On Arrival
☐ Locate & consult owner/ contact
☐ Ensure crowds & animals kept away
☐ Develop plan
Does the animal need to be secured?
Secure animal according to plan
Does the animal require euthansia?
☐ Call PVO
Should a specimen be taken?
Ready equipment
☐ Collect specimens in duplicate & label
Test 1 specimen
☐ Conduct Investigation & complete form
☐ Conduct IEC with owner/ victims/ community
☐ Clean area & dispose of all waste
Upon return
Deposit specimens & test with PVO
 □ Organize supplies & equipment □ Review investigation form & submit
☐ Review investigation form & submit

3d. IEC for anyone exposed to a suspect rabid animal

SUSPECT RABIES EXPOSURE ADVICE

- If you, your family or anyone in your community was bitten by a suspect rabid animal, report immediately to your ABTC
- If any of your animals, or other animals in your community were bitten by a suspect animal, contact your Animal Health Worker/ MAO
- If in any doubt, call the ABTC hotline:


XXX-XXX-XXXX

PAYO SA PINAGHIHINALAANG NA-EXPOSE SA RABIES

- Kung ikaw, iyong pamilya, o kung sino man sa iyong kumunidad ay nakagat ng isang pinaghihinalaang may rabies na hayop, ireport agad sa inyong ABTC
- Kung meron sa inyong mga alagang hayop o iba pang hayop sa inyong kumunidad na nakagat ng pinaghihinalaang hayop, kontakin ang inyong Animal Health Worker/ MAO
- Kung mayroong pag-aalinlangan, tumawag sa ABTC hotline: XXX-XXX-XXXX

3e. Procedures to follow when an animal has been in contact with an identified suspect rabid animal

Animal Exposed to a Suspect Rabid Animal



3f. Quarantine Notice for non-suspect animals that should be observed

English Quarantine Notice

Protect your community from Rabies! Keep this dog ALIVE and AWAY from people and other animals for 10 days

Based on an assessment it was decided this dog might have rabies and could pose a danger to your community. Please keep the dog alive and away from people for 10 days (quarantine). This gives us time to learn if people in your community were exposed to rabies. As the dog's caretaker, you must follow quarantine regulations. Do not kill the dog yourself because that could expose you and others to rabies. If the dog becomes ill or aggressive, a vet will safely kill it. If the dog is healthy after 10 days, it currently does not have rabies and can be safely released.

What you need to do:

- The investigating Animal Health Worker will put a collar on the dog. Do not remove the collar.
- Keep the dog on your property in a place away from people and animals
- Give the dog food and water during the 10-day quarantine period. Keeping the dog alive is the ONLY way to know for sure if it exposed others to rabies
- If the dog shows any of these signs during quarantine, contact your Animal Health Worker right away:
 - Lack of appetite
 - Lack of energy
 - Change in behaviour
 - Aggressive behaviour
 - Increased salivation
 - Difficulty walking
 - Change in bark
- If the dog shows signs of illness during quarantine, contact your Animal Health Worker/ MAO
- A vet will then safely capture the dog using drugs that cause a painless death. Any people and animals that had contact with the animal will require a medical assessment
- If you have any doubt, call the hotline: XXX-XXX-XXXX

QUARANTINE NOTICE

Bite date:

Quarantine release date – If the animal is healthy after 10 days the animal can be released on:

Caretaker:

- 1._____ I understand that I am responsible for monitoring the dog's health during quarantine
- 2._____ I understand that my dog must be kept on my property until the quarantine release data
- 3.____ I understand that my dog must be kept away from people and animals
- 4.____ I understand that I am responsible for ensuring that the dog is provided with food, water and shelter
- 5.____ I understand that the dog cannot be released until approved by the investigating officer
- 6.____ I understand that the health and medical treatment of others may depend on the results of this quarantine period 7.____ I understand that I must call the investigating officer if the animal becomes ill or dies

By the authority of the provincial government, your animal has been placed under quarantine to monitor it for signs of rabies. If you have further questions

Investigating Animal Health Worker:

Phone number:

Tagalog Quarantine Notice

Protektahan ang inyong kumunidad mula sa Rabies! Panatilihing BUHAY ang aso at MALAYO mula sa mga tao at iba pang hayop sa loob ng 10 araw

Base sa pagsusuri napag-desisyunan na itong aso ay maaaring may rabies at pwedeng magdulot ng panganib sa inyong kumunidad. Maari lamang panatilihing buhay ang aso at malayo sa mga tao sa loob ng 10 araw (kuwarentenas). Ito ay nagbibigay ng oras para malaman kung ang mga tao sa iyong kumunidad ay na-expose sa rabies. Bilang taga-pangalaga ng aso, dapat mong sundin ang regulasyon ng kuwarentenas. Huwag mong patayin ang aso dahil maaari mong maexpose ang iyong sarili at iba pa sa rabies. Kung ang aso ay nagkasakit or naging agresibo, ang beterinaryo ang papatay dito sa ligtas na paraan. Kung ang aso ay malusog sa loob ng 10 araw, ito ay walang rabies sa kasalukuyan at maaari ng pakawalan

- Ano ang iyong kailangang gawin:

 Ang nag iimbestigang Animal Health Worker ay mag lalagay ng panliig sa aso. Huwag tanggalin ang panliig.

 Panatilihing ang aso ay nasa iyong pangangalaga sa isang lugar na
- malayo sa mga tao at hayop
- Bigyan ang aso ng pagkain at tubig sa loob ng 10-araw na kuwarentenas. Ang pag papanatiling buhay ang aso lamang ang tanging paraan para malaman ng lubos kung ito'y nakapag-expose ng iba na sa rabies
- Kung ang aso ay nagpakita ng alin man sa mga sinyales na ito habang naka-kuwarentenas, agarang kontakin ang inyong Animal Health Worker
 - Kawalan ng ganang kumain
 - Kawalan ng lakas
 - Pagbabago sa pakikitungo
 - Pagiging agresibo
 - Nadagdagang paglalaway
 - Nahihirapang maglakad
 - Pagbabago sa pagtahol
- Kung ang aso ay nagpakita ng sintomas ng sakit habang naka-
- kuwarentenas, kontakin ang inyong Animal Health Worker/ MAO Ang beterinaryo ang siyang bibihag sa aso gamit ang gamot na siyang magdudulot ng walang sakit na pagkamatay. Sino mang tao o hayop na nkalapit sa nasabing hayop ay mangangailangan ng pagsusuring
- Kung ikaw ay mayroong pag-aalinlangan, tumawag sa hotline:

PAUNAWA NG KUWARENTENAS

Petsa ng kagat:

Petsa ng pagpapalaya sa Kuwarentenas – Kung ang hayop ay malusog pagkatapos ng 10 araw, ang hayop ay maaari ng

Tagapag-alaga:

- Nauunawaan ko na ako ang responsable sa pag subaybay ng kundisyon ng aso habang naka-kuwarentenas
- 2. Nauunawaan ko na ang aking aso ay kailangang manatili sa aking pangangalaga hanggang sa petsa ng pagpapalaya sa kuwarentenas
- Nauunawaan ko na ang aking aso ay kailangang panatilihing malayo sa mga tao at hayop
- Nauunawaan ko na ako ang responsable na tiyaking ang aso ay nabibigyan ng pagkain, tubig, at tirahan
- Nauunawaan ko na ang aso ay hindi papakawalan hangga't di pa inaaprobahan ng nag-iimestigang opisyal
- Nauunawaan ko na ang kalusugan at medikal na pag-gamot ng iba ay nakadepende sa resulta ng kuwarentenas
- Nauunawaan ko na kinakailangan kong tawagan ang nag-iimbestigang opisyal kung ang hayop ay biglang nagkasakit o namatay

Sa pamamagitan ng awtoridad ng pamahalaang panlalawigan, ang iyong hayop ay inilagay sa ilalim ng kuwarentenas upang subaybayan ito para sa mga sintomas ng rabies. Kung mayroon kayong karagdagang katanungan makipag-ugnayan lamang sa:

Nag-iimbestigang Manggagawa sa Kalusugan ng Hayop: Numero ng Telepono: