

# Discovery space and science with the PLACID stellar coronagraph

Ruben Tandon \*<sup>1</sup>, Liurong Lin <sup>1</sup>, Axel Potier <sup>1</sup>, Laurent Jolissaint <sup>2</sup>, Audrey Baur <sup>2</sup>, Derya Öztürk Çetni <sup>3</sup> and Jonas G. Kühn <sup>1</sup>

Email: ruben.tandon@unibe.ch

<sup>1</sup> Department of Space and Planetary Sciences, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland


<sup>2</sup> University of Applied Sciences HEIG-VD, Route de Cheseaux 1, 1401 Yverdon-les-Bains, Switzerland

<sup>3</sup> Atatürk University Astronomical Research and Application Center (ATASAM), Lütfü Ülkümen Street. No : 8/3, 25050, Yakutiye / Erzurum, Turkey



## Introduction


- PLACID (Programmable Liquid-crystal Active Coronagraph Imager for the DAG telescope) is a coronagraphic instrument, providing adaptive high-contrast imaging capabilities from H - to Ks - band
- PLACID uses a pixelated Spatial Light Modulator (SLM) to generate coronagraphic focal-plane masks (FPMs) for the first time on a telescope (4 m, DAG observatory, Erzurum, Turkey)
- Instrument delivered in March '24, first light expected by end of 2024
- Remote reconfiguration on-demand to adapt to e.g. observing conditions, multiple star coronagraphy, correcting aberrations, segmented primary mirrors (ELTs, HWO, ...)




Vortex FPM ( $n=2$ ) programmed for a binary star (can be rotated in time to perform ADI) [1]



Have a look at a related poster on simulations of an SLM-based coronagraph!




- Plots: no sky degradation
- Lab contrast curves
- Exposure time  $t_{exp}$ 
  - required for  $\text{SNR} = 5$
  - without coronagraph
  - no speckles
- Post-processing, ADI, CDI  
→ factor 10 improvement



## PLACID targets and discovery space

- Known exoplanets / candidates, Brown Dwarfs, circumstellar disks, binaries/triples
- Gaia, TESS, PLATO direct-imaging follow-ups in the North
- PLACID observational constraints:
  - Site: DEC:  $\geq -24^\circ$
  - TROI AO guide star:  $V \leq 13\text{ mag}$
  - On-sky FOV:  $16'' \times 9.6''$



## Outlook

- Obtaining on-sky contrast curves for PLACID
- Data reduction pipeline

## References

[1] Jonas G. Kühn et al. SLM-based digital adaptive coronagraphy: current status and capabilities. *Advances in Optical and Mechanical Technologies for Telescopes and Instrumentation III*, volume 10706 of SPIE Conference Series, page 107062N, July 2018. doi: 10.1117/12.2312554.

[2] Jonas G. Kühn et al. SLM-based Active Focal-Plane Coronagraphy: Status and future on-sky prospects. *Proceedings of SPIE*, February 2021. doi: 10.48550/arXiv.2102.03201. 1