



Active coronagraphy and
coherence differential
imaging in unpolarized light
with the Swiss Wideband
Active Testbed for
Coronagraphic High-Contrast
Imaging (SWATCHi) 2.0

Axel Potier¹, Audrey Baur², Liurong Lin¹, Ruben Tandon¹, Jonas Kühn¹

¹University of Bern

²Haute Ecole d'Ingénierie et de Gestion du Canton de Vaud

U^b **UNIVERSITÄT
BERN**




Concepts of active/adaptive coronagraphy


- **Definition:** Any component whose properties can be actively modified to increase the SNR of any astrophysical source.
- Could use reflective or transmissive components in any plane: Pupil, Focal, non conjugate.. Modulation of phase, amplitude or polarization.



Moving/Deformable mirrors



Digital Micromirror Devices




Spatial Light Modulators

Concepts of active/adaptive coronagraphy


- **Definition:** Any component whose properties can be actively modified to increase the SNR of any astrophysical source.
- Could use reflective or transmissive components in any plane: Pupil, Focal, non conjugate.. Modulation of phase, amplitude or polarization.
- **Adapting the coronagraph to environmental constrains:**
 - Changing atmosphere turbulence
 - Upcoming extremely large telescope facilities (ground- or space-based) will rely on large segmented primary mirrors, which merit function may evolve in time (i.e. dead/missing/defective segments)
 - Micrometeorites




Moving/Deformable mirrors




Digital Micromirror Devices



Spatial Light Modulators



Leboulleux et al. 2022




Rigby et al. 2022

Concepts of active/adaptive coronagraphy


- **Definition:** Any component whose properties can be actively modified to increase the SNR of any astrophysical source.
- Could use reflective or transmissive components in any plane: Pupil, Focal, non conjugate.. Modulation of phase, amplitude or polarization.
- **Adapting the coronagraph to environmental constrains:**
 - Changing atmosphere turbulence
 - Upcoming extremely large telescope facilities (ground- or space-based) will rely on large segmented primary mirrors, which merit function may evolve in time (i.e. dead/missing/defective segments)
 - Micrometeorites
- **Adapting the coronagraph to science cases constraints:**
 - binary/multiple stars systems
 - resolved stars
 - circumstellar disks
 - Infinite number of observing modes (discovery vs. follow-up/characterization).













Moving/Deformable mirrors



Digital Micromirror Devices



Spatial Light Modulators



Rigby et al. 2022

Leboulleux et al. 2022


HD 98800



NASA/JPL-Caltech/T. Pyle



Active/adaptive coronagraphy with SLMs

- We want to **get rid of pre-manufactured phase masks** and use programmable liquid crystal spatial light modulators (SLMs) instead:
 - > 2π phase retardance devices are commercially available up to H/K-band






Active/adaptive coronagraphy with SLMs

- We want to **get rid of pre-manufactured phase masks** and use programmable liquid crystal spatial light modulators (SLMs) instead:
 - > 2π phase retardance devices are commercially available up to H/K-band
- **Each pixel of a SLM can locally delay the optical path (phase) of incoming light:**
 - Video-rate adjustable birefringence
 - Requirement: linearly-polarized light !



Active/adaptive coronagraphy with SLMs

- We want to **get rid of pre-manufactured phase masks** and use programmable liquid crystal spatial light modulators (SLMs) instead:
 - > 2π phase retardance devices are commercially available up to H/K-band
- **Each pixel of a SLM can locally delay the optical path (phase) of incoming light:**
 - Video-rate adjustable birefringence
 - Requirement: linearly-polarized light !
- In principle, **any phase pattern can be programmed.**
 - μm -sized pixels provide exquisite sampling, also in the focal-plane



The DAG telescope

East Anatolia Observatory (DAG) is the new national observatory of Turkey

- Ritchey-Chrétien architecture **4-m primary mirror** and two Nasmyth foci
- VIS and NIR (up to 3um)
- Established near Erzurum
- Altitude: 3100m, 260 clear nights, median seeing 0.9''
- Under leadership of ATASAM



DAG/PLACID in a nutshell



u^b

UNIVERSITÄT
BERN

heig-vd




heig-vd




MACQUARIE
University
SYDNEY-AUSTRALIA


- PLACID stands for “Programmable Liquid-crystal Active Coronagraphic Imager for the DAG telescope”
 - **World's first SLM-based active high-contrast instrument**
 - H-band ($1.65 \mu\text{m}$) and Ks-band ($2.2 \mu\text{m}$)
 - Versatile instrument ideal for prototyping and mentoring
 - Final Design Review (FDR) passed in Dec 2021
 - **First light by end of 2024**
- PLACID will operate downstream of TROI A
 - «TuRkish adaptive Optics system for Infrared Astronomy»
 - **Extreme AO with pyWSF and 468-DM from AlpAO**
- ... and upstream of DIRAC
 - «DAG InfRAed Camera»
 - **HAWAII-2RG detector**



DAG/PLACID performance at Factory Acceptance Review




DAG/PLACID performance at Factory Acceptance Review




PLACID upgrade first goal: Beat polarization -> Increase throughput

- Use 2 off the shelf LCOS SLMs in cascade, half-wave plate in between




PLACID upgrade first goal: Beat polarization -> Increase throughput

- Use 2 off the shelf LCOS SLMs in cascade, half-wave plate in between




- Use 1 photothermal SLM prototype: active material whose index of refraction is changed by a local temperature gradient



See SPIE paper
#13097-150

PLACID upgrade second goal: coherent differential imaging



Composed of:


- HD 163296 disk
- Static speckles
- Turbulent speckles forming a halo

Results recently obtained with
VLT/SPHERE. To be optimized.


PLACID upgrade second goal: coherent differential imaging

- The residual stellar speckles are incoherent with the off-axis source

Total intensity



Coherent intensity



Composed of:

- HD 163296 disk
- Static speckles
- Turbulent speckles forming a halo

Calibration of the static speckles

Temporal modulation of the speckle through phase diversity


Results recently obtained with
VLT/SPHERE. To be optimized.

PLACID upgrade second goal: coherent differential imaging


- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps

Results recently obtained with
VLT/SPHERE. To be optimized.


Total intensity



Coherent intensity



Incoherent intensity



Composed of:

- HD 163296 disk
- Static speckles
- Turbulent speckles forming a halo

Calibration of the static speckles



Temporal modulation of the speckle through phase diversity

Composed of:

- HD 163296 disk
- Turbulent speckles forming a halo



PLACID upgrade second goal: coherent differential imaging

- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps




PLACID upgrade second goal: coherent differential imaging

- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps







Coherent speckle



PLACID upgrade second goal: coherent differential imaging



- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps




Incoherent off-axis
point source

PLACID upgrade second goal: coherent differential imaging


- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps



Coherent speckle




Incoherent off-axis
point source



PLACID upgrade second goal: coherent differential imaging

- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps


Naive implementation: synchronous noise detection in the temporal Fourier domain:



PLACID upgrade second goal: coherent differential imaging

- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps


Naive implementation: synchronous noise detection in the temporal Fourier domain:



PLACID upgrade second goal: coherent differential imaging

- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps


Naive implementation: synchronous noise detection in the temporal Fourier domain:



PLACID upgrade second goal: coherent differential imaging

- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps


Naive implementation: synchronous noise detection in the temporal Fourier domain:



PLACID upgrade second goal: coherent differential imaging


- The residual stellar speckles are incoherent with the off-axis source
- Creating coherence and incoherence maps

Naive implementation: synchronous noise detection in the temporal Fourier domain:



Design of SWATCHi 2.0 (University of Bern)

- Broadband (H-band)
- One Off-axis incoherent source to demonstrate CDI
- Motorized filter wheels
- 1 Multi-3.5 BMC DM \rightarrow 4mm entrance pupil
- F# = 160




Design of SWATCHi 2.0: latest technology for millisecond CDI

- Broadband (H-band)
- One Off-axis incoherent source to demonstrate CDI
- Motorized filter wheels
- 1 Multi-3.5 BMC DM \rightarrow 4mm entrance pupil
- F# = 160



- 11 pixels/resol
- 23 λ/D FoV



Design of SWATCHi 2.0: latest technology for millisecond CDI

- Broadband (H-band)
- One Off-axis incoherent source to demonstrate CDI
- Motorized filter wheels
- 1 Multi-3.5 BMC DM \rightarrow 4mm entrance pupil
- F# = 160


NEW SINCE 2021 !
High-speed > 200 Hz SLMs

1920 x 1152 Spatial Light Modulator
- High Resolution
- High Speed

With PCIe Controller
supporting frame rates up to 422 Hz



- Pixel size = 9.2um \rightarrow 28.5 spixel/resol




Design of SWATCHi 2.0: latest technology for polarization-insensitive SLM solution

- Broadband (H-band)
- One Off-axis incoherent source to demonstrate CDI
- Motorized filter wheels
- 1 Multi-3.5 BMC DM \rightarrow 4mm entrance pupil
- F# = 160


NEW SINCE 2024!



- Pixel size < 50um \rightarrow 5.2 spixel/resol



Design of SWATCHi 2.0: Testbed simulations for spatial sampling at the SLM



See Liurong Lin's poster !
SPIE paper #13096-128




Design of SWATCHi 2.0: latest technology for polarization-insensitive SLM solution

- Broadband (H-band)
- One Off-axis incoherent source to demonstrate CDI
- Motorized filter wheels
- 1 Multi-3.5 BMC DM \rightarrow 4mm entrance pupil
- F# = 160




NEW SINCE 2024!



- Pixel size < 50um \rightarrow 5.2 spixel/resol





Side project: characterizing SLMs with Digital Holographic Microscopy



SWATCHi 2.0 current status: alignment to be started!



SWATCHi 2.0 current status: alignment to be started!



RACE-GO team at SPIE'24 and steps forwards



Bern lab



2023

Design of SWATCHi 2.0 and hardware procurement

2024

Alignment/interfacing of SWATCHi 2.0

2025

Broadband tests

Full characterizat ion and comparison of SLM technologies

Developing new algorithms for PLACID

2026

Development a viable scheme for CDI

PLACID



Packing and shipping to Turkey

Target list

Development of PLACID exposure time calculator

Commissionning at DAG

PLACID operations:

- Validation of observing modes on known exoplanets, BD and disks
 - High-contrast imaging survey of binaries
 - On-sky tests of customized coronagraphs
- Data reduction pipeline

PLACID upgrade for CDI application

On-sky validation of the CDI

Simulations



Detailed simulations of high-contras

performance with SLMs and PLACID:
- Understand the impact of spatial and electrical resolution on coronagraph null

- Understand PLACID future on-sky results and artifacts

CDI in simulations:

- Deriving the optimal observing strategy with the CDI postprocessing technique

The PI



SPIE poster #13096-129



SPIE poster #13096-128



SPIE talk #13096-58





WE WANT YOU!

Postdoc position at the University of Bern

