EDS 223 Homework 1

Map Making Practice

Rachel Miller

2026-10-10

Table of contents

Why Salinas and Guayama? 1
Load Packages and Source Functions 2
Import Data 2
Data Wrangling 2
Visualize Data 3
Map Interpretation 4
References 4
Data Source 5

Why Salinas and Guayama?

The neighboring Salinas and Guayama municipalities in Puerto Rico are widely recognized as a "sacrifice zone", or an area "where residents are subjected to heightened levels of pollution and hazardous materials, despite the adverse impacts on their health" as defined by the Climate Reality Project (Sacrifice Zones, n.d.). The predominantly low-income and Afro-Puerto Rican Residents are subjected to disproportionate exposure to toxic emissions from several major industrial polluters concentrated in the Salinas-Guayama region, including two fossil-fired power plants (Grist, 2024) and a sterilization facility (Earthjustice, 2026).

Puerto Rico shares many characteristics with the surrounding small island developing states in the Caribbean, including high vulnerability to climate change impacts, but as a United States territory isn't designated as such. This prevents it from benefiting from international frameworks designed to support small island developing states. Concurrently, Puerto Ricans grapple with unequal political representation, leaving them even more vulnerable to environmental challenges like climate change and pollution.

Load Packages and Source Functions

```
library(tidyverse)
library(sf)
library(stars)
library(tmap)
library(here)
```

Import Data

```
# Read in geodatabase of EJScreen data at the Census Block Group level
ejscreen <- sf::st_read(here::here(
  "data", "ejscreen","EJSCREEN_2023_BG_StatePct_with_AS_CNMI_GU_VI.gdb"))
```

```
Reading layer `EJSCREEN_StatePctiles_with_AS_CNMI_GU_VI' from data source
  `/home/rachelcmiller/EDS223-HW/EDS223-HW1/data/ejscreen/EJSCREEN_2023_BG_StatePct_with_AS_
  using driver `OpenFileGDB'
Simple feature collection with 243021 features and 223 fields
Geometry type: MULTIPOLYGON
Dimension:     XY
Bounding box:  xmin: -19951910 ymin: -1617130 xmax: 16259830 ymax: 11554350
Projected CRS: WGS 84 / Pseudo-Mercator
```

Data Wrangling

```
# Filter to Puerto Rico
puerto_rico <- ejscreen %>%
  dplyr::filter(ST_ABBREV == "PR")

# Filter to Salinas-Guayama Municipalities
salinas_guayama <- ejscreen %>%
  dplyr::filter(CNTY_NAME %in% c("Salinas Municipio", "Guayama Municipio"))

# Keep only rows representing land areas
salinas_guayama_clean <- salinas_guayama[-c(32,55),]


# Find the average values for all variables within municipalities
```

```
pr_municipalities <- aggregate(
  puerto_rico,
  by = list(puerto_rico$CNTY_NAME),
  FUN = mean,na.rm = TRUE)
```

Visualize Data

```
# Create a map of percentage low income in Salinas-Guayama
tm_shape(salinas_guayama_clean) +
  tm_polygons(fill = "P_LOWINCPCT",
  fill.legend = tm_legend(title = "Percentage Low Income")) +
  tm_title("Fig.1 : Percentage Low Income Population in Salinas and Guyama") +
  tm_scalebar() +
  tm_compass(position = c("left", "top"))
```


Fig.1 : Percentage Low Income Population in Salinas and Guyama




```
# Create a map of percentile for toxic releases to air in Salinas-Guayama
tm_shape(salinas_guayama_clean) +
  tm_polygons(fill = "P_RSEI_AIR",
  fill.legend = tm_legend(title = "Percentile for Toxic Releases to Air")) +
  tm_title("Fig.2 : Percentile for Toxic Releases to Air in Salinas and Guayama") +
```

```
  tm_scalebar() +
  tm_compass(position = c("left", "top"))
```

Fig.2 : Percentile for Toxic Releases to Air in Salinas and Guayama



Map Interpretation

Figure 1 features a representation of the percentage of low income residents throughout the Salinas and Guayama municipalities in Puerto Rico. From this figure, the highest percentage low-income communities, with 80 to 100% low-income residents, are concentrated in the north-western and south-eastern areas. Figure 2, on the other hand, displays the percentile for toxic chemicals released to air relative to the broader United States in the same region. From Figure 2, the areas with a high percentile are similarly concentrated in the western and south-eastern areas. These areas are classified in the 60th-100th percentile, meaning that 40% or less of the U.S has a higher value than these areas. From comparing the two figures, the areas with high levels of toxic chemicals released to air closely coincide with the areas with the most low-income residents. This suggests that the lower income residents are disproportionately exposed to potential health impacts from these pollutants.

References

Earthjustice. (2026, October 5). New air monitoring finds elevated levels of Cancer-Causing ethylene oxide in Salinas, Puerto Rico - EarthJustice. https://earthjustice.org/press/2026/new-

air-monitoring-finds-elevated-levels-of-cancer-causing-ethylene-oxide-in-salinas-puerto-rico

Grist. (2024, May 18). As fossil fuel plants face retirement, a Puerto Rico community pushes for rooftop solar. https://grist.org/energy/as-fossil-fuel-plants-face-retirement-a-puerto-rico-community-pushes-for-rooftop-solar/

Sacrifice zones. (n.d.). The Climate Reality Project. https://www.climaterealityproject.org/sacrifice-zones

Data Source

The data is sourced from the United States Environmental Protection Agency's previous EJScreen: Environmental Justice Screening and Mapping Tool, and can be accessed from the following link: https://drive.google.com/file/d/1nG6Nj1bXfzQ-FOVMO8Km3eNy4SWu1YcIQ/view