

1 Chapter 12 Inference in a Changing World

Problem 12.4 We build on the HMM in section 12.1. Suppose the observer wants to make a prediction for the future state s_{t+1} , given the measurements x_1, \dots, x_t (i.e., the measurement at time $t+1$ has not been made yet). Derive the mean and variance of the posterior. You can assume the mean and variance of $p(s_t | x_1, \dots, x_{t-1})$ to be known.

Answer

We assume that at time t , the observer has a posterior over s_t :

$$p(s_t | x_1, \dots, x_t) = \mathcal{N}(s_t; \mu_{\text{post},t}, \sigma_{\text{post},t}^2).$$

The generative dynamics of the HMM define the transition model as:

$$p(s_{t+1} | s_t) = \mathcal{N}(s_{t+1}; s_t + \Delta, \sigma_s^2),$$

where Δ is the changing term and σ_s^2 is the process noise variance.

The posterior prediction for s_{t+1} would be:

Marginalizing over the uncertainty in s_t to get the predictive distribution over s_{t+1} :

$$p(s_{t+1} | x_1, \dots, x_t) = \int p(s_{t+1} | s_t) p(s_t | x_1, \dots, x_t) ds_t.$$

Since both $p(s_t | x_{1:t})$ and $p(s_{t+1} | s_t)$ are Gaussians, the result of the convolution is also a Gaussian:

$$p(s_{t+1} | x_1, \dots, x_t) = \mathcal{N}(\mu_{\text{pred},t+1}, \sigma_{\text{pred},t+1}^2),$$

where:

$$\begin{aligned} \mu_{\text{pred},t+1} &= \mu_{\text{post},t} + \Delta, \\ \sigma_{\text{pred},t+1}^2 &= \sigma_{\text{post},t}^2 + \sigma_s^2. \end{aligned}$$

All in all, the predicted future state s_{t+1} has a Gaussian distribution with:

- Mean: $\mu_{\text{post},t} + \Delta$
- Variance: $\sigma_{\text{post},t}^2 + \sigma_s^2$

This concludes the derivation of the predictive posterior for s_{t+1} . ■