

Nova Surgical Institute

NOVA SURGICAL INSTITUTE, 6640 VAN NUYS BLVD, STE 101, Van Nuys, CA, 91405

PROCEDURE NOTE

Date: 07/30/2025

RE: JUAREZ CEJA, JUAN

DOB: 07/25/1978

Preoperative diagnosis:

1. Lumbar facet arthropathy at bilateral L3-L4, L4-L5, L5-S1.
2. Lumbago.

Postoperative diagnosis:

1. Lumbar facet arthropathy at bilateral L3-L4, L4-L5, L5-S1.
2. Lumbago.

Procedure Performed:

1. Local anesthetic and steroid injection into Facet Joint at bilateral L3-L4, L4-L5, L5-S1.
2. Fluoroscopy supervision of the lumbar spine.
3. AP and Lateral views of the lumbar spine.

Surgeon: DAVID GOSTINE, M.D.

Informed Consent:

The procedure was described to the patient and the possible benefits, side effects and complications were discussed in detail. The patient was also offered alternative forms of treatment. The patient after accepting the possible complications signed an informed consent form.

Anesthesia:

Anesthesia services were provided in accordance with the patient's request.

Position:

The patient was positioned in the prone position on the fluoroscopy table. The lower back was prepped draped in a sterile manner.

Procedure of Lumbar Facet Joint Steroid Injection:

RE: JUAREZ CEJA, JUAN

July 30, 2025

Page 2 of 2

Utilizing fluoroscopic guidance in the vertical plane, the endplates of the L3-L4 vertebral bodies were aligned. The image intensifier was then rotated obliquely to the left and the interarticular spaces as well as the superior and inferior articulating processes were identified.

A 27 gauge spinal needle was inserted and directed into the interarticular space at the junction of the superior 2/3rd and inferior 1/3rd of the joint. The facet joint was then injected with 1cc of a mixture of 0.25% Bupivacaine and 10mg of Depo-Medrol.

The same procedure was repeated at the L4-L5 and L5-S1 level on the left side. The procedure was then repeated at the L3-L4, L4-L5 and L5-S1 on the right side.

The patient tolerated the procedure well without any complications and was taken to the recovery room in a stable condition. The vital signs remained stable; there were no sensory or motor changes in the lower extremities.

Plan:

Follow-up in two weeks for an evaluation of the procedure.

Sincerely,

A handwritten signature in black ink, consisting of a large, stylized 'D' followed by a horizontal line and a small loop.

DAVID GOSTINE, M.D.
Pain Management Specialist
California License No:

DOD: 07/30/2025

DOT: 07/30/2025