



Location: SHERMAN OAKS HOSPITAL

PATIENT NAME : JUAREZ, CEJA JUAN LUIS

D-O-B : 07-25-1978

STUDY DATE : 10-12-2024

APPROVAL DATE : 10-16-2024

TO SPEAK TO A RADIOLOGIST: 877-939-7378

PATIENT ID : A311585

ACCESSION NO : A677915

REFERRING PHYSICIAN : SALCEDO RAYMUNDO

RADIOLOGIST : ADIL MAZHAR

MRI OF LEFT ANKLE

TECHNIQUE: Multiplanar, multisequence MRI of the ankle was performed. Sequences were obtained in sagittal, coronal and axial planes and submitted for interpretation.

CLINICAL HISTORY: Pain.

FINDINGS:

ALIGNMENT: Normal.

FLUID:

Tibiotalar: Minimal tibiotalar effusion.

Subtalar: Minimal subtalar effusion.

MEDIAL:

Medial malleolus: Normal.

Tendons:

Posterior tibial tendon: Posterior tibial tendinosis.

Flexor digitorum longus: Normal.

Flexor hallucis longus: Tenosynovitis of the flexor hallucis longus tendon.

Ligaments:

Deltoid ligament complex: Intact.

Spring/plantar calcaneo- navicular ligament: Intact.

LATERAL:

Lateral malleolus: Bone marrow edema involving the lateral malleolus.

Tendons:

Peroneus longus: Normal.

Peroneus brevis: Peroneus brevis tendinosis (image 3, series 7).

Peroneal retinaculum: Intact.

Ligaments:

Anterior inferior tibiofibular(syndesmosis): Intact.

Posterior inferior tibiofibular(syndesmosis): Intact.

Anterior talofibular ligament: Sprain vs partial tear of the anterior talofibular ligament.

Calcaneofibular ligament: Intact.

Posterior talofibular ligament: Intact.

POSTERIOR:

Posterior talus: Normal.

Intermalleolar ligament: Intact.

Achilles tendon: Achilles tendinosis. Peritendinitis of the Achilles tendon.

Plantar fascia: Normal.

Enthesophyte: None.

Tendons:

Anterior tibial tendon: Normal.

Extensor hallucis longus: Normal.

Extensor digitorum longus: Normal.

Peroneus tertius: Absent.

JOINTS:

Tibiotalar joint: Normal.

Subtalar joint: Normal.

Tarsal joints: Minimal fluid signal is seen in some of the tarsal joints. There is a cystic focus adjacent to the talonavicular joint, which may relate to a distended joint recess or a ganglion cyst (image 21, series 3 and image 18, series 6).

OTHER FINDINGS:

Bones(other than subarticular marrow): Normal.

Muscles: Normal.

Tarsal tunnel: Normal.


Sinus tarsi: Minimal fluid signal is seen within the sinus tarsi.

Sesamoid bone:

Miscellaneous: There is mild Kaeger's fat pad edema. There is mild superficial subcutaneous edema overlying the lateral ankle (image 14, series 7). The dorsal artery of the foot is prominent (image 18, series 7).

Impression:

1. Minimal tibiotalar effusion.
2. Minimal subtalar effusion.
3. Posterior tibial tendinosis.
4. Tenosynovitis of the flexor hallucis longus tendon.
5. Bone marrow edema involving the lateral malleolus.
6. Peroneus brevis tendinosis (image 3, series 7).
7. Sprain vs partial tear of the anterior talofibular ligament.
8. Achilles tendinosis. Peritendinitis of the Achilles tendon.
9. There is mild Kaeger's fat pad edema. There is mild superficial subcutaneous edema overlying the lateral ankle (image 14, series 7). The dorsal artery of the foot is prominent (image 18, series 7).
10. Minimal fluid signal is seen in some of the tarsal joints. There is a cystic focus adjacent to the talonavicular joint, which may relate to a distended joint recess or a ganglion cyst (image 21, series 3 and image 18, series 6).
11. Minimal fluid signal is seen within the sinus tarsi.



ADIL MAZHAR
RADIOLOGIST

Time Finalized: 2024-10-16 06:50:25

