



Location: SHERMAN OAKS HOSPITAL

PATIENT NAME : JUAREZ, CEJA JUAN LUIS

D-O-B : 07-25-1978

STUDY DATE : 10-12-2024

APPROVAL DATE : 10-16-2024

TO SPEAK TO A RADIOLOGIST: 877-939-7378

PATIENT ID : A311585

ACCESSION NO : A677918

REFERRING PHYSICIAN : SALCEDO RAYMUNDO

RADIOLOGIST : ADIL MAZHAR

MRI OF RIGHT ANKLE

TECHNIQUE: Multiplanar, multisequence MRI of the ankle was performed. Sequences were obtained in sagittal, coronal and axial planes and submitted for interpretation.

CLINICAL HISTORY: Pain.

FINDINGS:

ALIGNMENT: Normal.

FLUID:

Tibiotalar: Moderate tibiotalar effusion.

Subtalar: Minimal subtalar effusion. A small to medium-sized cystic focus is seen posterior to the subtalar joint which may relate to a distended joint recess or a ganglion cyst (image 9, series 4).

MEDIAL:

Medial malleolus: Normal.

Tendons:

Posterior tibial tendon: Tenosynovitis of the posterior tibial tendon.

Flexor digitorum longus: Tenosynovitis of the flexor digitorum longus tendon.

Flexor hallucis longus: Tenosynovitis of the flexor hallucis longus tendon.

Ligaments:

Deltoid ligament complex: Intact.

Spring/plantar calcaneo- navicular ligament: Intact.

LATERAL:

Lateral malleolus: Bone marrow edema involving the lateral malleolus (image 11, series 7).

Tendons:

Peroneus longus: Normal.

Peroneus brevis: Normal.

Peroneal retinaculum: Intact.

Ligaments:

Anterior inferior tibiofibular(syndesmosis): Intact.

Posterior inferior tibiofibular(syndesmosis): Intact.

Anterior talofibular ligament: Full thickness tear of the anterior talofibular ligament (image 17, series 3).

Calcaneofibular ligament: Sprain vs partial tear of the calcaneofibular ligament (image 13, series 6).

Posterior talofibular ligament: Intact.

POSTERIOR:

Posterior talus: Stieda process of the posterior talus.

Intermalleolar ligament: Intact.

Achilles tendon: Achilles tendinosis. Peritendinitis of the Achilles tendon.

Plantar fascia: Normal.

Enthesophyte: None.

Tendons:

Anterior tibial tendon: Normal.

Extensor hallucis longus: Normal.

Extensor digitorum longus: Normal.

Peroneus tertius: Absent.

JOINTS:

Tibiotalar joint: Normal.

Subtalar joint: Normal.

Tarsal joints: Minimal fluid signal is seen in some of the tarsal joints.

OTHER FINDINGS:

Bones(other than subarticular marrow): Normal.

Muscles: Normal.

Tarsal tunnel: Normal.


Sinus tarsi: Minimal fluid signal is seen within the sinus tarsi.

Sesamoid bone:

Miscellaneous: There is mild calcaneal and Kaeger's fat pad edema.

Impression:

1. Moderate tibiotalar effusion.
2. Minimal subtalar effusion. A small to medium-sized cystic focus is seen posterior to the subtalar joint which may relate to a distended joint recess or a ganglion cyst (image 9, series 4).
3. Tenosynovitis of the posterior tibial tendon.
4. Tenosynovitis of the flexor digitorum longus tendon.
5. Tenosynovitis of the flexor hallucis longus tendon.
6. Bone marrow edema involving the lateral malleolus (image 11, series 7).
7. Full thickness tear of the anterior talofibular ligament (image 17, series 3).
8. Sprain vs partial tear of the calcaneofibular ligament (image 13, series 6).
9. Stieda process of the posterior talus.
10. Achilles tendinosis. Peritendinitis of the Achilles tendon.
11. There is mild calcaneal and Kaeger's fat pad edema.
12. Minimal fluid signal is seen in some of the tarsal joints.
13. Minimal fluid signal is seen within the sinus tarsi.



ADIL MAZHAR

RADIOLOGIST

Time Finalized: 2024-10-16 06:48:14

