

DIGITAL SUMS OF POWERS, FACTORIALS, AND LEAST COMMON MULTIPLES

DAVID G. RADCLIFFE

ABSTRACT. We present logarithmic lower bounds for digital sums of powers, multiples of powers, factorials, and least common multiples, emphasizing the elementary divisibility mechanisms behind these estimates. We then explain how Stewart's theorem, via lower bounds for linear forms in logarithms, gives the corresponding general result for powers.

1. INTRODUCTION

In this article, we prove lower bounds on digital sums of powers, multiples of powers, factorials, and the least common multiple of $\{1, \dots, n\}$, using only elementary number theory. We then explain how Stewart's theorem, via lower bounds for linear forms in logarithms, gives the corresponding general result for powers.

We were inspired by the following problem, which was posed and solved by Waław Sierpiński [6, Problem 209]:

Prove that the sum of digits of the number 2^n (in decimal system) increases to infinity with n .

Conjecturally, the number of nonzero base- b digits of a^n should grow linearly in n for multiplicatively independent a and b , but this remains unproved.

2. A DIVISIBILITY ARGUMENT FOR POWERS

Let $s_b(N)$ denote the sum of the base- b digits of N , and let $c_b(N)$ denote the number of nonzero base- b digits. Since

$$c_b(N) \leq s_b(N) \leq (b-1)c_b(N),$$

the functions c_b and s_b are equivalent up to constants depending only on b .

The following simple lemma is the basic mechanism.

Lemma 1. *Let $2 \leq a < b$ with $a \mid b$. Let $(e_k)_{k \geq 1}$ be integers such that $e_1 \geq 1$ and $a^{e_k} > b^{e_{k-1}}$ for $k \geq 2$. If N is divisible by a^{e_k} but not by b , then $c_b(N) \geq k$.*

Proof. The proof is by induction on k . For $k = 1$ the claim is immediate. For $k \geq 2$, write

$$N = b^{e_{k-1}}q + r, \quad 0 \leq r < b^{e_{k-1}}.$$

Since $N \geq a^{e_k} > b^{e_{k-1}}$, we have $q \geq 1$. Also r is divisible by $a^{e_{k-1}}$, because both N and $b^{e_{k-1}}q$ are; and r is not divisible by b , because N is not. Hence, by induction, $c_b(r) \geq k-1$. The

base- b expansion of N is obtained by adjoining the block of e_{k-1} low-order digits represented by r to the expansion of q , so

$$c_b(N) = c_b(q) + c_b(r) \geq 1 + (k-1) = k.$$

□

Choosing $e_1 = 1$ and $e_k = \lceil re_{k-1} \rceil$, where $r = \log b / \log a > 1$, gives $e_k < r^k / (r-1)$. Lemma 1 therefore implies the following explicit form.

Corollary 1. *Let $2 \leq a < b$ with $a \mid b$. If N is divisible by a^n but not by b , then*

$$c_b(N) > C \log n$$

for all sufficiently large n , where $C > 0$ depends only on a and b . Moreover, any $C < 1/\log(\log b / \log a)$ is admissible.

To apply this to powers a^n , one must remove the trailing zeros in base b . The next lemma ensures that, under a multiplicative independence hypothesis, some prime shared by a and b survives to high order after those trailing zeros are removed.

Lemma 2. *Let $a, b \geq 2$, and suppose $\log a / \log b$ is irrational. If*

$$a^n = b^m t,$$

with $t \in \mathbb{Z}_{\geq 1}$, then there is a prime $p \mid a$ and a constant $C > 0$, depending only on a and b , such that

$$\nu_p(t) \geq Cn,$$

where $\nu_p(t)$ denotes the p -adic valuation of t .

Proof. The vectors $(\nu_p(a))_p$ and $(\nu_p(b))_p$ are linearly independent over $\mathbb{Q}$. Hence there are primes p, q with

$$\nu_p(a)\nu_q(b) - \nu_q(a)\nu_p(b) > 0.$$

Comparing p -adic and q -adic valuations in $a^n = b^m t$ gives

$$n\nu_p(a) = m\nu_p(b) + \nu_p(t), \quad n\nu_q(a) \geq m\nu_q(b).$$

Eliminating m yields

$$\nu_p(t) \geq n \left(\nu_p(a) - \frac{\nu_q(a)}{\nu_q(b)} \nu_p(b) \right),$$

and the coefficient is positive by the choice of p, q . □

Let d be the largest divisor of a all of whose prime factors divide b . Equivalently, d is the smallest divisor of a such that a/d is coprime to b . Combining Corollary 1 with Lemma 2 gives a logarithmic lower bound when d and b are multiplicatively independent.

Theorem 1. *Let $a, b \geq 2$, let d be the smallest divisor of a such that $\gcd(a/d, b) = 1$, and suppose that $\log d / \log b$ is irrational. Then*

$$c_b(a^n) > C \log n$$

for all sufficiently large n , with $C > 0$ depending only on a and b .

Proof. Write $a = dg$ with $\gcd(g, b) = 1$, and write $a^n = b^m s$ with $b \nmid s$. Since $g^n \mid s$, put $s = g^n t$. Then $d^n = b^m t$. By Lemma 2, there is a prime $p \mid d$ with $\nu_p(t) \geq C_1 n$. Since $t \mid s$, we also have $\nu_p(s) \geq C_1 n$. Applying Corollary 1 to s gives $c_b(s) > C \log n$. Finally, a^n and s differ only by a power of b , so they have the same nonzero base- b digits. $\square$

The methods of this section are not sufficient to prove Stewart's theorem, because they apply equally to powers and their multiples. For example, every 3^n divides some number of the form $10^k + 8$, which has only two nonzero decimal digits. Thus a different idea is needed to study powers such as 3^n in base 10, where the base and the exponentiated integer are coprime.

3. FACTORIALS AND LEAST COMMON MULTIPLES

For factorials and least common multiples, the useful divisibility is not by a large prime power but by numbers of the form $b^r - 1$.

Lemma 3 (Balog–Dartyge[1], Stolarsky[8]). *If $m, r \geq 1$ and $b \geq 2$, and if $b^r - 1 \mid m$, then*

$$s_b(m) \geq (b - 1)r.$$

Proof. Group the base- b expansion of m into blocks of length r :

$$m = \sum_{i=0}^{k-1} B_i b^{ri}, \quad 0 \leq B_i < b^r.$$

Let $G(m) = \sum_i B_i$. Since $b^r \equiv 1 \pmod{b^r - 1}$, we have $G(m) \equiv m \pmod{b^r - 1}$. Moreover $G(m) < m$ whenever $m \geq b^r$, while $G(m) = m$ for $m < b^r$. Iterating G therefore carries any positive multiple of $b^r - 1$ down to the unique such multiple below b^r , namely $b^r - 1$. Subadditivity of digital sums gives

$$s_b(G(m)) \leq \sum_i s_b(B_i) = s_b(m),$$

so $s_b(m) \geq s_b(b^r - 1) = (b - 1)r$. $\square$

Taking $m = n!$ or $m = \Lambda_n = \text{lcm}(1, \dots, n)$, we obtain a logarithmic lower bound immediately. If $n \geq b^r - 1$, then $b^r - 1$ divides both $n!$ and Λ_n . With $r = \lfloor \log_b(n + 1) \rfloor$, Lemma 3 yields

$$s_b(n!) \geq (b - 1) \lfloor \log_b(n + 1) \rfloor, \quad s_b(\Lambda_n) \geq (b - 1) \lfloor \log_b(n + 1) \rfloor.$$

Thus both digital sums are $\gg_b \log n$. Similar arguments appear in work of Luca [3]; Sanna [4], building on this lemma and deeper methods, proved the stronger bound

$$s_b(n!) > C_b \log n \log \log \log n$$

for $n > e^e$, and the same estimate holds for $s_b(\Lambda_n)$.

Heuristically, after ignoring trailing zeros, the remaining digits of $n!$ and Λ_n should be roughly equidistributed among $\{0, 1, \dots, b - 1\}$. This suggests much larger orders of growth, namely $s_b(n!) \asymp n \log n$ and $s_b(\Lambda_n) \asymp n$, but such estimates are presently out of reach.

4. STEWART'S THEOREM

The general result for powers requires lower bounds for linear forms in logarithms. We first record a small consequence of Lemma 2.

Lemma 4. *Let $a, b \geq 2$ and suppose $\log a / \log b$ is irrational. If $a^n = b^r t$ with t not divisible by b , then*

$$\log t \asymp_{a,b} n.$$

Proof. The upper bound follows from $t \leq a^n$. The lower bound follows from Lemma 2, since t is divisible by $p^{\lceil Cn \rceil}$ for some prime p . $\square$

Theorem 2 (Stewart[7]). *Let $a, b \geq 2$, and suppose $\log a / \log b$ is irrational. Then there is a constant C , depending only on a and b , such that*

$$c_b(a^n) > \frac{\log n}{\log \log n + C}$$

for all sufficiently large n .

Proof. Let $N = a^n$, and write its nonzero base- b digits in the form

$$N = b^m (d_1 b^{-m_1} + \cdots + d_k b^{-m_k}),$$

where

$$1 = m_1 < m_2 < \cdots < m_k \leq m, \quad 1 \leq d_j \leq b-1.$$

Here m_j measures the distance of the j -th nonzero digit from the leading position: the j -th nonzero digit occurs at exponent $m - m_j$.

We shall prove that consecutive nonzero digit positions cannot be separated by too large a multiplicative factor. Fix $i < k$. Split N at the i -th nonzero digit:

$$q = b^{m_i} \sum_{j \leq i} d_j b^{-m_j}, \quad r = b^m \sum_{j > i} d_j b^{-m_j}.$$

Then q and r are positive integers and

$$N = b^{m-m_i} q + r.$$

Moreover,

$$b^{m_i-1} \leq q < b^{m_i},$$

because q is the integer formed by the first m_i digits of N , and its leading digit is nonzero. Similarly, since the first nonzero digit of r occurs at exponent $m - m_{i+1}$, we have

$$b^{m-m_{i+1}} \leq r < b^{m-m_{i+1}+1}.$$

Since $b^{m-1} < N < b^m$, it follows that

$$b^{-m_{i+1}} < \frac{r}{N} < b^{-m_{i+1}+2}.$$

Now define the linear form

$$\Lambda = -n \log a + (m - m_i) \log b + \log q.$$

The linear form Λ is nonzero. Indeed, if $\Lambda = 0$, then $N = b^{m-m_i} q$, which would force $r = 0$, contradicting the fact that there is at least one nonzero digit after the i -th one.

Applying a theorem of Baker and Wüstholz [2, p. 20] to this form yields the lower bound

$$\log |\Lambda| > -C_1 \max\{1, \log q\} \log n.$$

If $m_i \geq 3$, then $q \geq b^{m_i-1} \geq b^2$, so in particular $\log q > 1$, and the preceding estimate becomes

$$\log |\Lambda| > -C_1 \log q \log n.$$

Using $q < b^{m_i}$, this gives

$$\log |\Lambda| > -C_2 m_i \log n.$$

On the other hand,

$$e^\Lambda = 1 - \frac{r}{N}.$$

Since $q \geq 1$ and $r < b^{m-m_i}$, the decomposition $N = b^{m-m_i}q + r$ gives $N > 2r$. Thus $r/N < 1/2$, and so

$$|\Lambda| = -\log(1 - r/N) < 2r/N.$$

Using $r/N < b^{-m_{i+1}+2}$, we obtain

$$\log |\Lambda| < \log 2 + (-m_{i+1} + 2) \log b.$$

Combining the lower and upper bounds for $\log |\Lambda|$ yields

$$-C_2 m_i \log n < \log 2 + (-m_{i+1} + 2) \log b,$$

and hence

$$m_{i+1} \leq C_3 m_i \log n.$$

This proves the desired estimate when $m_i \geq 3$.

If $m_i < 3$, then q is bounded in terms of b . In this case the Baker–Wüstholz lower bound gives simply

$$\log |\Lambda| > -C_4 \log n.$$

The same upper bound

$$\log |\Lambda| < \log 2 + (-m_{i+1} + 2) \log b$$

then implies

$$m_{i+1} \leq C_5 \log n.$$

Since $m_i \geq 1$, this again gives

$$m_{i+1} \leq C_6 m_i \log n.$$

Thus for every $i < k$,

$$\frac{m_{i+1}}{m_i} \leq C_6 \log n.$$

Taking logarithms gives

$$\log m_{i+1} - \log m_i \leq \log \log n + C_7.$$

Summing over $i = 1, \dots, k-1$, and using $m_1 = 1$, we get

$$\log m_k \leq (k-1)(\log \log n + C_7).$$

It remains to relate m_k to n . Let T be the number of trailing zeroes in the base- b expansion of $N = a^n$, and write

$$N = b^T t, \quad b \nmid t.$$

The last nonzero digit of N occurs at exponent T . Since that same digit has exponent $m - m_k$, we have

$$T = m - m_k.$$

Therefore $t = N/b^T$ has exactly m_k base- b digits. Equivalently,

$$b^{m_k-1} \leq t < b^{m_k},$$

so

$$m_k = \frac{\log t}{\log b} + O(1).$$

By Lemma 4,

$$\log t \asymp_{a,b} n.$$

Hence

$$m_k \asymp_{a,b} n,$$

and therefore

$$\log m_k = \log n + O_{a,b}(1).$$

Combining this with the previous inequality gives

$$\log n + O_{a,b}(1) \leq (k-1)(\log \log n + C_7).$$

After increasing the constant in the denominator, we obtain

$$k > \frac{\log n}{\log \log n + C}$$

for all sufficiently large n . Since $k = c_b(a^n)$, this proves the theorem. $\square$

The irrationality hypothesis is necessary: if $\log a / \log b = r/s \in \mathbb{Q}$, then $a^{sn} = b^{rn}$ has only one nonzero base- b digit for every n . Thus Stewart's theorem supplies the quantitative form of the Senge–Straus criterion [5]: the nonzero digit count of a^n in base b tends to infinity exactly when a and b are multiplicatively independent.

REFERENCES

- [1] A. Balog and C. Dartyge, “On the sum of the digits of multiples,” *Moscow Journal of Combinatorics and Number Theory* **2** (2012), 3–15.
- [2] A. Baker and G. Wüstholz, “Logarithmic forms and group varieties,” *Journal für die reine und angewandte Mathematik* **442** (1993), 19–62. <https://doi.org/10.1515/crll.1993.442.19>
- [3] F. Luca, “The number of non-zero digits of $n!$,” *Canadian Mathematical Bulletin* **45** (2002), 115–118.
- [4] C. Sanna, “On the sum of digits of the factorial,” *Journal of Number Theory* **147** (2015), 836–841.
- [5] H. G. Senge and E. G. Straus, “PV-numbers and sets of multiplicity,” *Periodica Mathematica Hungarica* **3** (1973), 93–100.
- [6] W. Sierpiński, *250 Problems in Elementary Number Theory*, American Elsevier, 1970.
- [7] C. L. Stewart, “On the representation of an integer in two different bases,” *Journal für die reine und angewandte Mathematik* **319** (1980), 63–72.
- [8] K. B. Stolarsky, “Integers whose multiples have anomalous digital frequencies,” *Acta Arithmetica* **38** (1980), 117–128.