

Elementary bounds on digital sums of powers

David G. Radcliffe

March 28, 2026

Some notation

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

For integers $N \geq 0$ and $b \geq 2$, the *base- b expansion* of N is the unique representation of the form

$$N = \sum_{i=0}^{\infty} d_i b^i, \quad d_i \in \{0, 1, \dots, b-1\}.$$

The integers d_i are the *base- b digits* of N ; all but finitely many of these digits are zero.

We write $s_b(N)$ for the sum of the base- b digits of N , and $c_b(N)$ for the number of nonzero digits in that expansion.

A question to start with

Does the sum of decimal digits of 2^n increase to infinity with n ?

n	2^n	$s_{10}(2^n)$
5	32	5
10	1024	7
15	32768	26
20	1048576	31
25	33554432	29

It looks like $\lim_{n \rightarrow \infty} s_{10}(2^n) = \infty$, but the sequence is not monotonic.

The last digit of 2^n

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

We write the decimal expansion of 2^n as

$$2^n = \sum_{i=0}^{\infty} d_i 10^i,$$

where each digit satisfies $d_i \in \{0, 1, \dots, 9\}$.

First easy fact

The last digit d_0 is nonzero, because a power of 2 is never divisible by 10.

So every 2^n already has *at least one* nonzero digit.

The first block of digits

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

Assume $n \geq 4$. Then 2^n is divisible by $2^4 = 16$.

Look at the last four digits of 2^n :

$$2^n \bmod 10^4.$$

This remainder is divisible by 16.

Now suppose the three digits d_1, d_2, d_3 were all zero. Then the last four digits would look like

$$000d_0,$$

so the remainder would be less than 10, which is impossible.

Conclusion

At least one of the digits d_1, d_2, d_3 must be nonzero.

The second block of digits

Now assume $n \geq 14$. Then 2^n is divisible by 2^{14} , and

$$2^{14} = 16384 > 10^4.$$

Look at the last 14 digits:

$$2^n \bmod 10^{14}.$$

This remainder is divisible by 2^{14} .

Now suppose the ten digits $d_4, \dots, d_{13}$ were all zero. Then the last 14 digits would look like

$$0000000000d_3d_2d_1d_0$$

so the remainder would be less than 10^4 , which is a contradiction.

Conclusion

At least one digit in the block $d_4, d_5, \dots, d_{13}$ is nonzero.

What is the pattern?

We keep choosing larger and larger cutoffs:

$$4, 14, 47, 157, \dots$$

so that each new power of 2 is bigger than the previous power of 10.
For example,

$$2^{14} > 10^4, \quad 2^{47} > 10^{14}, \quad 2^{157} > 10^{47}.$$

Then the same argument says:

$$\begin{aligned} n \geq 4 &\implies d_1 + d_2 + d_3 > 0, \\ n \geq 14 &\implies d_4 + \dots + d_{13} > 0, \\ n \geq 47 &\implies d_{14} + \dots + d_{46} > 0, \text{ etc.} \end{aligned}$$

Why this works

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

As n grows, more and more of these disjoint blocks become available.

Each block contains *at least one* nonzero digit.

So the number of nonzero digits of 2^n tends to infinity.

Consequently, the sum of digits of 2^n also tends to infinity.

Generalization to other bases

Since we are using no special properties of 2 and 10 (except $2 \mid 10$) we can generalize this result beyond $a = 2$ and $b = 10$.

Theorem (Radcliffe)

Let $2 \leq a < b$ with $a \mid b$. Suppose N is divisible by a^n but not b . Then there exists $C > 0$ (depending on a, b) such that

$$s_b(N) \geq c_b(N) > C \log n$$

for n sufficiently large.

General Bases

The condition that $a \mid b$ can be weakened considerably.

Corollary

Let $a, b \geq 2$. Let d be the smallest factor of a with $\gcd(a/d, b) = 1$. If $\log(d)/\log(b)$ is irrational, then

$$s_b(a^n) \geq c_b(a^n) > C \log n$$

for sufficiently large n .

Example: $a = 6$ and $b = 12$.

Stewart's Theorem

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

Theorem (Stewart, 1980)

Let $a, b \geq 2$ with $\log(a)/\log(b)$ irrational. Then

$$s_b(a^n) \geq c_b(a^n) > \frac{\log n}{\log \log n + C} - 1$$

for all $n > 4$, where C depends only on a and b .

Note: The irrationality condition is necessary!

If $\log(a)/\log(b) = r/s \in \mathbb{Q}$, then

$$a^{ns} = b^{nr}$$

has only one nonzero digit in base b .

Comparison with our results

- The condition of Stewart's theorem more general than ours.
- However, the conclusion only gives a sublogarithmic lower bound.
- Our results require only elementary number theory.
- Stewart's theorem requires advanced methods.

Question: Can one prove a superlogarithmic lower bound?

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

Baker's Theorem

The proof of Stewart's theorem requires a deep result from transcendental number theory (linear forms in logarithms):

Theorem (Baker-Wüstholz, 1993)

Let $\Lambda = b_1 \log \alpha_1 + \cdots + b_n \log \alpha_n$ where:

- $b_1, \dots, b_n$ are integers with $|b_i| \leq B$
- $\alpha_1, \dots, \alpha_n$ are algebraic with heights $\leq A_i$

If $\Lambda \neq 0$, then

$$\log |\Lambda| > -(16nd)^{2(n+2)} \log A_1 \cdots \log A_n \log B$$

where $d = [\mathbb{Q}(\alpha_1, \dots, \alpha_n) : \mathbb{Q}]$.

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

Proof Strategy

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

Let

$$a^n = b^m \left(d_1 b^{-m_1} + \cdots + d_k b^{-m_k} \right),$$

where $m = \lceil \log_b a^n \rceil$, $k = c_b(a^n)$, $d_i \in \{1, \dots, b-1\}$, and

$$1 = m_1 < \cdots < m_k \leq m.$$

Truncate the base- b expansion of a^n at the i -th nonzero digit:

$$a^n = b^{m-m_i} q_i + r_i$$

Goal: Show digit gaps m_{i+1}/m_i cannot be too large.

Main steps of proof

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

- 1 Approximate a^n by truncating to the i -th nonzero digit.
- 2 Estimate m_{i+1}/m_i in terms of the (relative) truncation error $a^{-n}r_i$.
- 3 Apply Baker's theorem to obtain a lower bound on the truncation error, using the fact that $\Lambda \asymp a^{-n}r_i$.
- 4 Deduce thereby an upper bound on the digit gap, using the fact that $m_{i+1}/m_i \asymp \log(a^{-n}r_i)/\log(q_i)$.
- 5 Compute a lower bound on the number of gaps, and hence the number of nonzero digits.

Thank you!

Elementary
bounds on
digital sums of
powers

David G.
Radcliffe

Sierpinski's
problem

Generalization
to Other
Bases

Stewart's
Theorem

Thank you

Link to ArXiv preprint

