




Pernambuco




Resíduos studentizados vs Ajustados




QQ-plot dos resíduos




Histograma dos resíduos



Resíduos vs Tempo



Cook's distance



Pernambuco

Davies  $p = 0$

Shapiro–Wilk  $p = 0.8795$

Breusch–Pagan  $p = 0.3418$

Durbin–Watson  $p = 0.0101$

| Ano  | Resíduo | Resíduo padronizado | Resíduo studentizado | Cook's distance |
|------|---------|---------------------|----------------------|-----------------|
| 2015 | 0.0072  | 1.9378              | 2.6351               | 1.4082          |
| 2016 | −0.0034 | −0.7047             | −0.6769              | 0.0532          |
| 2017 | −0.0086 | −1.6387             | −1.9324              | 0.1678          |
| 2018 | −0.0012 | −0.2417             | −0.2247              | 0.0063          |
| 2019 | 0.006   | 1.6316              | 1.9189               | 0.9983          |
| 2020 | 0.0015  | 0.3752              | 0.3509               | 0.0387          |
| 2021 | 0.0011  | 0.2334              | 0.2169               | 0.0057          |
| 2022 | 0.0006  | 0.1145              | 0.1061               | 0.0007          |
| 2023 | −0.0065 | −1.2283             | −1.2839              | 0.0833          |
| 2024 | −0.001  | −0.1975             | −0.1834              | 0.0041          |
| 2025 | 0.0042  | 1.0419              | 1.0494               | 0.2985          |