




Rio de Janeiro




Resíduos studentizados vs Ajustados




QQ-plot dos resíduos




Histograma dos resíduos



Resíduos vs Tempo



Cook's distance



Rio de Janeiro

Davies  $p = 0.0001$   
Shapiro–Wilk  $p = 0.8155$   
Breusch–Pagan  $p = 0.2224$   
Durbin–Watson  $p = 0.0521$

| Ano  | Resíduo | Resíduo padronizado | Resíduo studentizado | Cook's distance |
|------|---------|---------------------|----------------------|-----------------|
| 2015 | 0.003   | 0.6471              | 0.6179               | 0.5234          |
| 2016 | -0.0059 | -0.6471             | -0.6179              | 0.0523          |
| 2017 | 0.003   | 0.6471              | 0.6179               | 0.5234          |
| 2018 | 0.0148  | 1.7234              | 2.1028               | 0.5304          |
| 2019 | -0.0019 | -0.1983             | -0.1841              | 0.0037          |
| 2020 | -0.0154 | -1.5134             | -1.7082              | 0.1245          |
| 2021 | -0.013  | -1.2375             | -1.2962              | 0.0577          |
| 2022 | 0.0049  | 0.4633              | 0.4357               | 0.0081          |
| 2023 | 0.0129  | 1.2631              | 1.3309               | 0.0867          |
| 2024 | -0.0035 | -0.367              | -0.3431              | 0.0127          |
| 2025 | 0.0013  | 0.1492              | 0.1384               | 0.004           |