

Project Concept: STEMM Lab – A Real-World STEMM Games

STEMM Lab is a mobile app that transforms real-world physical activities into engaging, game-based **Science, Technology, Engineering, Mathematics, and Medicine (STEMM)** learning experiences. Students complete hands-on challenges using everyday materials, their bodies, and their surroundings, while the app captures data using the phone's camera, sensors, GPS, timers, and analytics.

For each activity, students upload videos, record results, rate the activity, and comment on outcomes, with GPS location tagging enabled. Teams compete on leaderboards, receive timed challenges, and iteratively improve their designs like real scientists and engineers. The app will have appropriate screens for each activity.

This app will be designed for upper Primary School and lower High School students.

App Start-Up

- Enter Team Name
 - Enter First Name of Each Team Member
 - Grade or Year Level
 - Team Discriminator: *Assigned by app*
-

Contents

Project Concept: STEMM Lab – A Real-World STEMM Games.....	1
Engineering Challenges.....	2
Activity 1: Parachute Drop Challenge.....	2
Activity 2: Sound Pollution Hunter.....	4
Activity 3: Hand Fan Challenge.....	6
Activity 4: Earthquake-Resistant Structure.....	8
Health and Medical Sciences.....	10
Activity 5: Human Performance Lab – Stretch Speed & Gracefulness.....	11
Activity 6: Reaction Board Challenge.....	13

Engineering Challenges

Activity 1: Parachute Drop Challenge - DONE

(Engineering + Physics)

Overview

Students design, build, and test a parachute for a small toy to reduce its landing speed and impact force. Teams iterate their designs under time and material constraints, aiming to achieve the slowest and safest landing within a target area.

Equipment

- Mobile phone with *STEMM Lab app*
- Small toy (e.g. army toy soldier)
- Table or elevated surface
- Paper or plastic
- String
- Scissors
- Tape


Instructions

1. Drop the toy without a parachute and record the fall (baseline test).
2. Build a parachute using provided materials.
3. Drop the toy from the same height and record the fall.
4. Review speed and landing accuracy results in the app.
5. Redesign and test up to three prototypes within 20 minutes.
6. Upload videos, results, and team reflections.

Diagram

- Toy attached to parachute
- Drop height marked
- Target landing zone shown on floor

(Simple labelled sketch)



Write-up (on paper):

Predict which parachute design was the best.

Sketch each of the design

Recode the times of the design

Were you correct in timings?

What design was the easiest to make?

	<i>How will it take to hit the ground?</i>	<i>Time (time to first hit the ground)</i>	<i>Were you right?</i>	<i>Time (time from first hitting the ground and stop moving) – need slow motion.</i>
<i>Action 1 (e.g. No parachute as the baseline)</i>				
<i>Action 2 (e.g. use plastic with four corners tied to the toy)</i>				
<i>Action 3</i>				

Discussion: Parachutes and Forces (Plain Language)

Gravity pulls objects downward, causing them to speed up as they fall. A parachute increases air resistance (also called drag). Drag acts upward, opposing the motion and slowing the fall. A slower fall reduces the force when the toy hits the ground, making the landing safer. Engineers improve parachute designs through repeated testing and redesign.

Forces Acting on the Toy

Force	Formula
Downward (weight)	$Weight = mass \times g$
Upward (drag)	Drag force from the parachute
Net (total) force	$Net\ Force = Weight - Drag\ Force$

Newton's Second Law:

$$Net\ Force = mass \times acceleration$$

Calculations

Step 1: Measure the Drop Height

Measure the height of the table or drop surface (distance fallen).

Step 2: Measure the Time

Drop the toy (do not throw it).

Record the time taken to first hit the ground using a phone timer or video.

Step 3: Calculate Final Velocity

Since the toy is dropped, the initial velocity is 0 m/s.

$$Final\ velocity = \frac{distance}{time}$$

Example:

$$1.0\ m / 0.5\ s = 2.0\ m/s$$

Step 4: Calculate Acceleration

$$Acceleration = \frac{Final\ velocity - Initial\ velocity}{time}$$

Initial velocity = 0:

$$Acceleration = 2.0 / 0.5 = 4.0\ m/s^2$$

Step 5: Calculate Net Force

$$Net\ Force = mass \times acceleration$$

Example (mass = 0.20 kg):

$$Net\ Force = 0.20 \times 4.0 = 0.8\ N$$

Step 6: Calculate Drag Force

Calculate the weight:

$$Weight = mass \times g = 0.20 \times 9.8 = 1.96 \text{ N}$$

Calculate drag:

$$Drag \text{ Force} = Weight - Net \text{ Force} = 1.96 - 0.8 = 1.16 \text{ N}$$

G-Force (Stopping Acceleration)

G-force describes how quickly the object slows down when it hits the ground. It is measured in multiples of $g = 9.8 \text{ m/s}^2$.

Using the STEMM App Slow-Motion Video:

Identify the moment the toy first hits the ground.

Measure the time taken for the toy to stop moving (contact time).

Case 1: Object Does Not Bounce

Object goes from impact speed downward to 0 m/s.

$$\Delta v = v_{\text{impact}}$$

G-force formula:

$$g - \text{force} = \frac{\Delta v}{t_{\text{contact}}} \div 9.8$$

Example:

Impact speed = 2.0 m/s

Contact time = 0.05 s

$$g - \text{force} = 2.0 / 0.05 \div 9.8 \approx 4.1 \text{ g}$$

Case 2: Object Bounces

Object goes from impact speed downward to upward velocity at separation (v_{up}).

$$\Delta v = v_{\text{down}} + v_{\text{up}}$$

G-force formula:

$$g - \text{force} = \frac{\Delta v}{t_{\text{contact}}} \div 9.8$$

- **Finding v_{up} if rebound height is unknown:**

Measure time to maximum height after bounce (t_{up}) using slow-motion video.

Calculate upward velocity:

$$v_{up} = g \cdot t_{up}$$

- **Example:**

Impact speed downward: 2.0 m/s

Time to max height after bounce: 0.15 s $\rightarrow v_{up} = 9.8 \cdot 0.15 \approx 1.47 \text{ m/s}$

Contact time: 0.02 s

$$\Delta v = 2.0 + 1.47 = 3.47 \text{ m/s} \quad g\text{-force} = 3.47 / 0.02 \div 9.8 \approx 17.7 \text{ g}$$

- **Observation:** Bouncing increases g-force because the velocity change during contact is larger.

Case	Δv	g-force formula
No bounce	v_{impact}	$g = v_{impact} / t_{contact} \div 9.8$
Bounce	$v_{impact} + v_{up}$	$g = (v_{impact} + v_{up}) / t_{contact} \div 9.8$

Tips for slow-motion video:

- Use a ruler in frame for scale
 - Identify first contact for contact time
 - Identify when object leaves surface for bounce calculation
-

Typical G-Force Ranges and Injury Risk

G-Force Range	Examples	Likely Effects
1–5 g	Standing up quickly, elevators, amusement rides	No injury
5–10 g	Hard falls while running, minor car braking	Possible bruising or strains

G-Force Range	Examples	Likely Effects
10–30 g	Sports collisions, bicycle crashes, car crashes with seatbelts	Serious injuries possible (broken bones, concussions)
30–50 g	Severe car crashes, falls onto hard surfaces	High risk of severe injury
50+ g	Very sudden stops with no cushioning	Life-threatening injuries likely

Important: Duration matters. A brief spike can be survivable, while sustained g-forces are more dangerous.

Student Focus

Primary School students

- Measure time
- Calculate final speed

High School students

- Calculate final velocity
- Calculate acceleration
- Calculate net force
- Calculate drag force
- Calculate g-force

Curriculum Links

Science

- ACSSU076 / ACSSU117 – Forces affect motion
- ACSIS124 – Planning and conducting investigations
- ACSIS126 – Analysing patterns in data

Design & Technologies

- ACTDEP036 – Generate, test, and improve solutions

Mathematics

- ACMMG108 – Measuring speed
 - ACMSP147 – Comparing data and averages
-

Activity 2: Sound Pollution Hunter - DONE

(Environmental Science)

Overview

Students measure and compare sound levels in different classroom activities.


Equipment

- Mobile phone with *STEMM Lab app*

Instructions

1. Measure noise from different actions (dropping objects (pens, books) talking, walking, stamping your feet).
2. Record sound levels and locations.
3. Map loud and quiet zones.

Diagram:



Write-up (on paper):

Predict which action created the loudest sound

Record the results

Were you right?

Any surprises?

Should we wear ear muffs in your classroom?

	<i>Prediction (louder or softer than)</i>	<i>Outcome (dB)</i>	<i>Were you right?</i>
<i>Action 1 (e.g. dropping a book on the table)</i>			

Action 2			
Action 3			

Discussion

Sound intensity varies depending on energy and surfaces. Prolonged loud noise can impact health and concentration.

Sound Levels and Hearing Damage Risk

Sound Level (dB)	Example Sounds	Risk to Hearing
0–30 dB	Whisper, quiet library	No risk
30–60 dB	Normal conversation, classroom noise	Safe for long periods
60–85 dB	Busy traffic, vacuum cleaner	Generally safe, but long exposure can cause fatigue
85–90 dB	Lawn mower, loud classroom, heavy traffic	Hearing damage possible after long exposure
90–100 dB	Motorbike, power tools, loud music	Hearing damage likely after short exposure
100–110 dB	Nightclub, rock concert, chainsaw	Serious hearing damage in minutes
110–120 dB	Siren close by, car horn at 1 m	Painful; immediate damage possible
120–130 dB	Jet engine at close range	Immediate and severe hearing damage
140+ dB	Explosion, gunshot	Instant, permanent hearing damage

Curriculum Links

- ACSSU073 – Sound and energy
 - ACPPS053 – Health and wellbeing
-

Activity 3: Hand Fan Challenge - DONE

(Physics – Air Movement)

Overview

Students test how air movement affects flexible materials.


Equipment

- Paper and cardboard
- Scissors
- Mobile phone
- Sticky Tape
- STEMM Mobile App

Instructions

1. Stand paper upright on a table.
2. Fan air from 30 cm away.
3. Observe and record movement.
4. Repeat with different fan designs and fan distance (15cm, 30, 45cm)
5. Repeat with a cardboard instead of a paper vertical.

Diagram:



Write-up (on paper):

Predict which fan design makes the paper move the most.

Record the results

Were you right?

Any surprises?

*How does **material stiffness** affect the bend angle?*

*How does **fan design** influence air velocity and resulting paper movement?*

*How does **distance from the fan** affect bending?*

	<i>Bend (in degrees)</i>	<i>Outcome (in degrees)</i>	<i>Observation Notes: Were you right?</i>
<i>Design 1 (e.g. 1cm back and forward folds)</i>	30°		
<i>Design 2 (e.g. no folds)</i>			
<i>Design 3</i>			

Discussion

Moving air applies force to objects. Paper bends due to flexibility (plasticity), and repeated bending can weaken it.

Tip: Approximate force using $F \approx k \cdot \theta$. Students can **rank forces by stiffness and bend angle** without exact units if needed.

Optional Challenge

- Estimate a **stiffness coefficient** k for each material by measuring bend angle under a constant fan speed.

1. What k is

For bending a sheet of paper:

$$F \approx k \cdot \theta$$

- F = force applied (N)
- θ = bend angle (radians)
- k = stiffness coefficient, representing **resistance to bending**

In reality, k is related to **flexural rigidity** $D = \frac{Et^3}{12(1-\nu^2)}$, but for classroom experiments we can use **approximate empirical values**.

2. Approximate k values for different papers

Material	Thickness (mm)	Stiffness k (N/rad)	Notes
Thin printer paper	0.1	0.05	Bends very easily
Standard card stock	0.25	0.2	Moderate bend
Thin cardboard	0.5	0.5	Much harder to bend
Corrugated cardboard	3	2–3	Very stiff, almost no bend

These are **rough classroom values** to let students estimate forces. The point is to see **relative differences** between materials.

3. Example calculation

Suppose:

- Thin paper, $k = 0.05 \text{ N/rad}$
- Bend angle observed: $30^\circ \rightarrow \theta = 30^\circ \approx 0.524 \text{ rad}$

$$F \approx k \cdot \theta = 0.05 \cdot 0.524 \approx 0.026 \text{ N}$$

- For thicker cardboard, $k = 0.5$, same bend angle:

$$F \approx 0.5 \cdot 0.524 \approx 0.26 \text{ N}$$

So, the **force required increases strongly with stiffness**.

Curriculum Links

- ACSSU076 – Forces and motion

Activity 4: Earthquake-Resistant Structure - DONE

(Engineering + Earth Science)

Overview

Students design structures that withstand vibration, simulating earthquakes.


Equipment

- Cardboard, paper, scissors, sticky tape, plastic/paper cups.
- Mobile phone with vibration sensor

Instructions

1. Build an anti-vibration layer, by folding paper/cardboard.
2. Place a flat cardboard platform on top.
3. Place the phone in the centre and activate vibration mode on the STEMM App.
4. Modify the structure to reduce movement (e.g. more pillars, more folds, etc)

Diagram



Write-up (on paper):

Predict which fold design makes the phone move the least.

Record the results

Were you right?

Any surprises?

	<i>Phone moves</i>	<i>Outcome (in degrees)</i>	<i>Were you right?</i>
<i>Design 1 (e.g. 4 folds + 4 pillars)</i>	<i>e.g. +/- 1cm</i>	<i>4cm</i>	
<i>Design 2 (e.g. 10 folds + 4 pillars)</i>			

<i>Design 3 (e.g. 3 folds and 6 pillars)</i>			
--	--	--	--

Discussion

Earthquakes cause ground vibrations that can collapse poorly designed structures. Engineers design buildings to absorb and distribute energy safely.

Curriculum Links

- ACSSU096 – Earth processes
- ACTDEP036 – Testing and improving designs

Health and Medical Sciences

Activity 5: Human Performance Lab – Stretch Speed & Gracefulness - DONE

(Medical Science + Biomechanics)

Overview

Students investigate how the human body moves by measuring speed, smoothness, and coordination during controlled stretching activities.

Equipment


- Mobile phone with *STEMM Lab app*
- Open space to move safely

Instructions


1. Hold the phone firmly in one hand. Activate the App vibration sensor.
2. Perform guided movement slowly as shown in the app. Record the vibration.
3. Repeat the activity with vibration feedback enabled.
4. Review speed, smoothness, and range-of-motion data.
5. Upload results and reflect as a group.

Diagram

- Student holding phone
- Direction of movement arrows



Movement 1



Movement 2



Movement 3

Write-up (on paper):

Which movement was the hardest to keep the vibration low?

Record the results

Were you right?

Any surprises?

	<i>Predict Phone Vibration Sensor (absolute)</i>	<i>Outcome (time + movement)</i>	<i>Were you right?</i>
<i>Attempt 1</i>	<i>e.g. +/- 1cm</i>	<i>5 mm in 20 seconds</i>	
<i>Attempt 2</i>		<i>5 mm in 5 seconds</i>	
<i>Attempt 3</i>			

Discussion (Plain Language)

Muscles and joints work together to create movement. Faster movements often reduce control, while smoother movements show better coordination. Sensors in the phone measure how quickly and smoothly the body moves, helping students understand biomechanics and fatigue.

Curriculum Links

Health & Physical Education

- ACPPS051 – Movement skills
- ACPPS054 – Physical performance

Science (Biology)

- ACSSU176 – Structure and function of body systems
-

Activity 6: Reaction Board Challenge - DONE

(Neuroscience + Mathematics)

Overview

Students measure reaction time, coordination, and improvement through repeated digital and physical challenges.

Equipment

- Mobile phone with *STEMM Lab app*
- Clear working space

Instructions

Phase 1 – Tap Reaction

1. Tap the screen as soon as the hidden button appears.
2. Record reaction time.

Rotate through each team member

Phase 2 – Swap Hands

3. Repeat using the non-dominant hand.
4. Compare results.

Rotate through each team member


Phase 3 – Tracing Challenge

5. Trace a moving shape on the screen.
6. Review accuracy and delay.

Rotate through each team member

Diagram

- Phone screen with reaction button
- Tracing path illustration



Write-up (on paper):

Predict your reaction time.

Record the results

Were you right?

Any surprises?

	<i>Reaction Time Prediction</i>	<i>Outcome (time + movement)</i>	<i>Were you right?</i>
<i>Attempt 1</i>	<i>e.g. +/- 1cm</i>	<i>6 seconds delay</i>	
<i>Attempt 2</i>		<i>3 seconds delay</i>	
<i>Attempt 3</i>			

Discussion (Plain Language)

Reaction time measures how quickly the brain processes information and sends signals to muscles. Practice can improve speed and coordination. Comparing hands shows how dominance affects performance.

Curriculum Links

Science Inquiry

- ACSIS130 – Collecting and analysing data

Mathematics

- ACMSP147 – Averages and variation

Health

- ACPPS057 – Understanding physical performance

Activity 7: Breathing Pace Trainer

(Medical Science)

Overview

Students analyse breathing patterns at rest and after exercise.

Equipment


- Mobile phone with *STEMM Lab app*
- Flat surface or mat

Instructions

1. Place the phone gently on the chest.
2. Record breathing at rest.

3. Perform light exercise.
 1. Jog one minute on the spot
 2. 100 star jump
4. Record breathing again and compare results.

Rotate for each team member



Write-up (on paper):

Predict which fold design makes the phone move the least.

Record the results

Were you right?

Any surprises?

	<i>Predict Breathe per minute</i>	<i>Outcome (time + movement)</i>	<i>Were you right?</i>
<i>Breathing at Rest</i>	<i>e.g. 6 breath per minute</i>		
<i>After Exercise 1</i>			
<i>After Exercise 2</i>			

Discussion

Breathing rate increases during exercise to supply more oxygen to muscles. Sensors detect chest movement, helping students visualise breathing patterns.

Curriculum Links

- ACSSU176 – Body systems
 - ACPPS054 – Physical activity and health
-