

Osteoarthritis

Osteoarthritis (OA) also known as **degenerative arthritis, degenerative joint disease**, or **osteoarthrosis**, is a type of **joint disease** that results from breakdown of **joint cartilage** and underlying **bone**.^[1] The most common symptoms are **joint pain** and stiffness. Initially, symptoms may occur only following exercise, but over time may become constant. Other symptoms may include **joint swelling**, decreased **range of motion**, and when the back is affected weakness or numbness of the arms and legs. The most commonly involved joints are those near the ends of the fingers, at the base of the thumb, neck, lower back, knees, and hips. Joints on one side of the body are often more affected than those on the other. Usually the problems come on over years. It can affect work and normal daily activities. Unlike other types of **arthritis**, only the joints are typically affected.^[2]

Causes include previous joint injury, abnormal joint or limb development, and **inherited** factors. Risk is greater in those who are **overweight**, have one leg of a different length, and have jobs that result in high levels of joint stress.^{[2][3]} Osteoarthritis is believed to be caused by mechanical stress on the joint and low grade inflammatory processes.^[4] It develops as cartilage is lost with eventually the underlying bone becoming affected.^[2] As pain may make it difficult to exercise, **muscle loss** may occur.^{[3][5]} Diagnosis is typically based on signs and symptoms with **medical imaging** and other tests occasionally used to either support or rule out other problems. Unlike in **rheumatoid arthritis**, which is primarily an **inflammatory** condition, the joints do not typically become hot or red.^[2]

Treatment includes exercise, efforts to decrease joint stress, **support groups**, and **pain medications**. Efforts to decrease joint stress include resting, the use of a **cane**, and **braces**. Weight loss may help in those who are overweight. Pain medications may include **paracetamol** (acetaminophen). If this does not work **NSAIDs** such as **naproxen** may be used but these medications are associated with greater side effects. **Opioids** if used are generally only recommended short term due to the risk of **addiction**.^[2] If pain interferes with normal life despite other treatments, **joint replacement** surgery may help. An artificial joint, however, only lasts a limited amount of time.^[3] Outcomes for most people with osteoarthritis are good.^[2]

OA is the most common form of arthritis with disease of the knee and hip affecting about 3.8% of people as of 2010.^{[2][6]} Among those over 60 years old about 10% of males and 18% of females are affected.^[3] It is the cause of about 2% of **years lived with disability**.^[6] In Australia

about 1.9 million people are affected,^[7] and in the United States about 27 million people are affected.^[2] Before 45 years of age it is more common in men, while after 45 years of age it is more common in women. It becomes more common in both sexes as people become older.^[2]

1 Signs and symptoms

The main symptom is **pain**, causing **loss of ability** and often stiffness. "Pain" is generally described as a sharp ache or a burning sensation in the associated **muscles** and **tendons**. OA can cause a crackling noise (called "**crepitus**") when the affected joint is moved or touched and people may experience muscle **spasms** and contractions in the tendons. Occasionally, the joints may also be filled with fluid.^[8] Some people report increased pain associated with cold temperature, high humidity, and/or a drop in barometric pressure, but studies have had mixed results.^[9]


OA commonly affects the hands, feet, **spine**, and the large **weight bearing** joints, such as the **hips** and knees, although in theory, any joint in the body can be affected. As OA progresses, the affected joints appear larger, are stiff and painful, and usually feel better with gentle use but worse with excessive or prolonged use, thus distinguishing it from **rheumatoid arthritis**.

In smaller joints, such as at the fingers, hard bony enlargements, called **Heberden's nodes** (on the **distal interphalangeal joints**) and/or **Bouchard's nodes** (on the proximal interphalangeal joints), may form, and though they are not necessarily painful, they do limit the movement of the fingers significantly. OA at the toes leads to the formation of **bunions**, rendering them red or swollen. Some people notice these physical changes before they experience any pain.

OA is the most common cause of a **joint effusion** of the knee.^[10]

2 Risk factors

Damage from mechanical stress with insufficient self repair by joints is believed to be the primary cause of osteoarthritis.^[11] Sources of this stress may include misalignments of bones caused by congenital or pathogenic causes; mechanical injury; excess body weight; loss of strength in the muscles supporting a joint; and impair-



Osteoarthritis most often occurs in the hands (at the ends of the fingers and thumbs), neck, lower back, knees, and hips

ment of peripheral nerves, leading to sudden or uncoordinated movements.^[11] However **exercise**, including running in the absence of injury, has not been found to increase the risk.^[12] Nor has cracking one's knuckles been found to play a role.^[13]

2.1 Primary

A number of studies have shown that there is a greater prevalence of the disease among **siblings** and especially

identical twins, indicating a hereditary basis.^[14] Although a single factor is not generally sufficient to cause the disease, about half of the variation in susceptibility has been assigned to genetic factors.^[15]

As early human ancestors evolved into bipeds, changes occurred in the pelvis, hip joint and spine which increased the risk of osteoarthritis.^[16] Additionally genetic variations that increase the risk were likely not selected against because usually problems only occur after reproductive success.^[17]

The development of OA is correlated with a history of previous joint injury and with obesity, especially with respect to knees.^[18] Since the correlation with obesity has been observed not only for knees but also for non-weight bearing joints and the loss of body fat is more closely related to symptom relief than the loss of body weight, it has been suggested that there may be a metabolic link to body fat as opposed to just mechanical loading.^[19]

Changes in sex hormone levels may play a role in the development of OA as it is more prevalent among post-menopausal women than among men of the same age.^{[20][21]} A study of mice found natural female hormones to be protective while injections of the male hormone **dihydrotestosterone** reduced protection.^[22]

2.2 Secondary



lateral



front


Secondary osteoarthritis (due to an old injury with fracture) of the ankle in a woman of 82 years old

This type of osteoarthritis is caused by other factors but the resulting pathology is the same as for primary os-

teoarthritis:

- **Alkaptonuria**
- **Congenital disorders of joints**
- **Diabetes** doubles the risk of having a joint replacement due to OA and people with diabetes have joint replacements at a younger age than those without diabetes.^[23]
- **Ehlers-Danlos Syndrome**
- **Hemochromatosis and Wilson's disease**
- **Inflammatory diseases** (such as **Perthes' disease**), (**Lyme disease**), and all chronic forms of arthritis (e.g., **costochondritis**, **gout**, and **rheumatoid arthritis**). In gout, uric acid crystals cause the cartilage to degenerate at a faster pace.
- **Injury to joints or ligaments** (such as the **ACL**), as a result of an accident or orthopedic operations.
- **Ligamentous** deterioration or instability may be a factor.
- **Marfan syndrome**
- **Obesity**
- **Joint infection**

3 Pathophysiology



Hip joint with osteoarthritis^[24]

While OA is a degenerative joint disease that may cause gross cartilage loss and morphological damage to other joint tissues, more subtle biochemical changes occur in the earliest stages of OA progression. The water content of healthy cartilage is finely balanced by compressive force driving water out & swelling pressure drawing water in.^[25] Collagen fibres exert the compressive force, whereas the **Gibbs–Donnan effect** & cartilage proteoglycans create osmotic pressure which tends to draw water in.^[25]

However, during onset of OA, the collagen matrix becomes more disorganized and there is a decrease in proteoglycan content within cartilage. The breakdown of collagen fibers results in a net increase in water content.^{[26][27][28][29][30]} This increase occurs because whilst there is an overall loss of proteoglycans (and thus a decreased osmotic pull),^{[27][31]} it is outweighed by a loss of collagen.^{[25][31]} Without the protective effects of the proteoglycans, the **collagen** fibers of the cartilage can become susceptible to degradation and thus exacerbate the degeneration. **Inflammation** of the **synovium** (joint cavity lining) and the surrounding **joint capsule** can also occur, though often mild (compared to what occurs in rheumatoid arthritis). This can happen as breakdown products from the cartilage are released into the synovial space, and the cells lining the joint attempt to remove them.

Other structures within the joint can also be affected.^[32] The **ligaments** within the joint become thickened and **fibrotic** and the **menisci** can become damaged and wear away.^[33] Menisci can be completely absent by the time a person undergoes a **joint replacement**. New bone outgrowths, called “spurs” or **osteophytes**, can form on the margins of the joints, possibly in an attempt to improve the congruence of the **articular cartilage** surfaces in the absence of the menisci. The **subchondral bone** volume increases and becomes less mineralized (hypomineralization).^[34] All these changes can cause problems functioning. The **pain** in an osteoarthritic joint has been related to thickened **synovium**^[35] and **subchondral bone lesions**.^[36]

4 Diagnosis

Diagnosis is made with reasonable certainty based on history and clinical examination.^{[37][38]} **X-rays** may confirm the diagnosis. The typical changes seen on X-ray include: joint space narrowing, subchondral **sclerosis** (increased bone formation around the joint), subchondral **cyst** formation, and **osteophytes**.^[39] Plain films may not correlate with the findings on physical examination or with the degree of pain.^[40] Usually other imaging techniques are not necessary to clinically diagnose OA.

In 1990, the **American College of Rheumatology**, using

data from a multi-center study, developed a set of criteria for the diagnosis of hand OA based on hard tissue enlargement and swelling of certain joints.^[41] These criteria were found to be 92% sensitive and 98% specific for hand OA versus other entities such as rheumatoid arthritis and spondyloarthropathies.^[42]

Related pathologies whose names may be confused with OA include pseudo-arthritis. This is derived from the Greek words pseudo, meaning “false”, and arthrosis, meaning “joint.” Radiographic diagnosis results in diagnosis of a fracture within a joint, which is not to be confused with OA which is a degenerative pathology affecting a high incidence of distal phalangeal joints of female patients. A polished ivory-like appearance may also develop on the bones of the affected joints, reflecting a change called **eburnation**.^[43]

- Severe osteoarthritis and osteopenia of the carpal joint and 1st carpometacarpal joint.
- MRI of osteoarthritis in the knee, with characteristic narrowing of the joint space.
- Primary osteoarthritis of the left knee. Note the **osteophytes**, narrowing of the joint space (arrow), and increased subchondral bone density (arrow).
- Damaged cartilage from sows. (a) cartilage erosion (b) cartilage ulceration (c) cartilage repair (d) osteophyte (bone spur) formation.
- Histopathology of osteoarthrosis of a knee joint in an elderly female.
- Histopathology of osteoarthrosis of a knee joint in an elderly female.
- In a healthy joint, the ends of bones are encased in smooth cartilage. Together, they are protected by a joint capsule lined with a synovial membrane that produces synovial fluid. The capsule and fluid protect the cartilage, muscles, and connective tissues.
- With osteoarthritis, the cartilage becomes worn away. Spurs grow out from the edge of the bone, and synovial fluid increases. Altogether, the joint feels stiff and sore.
- thumb|Osteoarthritis

4.1 Classification


A number of classification systems are used for gradation of osteoarthritis:

- **WOMAC** scale
- **Kellgren-Lawrence** grading scale

OA can be classified into either primary or secondary depending on whether or not there is an identifiable underlying cause.

Both primary generalized nodal OA and erosive OA (EOA, also called inflammatory OA) are sub-sets of primary OA. EOA is a much less common, and more aggressive inflammatory form of OA which often affects the distal interphalangeal joints of the hand and has characteristic articular erosive changes on x-ray.^[44]

5 Management



People with osteoarthritis should do different kinds of exercise for different benefits to the body.

Lifestyle modification (such as weight loss and exercise) and analgesics are the mainstay of treatment. Acetaminophen (also known as paracetamol) is recommended first line with NSAIDs being used as add on therapy only if pain relief is not sufficient.^[45] This is due to the relative greater safety of acetaminophen.^[45]

5.1 Lifestyle modification

For overweight people, **weight loss** may be an important factor.^[46] Patient education has been shown to be helpful in the self-management of arthritis.^[46] It decreases pain, improves function, reduces stiffness and fatigue, and reduces medical usage.^[46] Patient education can provide on average 20% more pain relief when compared to NSAIDs alone in patients with hip OA.^[46]

5.2 Physical measures

Moderate exercise is beneficial with respect to pain and function in those with osteoarthritis of the knee and hip.^{[47][48]} These exercises should occur at least three times per week.^[49] While some evidence supports certain **physical therapies**, evidence for a combined program is limited.^[50] There is not enough evidence to determine the effectiveness of **massage therapy**.^[51] The evidence for **manual therapy** is inconclusive.^[52] Functional, gait, and balance training has been recommended to address impairments of position sense, balance, and strength in individuals with lower extremity arthritis as these can contribute to higher falls in older individuals.^[53]

Lateral wedge insoles do not appear to be useful in osteoarthritis of the knee.^{[54][55]} Knee braces may be useful.^[56] For pain management heat can be used to relieve stiffness, and cold can relieve muscle spasms and pain.^[57]

5.3 Medication

The analgesic **acetaminophen** is the first line treatment for OA.^{[45][59]} However, a 2015 review found acetaminophen to only have a small short term benefit.^[60] For mild to moderate symptoms effectiveness is similar to **non-steroidal anti-inflammatory drugs (NSAIDs)**, though for more severe symptoms NSAIDs may be more effective.^[45] NSAIDs such as **naproxen** while more effective in severe cases are associated with greater side effects such as **gastrointestinal bleeding**.^[45] Another class of NSAIDs, **COX-2 selective inhibitors** (such as **celecoxib**) are equally effective to NSAIDs with lower rates of adverse gastrointestinal effects but higher rates of cardiovascular disease such as **myocardial infarction**.^[61] They are also more expensive than non-specific NSAIDs.^[62] Oral **opioids**, including both weak opioids such as **tramadol** and stronger opioids, are also often prescribed. Their appropriateness is uncertain and opioids are often recommended only when first line therapies have failed or are contraindicated.^{[63][63][64]} This is due to a small benefit and relatively large risk of side effects.^[65] Oral **steroids** are not recommended in the treatment of OA.^[59]

There are several NSAIDs available for **topical** use including **diclofenac**. Topical and oral diclofenac work

equally well with topical having a greater risk of mild skin reactions but no greater risk of gastrointestinal adverse effects.^[66] Transdermal **opioid pain medications** are not typically recommended in the treatment of osteoarthritis.^[67] **Topical capsaicin** is controversial with some reviews finding benefit^{[68][69]} and others not.^[70]

Joint injections of glucocorticoids (such as **hydrocortisone**) leads to short term pain relief that may last between a few weeks and a few months.^[71] Injections of **hyaluronic acid** have not been found to lead to much improvement compared to placebo^{[72][73]} but have been associated with harm.^[73] The effectiveness of injections of **platelet-rich plasma** is unclear; there are suggestions that such injections improve function but not pain and are associated with increased risk.^{[74][75]}

5.4 Surgery

If problems are significant and more conservative management is ineffective, **joint replacement surgery** or resurfacing may be recommended. Evidence supports joint replacement for both knees and hips as it is both clinically effective,^{[76][77]} and cost-effective.^{[78][79]} Surgery to transfer articular cartilage from a non-weight-bearing area to the damaged area is one possible procedure that has some success but there are problems getting the transferred cartilage to integrate well with the existing cartilage at the transfer site.^[80]

Osteotomy may be useful in people with knee osteoarthritis but has not been well studied.^[81] **Arthroscopic surgery** is largely not recommended as it does not improve outcomes in knee osteoarthritis.^{[82][83]} Additionally arthroscopy may result in harm.^[84]

5.5 Alternative medicine

Many **dietary supplements** are sold as treatments for OA. Since **glucosamine** is a precursor for a component of **cartilage**, it has been studied for prevention and treatment. The effectiveness of glucosamine is controversial.^{[85][86]} Most recent reviews found it to be equal to^{[87][88]} or only slight better than **placebo**.^{[89][90]} A difference may exist between glucosamine sulfate and glucosamine hydrochloride, with glucosamine sulfate showing a benefit and glucosamine hydrochloride not.^[91] The evidence for glucosamine sulfate having an effect on OA progression is somewhat unclear and if present likely modest.^[92] The **Osteoarthritis Research Society International** recommends that glucosamine be discontinued if no effect is observed after six months^[93] and the **National Institute of Clinical Excellence** no longer recommends its use.^[5] Despite the difficulty in determining the efficacy of glucosamine, it remains a viable treatment option.^[94] Its use as a therapy for osteoarthritis is usually safe.^[94]


Phytodolor,^[68] SAME,^[95] and SKI 306X (a Chinese herbal mixture)^[69] may be effective in improving pain, and there is some evidence to support the use of *cat's claw* as an anti-inflammatory.^[96] There is tentative evidence to support avocado/soybean unsaponifiables (ASU),^{[69][97]} methylsulfonylmethane,^[68] and rose hip.^[68] A few high-quality studies of *Boswellia serrata* show consistent, but small, improvements in pain and function among people with osteoarthritis.^[98]

There is little evidence supporting benefits for some supplements, including: the Ayurvedic herbal preparations with brand names Articulín F and Eazmov, collagen, devil's claw, Duhuo Jisheng Wan (a Chinese herbal preparation), fish liver oil, ginger, the herbal preparation gitadyl, glucosamine, hyaluronic acid, omega-3 fatty acids, the brand-name product Reumalax, stinging nettle, turmeric, vitamins A, C, and E in combination, vitamin E alone, vitamin K and willow bark. There is insufficient evidence to make a recommendation about the safety and efficacy of these treatments.^{[68][96]}

While acupuncture leads to improvements in pain relief, this improvement is small and may be of questionable importance. Waiting list-controlled trials for peripheral joint osteoarthritis do show clinically relevant benefits, but these may be due to placebo effects.^[99] Acupuncture does not seem to produce long-term benefits.^[100] While electrostimulation techniques such as TENS have been used for twenty years to treat osteoarthritis in the knee, there is no conclusive evidence to show that it reduces pain or disability.^[101]

A Cochrane review of low level laser therapy found unclear evidence of benefit.^[102] Another review found short term pain relief for osteoarthritic knees.^[103]

6 Epidemiology



Disability-adjusted life year for OA per 100,000 inhabitants in 2004.^[104]

Globally approximately 250 million people have osteoarthritis of the knee (3.6% of the population).^[105] OA affects nearly 27 million people in the United States, accounting for 25% of visits to primary care physicians, and half of all NSAID prescriptions. It is estimated that 80% of the population have radiographic evidence

of OA by age 65, although only 60% of those will have symptoms.^[106]

As of 2004, OA globally causes moderate to severe disability in 43.4 million people.^[107]

In the United States, there were approximately 964,000 hospitalizations for osteoarthritis in 2011, a rate of 31 stays per 10,000 population.^[108] With an aggregate cost of \$14.8 billion (\$15,400 per stay), it was the second-most expensive condition seen in U.S. hospital stays in 2011. By payer, it was the second-most costly condition billed to Medicare and private insurance.^{[109][110]}

7 History

Evidence for OA found in the fossil record is studied by paleopathologists, specialists in ancient disease and injury. OA has been reported in fossils of the large carnivorous dinosaur *Allosaurus fragilis*.^[111]

7.1 Etymology

OA is derived from the Greek word part *osteo-*, meaning “of the bone”, combined with *arthritis*: *arthr-*, meaning “joint”, and *-itis*, the meaning of which has come to be associated with inflammation.^[112] The *-itis* of OA could be considered misleading as inflammation is not a conspicuous feature. Some clinicians refer to this condition as *osteoarthrosis* to signify the lack of inflammatory response.

8 Research

There are ongoing efforts to determine if there are agents that modify outcomes in OA. Sprifermin is one candidate drug. There is also tentative evidence that strontium ranelate may decrease degeneration in OA and improve outcomes.^{[113][114]}

As well as attempting to find disease-modifying agents for OA, there is emerging evidence that a system-based approach is necessary to find the causes of OA.^[115] Changes may occur before clinical disease is evident due to abnormalities in biomechanics, biology and/or structure of joints that predispose them to develop clinical disease. Research is thus focusing on defining these early pre-OA changes using biological, mechanical, and imaging markers of OA risk, emphasising multi-disciplinary approaches, and looking into personalized interventions that can reverse OA risk in healthy joints before the disease becomes evident.

Gene transfer strategies aim to target the disease process rather than the symptoms.^[116]

8.1 Biomarkers

Guidelines outlining requirements for inclusion of soluble **biomarkers** in OA clinical trials were published in 2015,^[117] but as yet, there are no validated **biomarkers** for OA. A 2015 systematic review of **biomarkers** for OA looking for molecules that could be used for risk assessments found 37 different biochemical markers of **bone** and **cartilage** turnover in 25 publications.^[118] The strongest evidence was for urinary C-terminal **telopeptide** of **collagen type II** (uCTX-II) as a prognostic marker for knee OA progression and serum cartilage oligomeric protein (**COMP**) levels as a prognostic marker for incidence of both knee and hip OA. A review of biomarkers in hip OA also found associations with uCTXII.^[119]

One problem with using a specific **collagen type II** **biomarker** from the breakdown of articular cartilage is that the amount of cartilage is reduced (worn away) over time with progression of the disease so a patient can eventually have very advanced OA with none of this **biomarker** detectable in their **urine**. Another problem with a systemic **biomarker** is that a patient can have OA in multiple joints at different stages of disease at the same time, so the **biomarker** source cannot be determined. Some other **collagen** breakdown products in the **synovial fluid** correlated with each other after acute injuries (a known cause of secondary OA) but did not correlate with the severity of the injury.^[120]

9 References

- [1] *Atlas of Osteoarthritis*. Springer. 2015. p. 21. ISBN 9781910315163.
- [2] "Osteoarthritis". *National Institute of Arthritis and Musculoskeletal and Skin Diseases*. April 2015. Retrieved 13 May 2015.
- [3] Glyn-Jones, S; Palmer, AJ; Agricola, R; Price, AJ; Vincent, TL; Weinans, H; Carr, AJ (3 March 2015). "Osteoarthritis". *Lancet* **386**: 376–87. doi:10.1016/S0140-6736(14)60802-3. PMID 25748615.
- [4] Berenbaum F (2013). "Osteoarthritis as an inflammatory disease (osteoarthritis is not osteoarthrosis!)". *Osteoarthritis and Cartilage* **21** (1): 16–21. doi:10.1016/j.joca.2012.11.012. PMID 23194896.
- [5] Conaghan P (2014). "Osteoarthritis — Care and management in adults" (PDF).
- [6] March, L; Smith, EU; Hoy, DG; Cross, MJ; Sanchez-Riera, L; Blyth, F; Buchbinder, R; Vos, T; Woolf, AD (June 2014). "Burden of disability due to musculoskeletal (MSK) disorders.". *Best practice & research. Clinical rheumatology* **28** (3): 353–66. doi:10.1016/j.berh.2014.08.002. PMID 25481420.
- [7] Elsternwick (2013). "A problem worth solving.". *Arthritis and Osteoporosis Victoria*.
- [8] MedlinePlus Encyclopedia *Osteoarthritis*
- [9] de Figueiredo EC, Figueiredo GC, Dantas RT (December 2011). "Influência de elementos meteorológicos na dor de pacientes com osteoartrite: Revisão da literatura" [Influence of meteorological elements on osteoarthritis pain: a review of the literature]. *Rev Bras Reumatol* (in Portuguese) **51** (6): 622–8. doi:10.1590/S0482-50042011000600008. PMID 22124595.
- [10] "Water on the knee". MayoClinic.com.
- [11] Brandt KD, Dieppe P, Radin E (January 2009). "Etiopathogenesis of osteoarthritis". *Med. Clin. North Am.* **93** (1): 1–24, xv. doi:10.1016/j.mcna.2008.08.009. PMID 19059018.
- [12] Bosomworth NJ (September 2009). "Exercise and knee osteoarthritis: benefit or hazard?". *Can Fam Physician* **55** (9): 871–8. PMC 2743580. PMID 19752252.
- [13] Deweber K, Olszewski M, Ortolano R (2011). "Knuckle cracking and hand osteoarthritis". *J Am Board Fam Med* **24** (2): 169–74. doi:10.3122/jabfm.2011.02.100156. PMID 21383216.
- [14] Valdes AM, Spector TD (August 2008). "The contribution of genes to osteoarthritis". *Rheum. Dis. Clin. North Am.* **34** (3): 581–603. doi:10.1016/j.rdc.2008.04.008. PMID 18687274.
- [15] Spector TD, MacGregor AJ (2004). "Risk factors for osteoarthritis: genetics". *Osteoarthr. Cartil.* 12 Suppl A: S39–44. doi:10.1016/j.joca.2003.09.005. PMID 14698640.
- [16] Hogervorst T, Bouma HW, de Vos J (August 2009). "Evolution of the hip and pelvis.". *Acta orthopaedica. Supplementum* **80** (336): 1–39. doi:10.1080/17453690610046620. PMID 19919389.
- [17] van der Kraan PM, van den Berg WB (April 2008). "Osteoarthritis in the context of ageing and evolution. Loss of chondrocyte differentiation block during ageing.". *Ageing Research Reviews* **7** (2): 106–13. doi:10.1016/j.arr.2007.10.001. PMID 18054526.
- [18] Coggon D, Reading I, Croft P, McLaren M, Barrett D, Cooper C (May 2001). "Knee osteoarthritis and obesity". *Int. J. Obes. Relat. Metab. Disord.* **25** (5): 622–7. doi:10.1038/sj.ijo.0801585. PMID 11360143.
- [19] Pottie P, Presle N, Terlain B, Netter P, Mainard D, Berenbaum F (November 2006). "Obesity and osteoarthritis: more complex than predicted!". *Ann. Rheum. Dis.* **65** (11): 1403–5. doi:10.1136/ard.2006.061994. PMC 1798356. PMID 17038451.
- [20] Linn S, Murtaugh B, Casey E (May 2012). "Role of sex hormones in the development of osteoarthritis". *PM R* **4** (5 Suppl): S169–73. doi:10.1016/j.pmrj.2012.01.013. PMID 22632696.
- [21] Tanamas SK, Wijethilake P, Wluka AE, Davies-Tuck ML, Urquhart DM, Wang Y, Cicuttini FM (June 2011). "Sex hormones and structural changes in osteoarthritis: a systematic review". *Maturitas* **69** (2): 141–56. doi:10.1016/j.maturitas.2011.03.019. PMID 21481553.

- [22] Ma HL, Blanchet TJ, Peluso D, Hopkins B, Morris EA, Glasson SS (June 2007). "Osteoarthritis severity is sex dependent in a surgical mouse model". *Osteoarthr. Cartil.* **15** (6): 695–700. doi:10.1016/j.joca.2006.11.005. PMID 17207643.
- [23] King KB, Rosenthal AK (2015). "The adverse effects of diabetes on osteoarthritis: update on clinical evidence and molecular mechanisms". *Osteoarthritis Cartilage* **23** (6): 841–50. doi:10.1016/j.joca.2015.03.031. PMID 25837996.
- [24] "OpenStax CNX". *cnx.org*. Retrieved 2015-10-14.
- [25] Maroudas AI (April 1976). "Balance between swelling pressure and collagen tension in normal and degenerate cartilage". *Nature* **260** (5554): 808–9. doi:10.1038/260808a0. PMID 1264261.
- [26] Bollet AJ, Nance JL (July 1966). "Biochemical Findings in Normal and Osteoarthritic Articular Cartilage. II. Chondroitin Sulfate Concentration and Chain Length, Water, and Ash Content". *J. Clin. Invest.* **45** (7): 1170–7. doi:10.1172/JCI105423. PMC 292789. PMID 16695915.
- [27] Brocklehurst R, Bayliss MT, Maroudas A, Coysh HL, Freeman MA, Revell PA, Ali SY (January 1984). "The composition of normal and osteoarthritic articular cartilage from human knee joints. With special reference to unicompartmental replacement and osteotomy of the knee". *J Bone Joint Surg Am* **66** (1): 95–106. PMID 6690447.
- [28] Chou MC, Tsai PH, Huang GS, Lee HS, Lee CH, Lin MH, Lin CY, Chung HW (April 2009). "Correlation between the MR T2 value at 4.7 T and relative water content in articular cartilage in experimental osteoarthritis induced by ACL transection". *Osteoarthr. Cartil.* **17** (4): 441–7. doi:10.1016/j.joca.2008.09.009. PMID 18990590.
- [29] Grushko G, Schneiderman R, Maroudas A (1989). "Some biochemical and biophysical parameters for the study of the pathogenesis of osteoarthritis: a comparison between the processes of ageing and degeneration in human hip cartilage". *Connect. Tissue Res.* **19** (2–4): 149–76. doi:10.3109/03008208909043895. PMID 2805680.
- [30] Mankin HJ, Thrasher AZ (January 1975). "Water content and binding in normal and osteoarthritic human cartilage". *J Bone Joint Surg Am* **57** (1): 76–80. PMID 1123375.
- [31] Venn M, Maroudas A (April 1977). "Chemical composition and swelling of normal and osteoarthrotic femoral head cartilage. I. Chemical composition". *Ann. Rheum. Dis.* **36** (2): 121–9. doi:10.1136/ard.36.2.121. PMC 1006646. PMID 856064.
- [32] Madry H, Luyten FP, Facchini A (2012). "Biological aspects of early osteoarthritis". *Knee Surg. Sports Traumatol. Arthrosc.* **20** (3): 407–22. doi:10.1007/s00167-011-1705-8. PMID 22009557.
- [33] Englund M, Roemer FW, Hayashi D, Crema MD, Guermazi A (2012). "Meniscus pathology, osteoarthritis and the treatment controversy". *Nat. Rev. Rheumatol.* **8** (7): 412–9. doi:10.1038/nrrheum.2012.69. PMID 22614907.
- [34] Li G, Yin J, Gao J, Cheng TS, Pavlos NJ, Zhang C, Zheng MH (2013). "Subchondral bone in osteoarthritis: insight into risk factors and microstructural changes". *Arthritis Research & Therapy* **15** (6): 223. doi:10.1186/ar4405. PMID 24321104.
- [35] Hill CL, Gale DG, Chaisson CE, Skinner K, Kazis L, Gale ME, Felson DT (2001). "Knee effusions, popliteal cysts, and synovial thickening: association with knee pain in osteoarthritis". *J. Rheumatol.* **28** (6): 1330–7. PMID 11409127.
- [36] Felson DT, Chaisson CE, Hill CL, Totterman SM, Gale ME, Skinner KM, Kazis L, Gale DR (2001). "The association of bone marrow lesions with pain in knee osteoarthritis". *Ann Intern Med.* 2001 Apr 3;134(7):541-9 **134** (7): 541–9. doi:10.7326/0003-4819-134-7-200104030-00007. PMID 11281736.
- [37] Zhang W, Doherty M, Peat G, Bierma-Zeinstra MA, Arden NK, Bresnihan B, Herrero-Beaumont G, Kirschner S, Leeb BF, Lohmander LS, Mazières B, Pavelka K, Punzi L, So AK, Tuncer T, Watt I, Bijlsma JW (March 2010). "EULAR evidence-based recommendations for the diagnosis of knee osteoarthritis". *Ann. Rheum. Dis.* **69** (3): 483–9. doi:10.1136/ard.2009.113100. PMID 19762361.
- [38] Bierma-Zeinstra SM, Oster JD, Bernsen RM, Verhaar JA, Ginai AZ, Bohnen AM (August 2002). "Joint space narrowing and relationship with symptoms and signs in adults consulting for hip pain in primary care". *J. Rheumatol.* **29** (8): 1713–8. PMID 12180735.
- [39] *Osteoarthritis (OA): Joint Disorders* at Merck Manual of Diagnosis and Therapy Professional Edition
- [40] Phillips CR, Brasington RD (2010). "Osteoarthritis treatment update: Are NSAIDs still in the picture?". *Journal of Musculoskeletal Medicine* **27** (2).
- [41] Kalunian KC (2013). "Patient information: Osteoarthritis symptoms and diagnosis (Beyond the Basics)". UpToDate. Retrieved 15 February 2013.
- [42] Altman R, Alarcón G, Appelrouth D, Bloch D, Borenstein D, Brandt K, Brown C, Cooke TD, Daniel W, Gray R (November 1990). "The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the hand". *Arthritis Rheum.* **33** (11): 1601–10. doi:10.1002/art.1780331101. PMID 2242058.
- [43] Vasan N, Tao L, Vikas B (2010). *First Aid for the USMLE Step 1, 2010 (First Aid USMLE)*. McGraw-Hill Medical. p. 378. ISBN 0-07-163340-5.
- [44] Punzi L, Ramonda R, Sfriso P (October 2004). "Erosive osteoarthritis". *Best Pract Res Clin Rheumatol* **18** (5): 739–58. doi:10.1016/j.berh.2004.05.010. PMID 15454130.
- [45] Flood J (March 2010). "The role of acetaminophen in the treatment of osteoarthritis". *Am J Manag Care* **16** (Suppl Management): S48–54. PMID 20297877.

- [46] Cibulka MT, White DM, Woehrle J, Harris-Hayes M, Ensey K, Fagerson TL, Slover J, Godges JJ (April 2009). "Hip pain and mobility deficits—hip osteoarthritis: clinical practice guidelines linked to the international classification of functioning, disability, and health from the orthopaedic section of the American Physical Therapy Association". *J Orthop Sports Phys Ther* **39** (4): A1–25. doi:10.2519/jospt.2009.0301. PMID 19352008.
- [47] Hagen KB, Dagfinrud H, Moe RH, Østerås N, Kjekshus I, Grotle M, Smedslund G (2012). "Exercise therapy for bone and muscle health: an overview of systematic reviews". *BMC Med* **10**: 167. doi:10.1186/1741-7015-10-167. PMC 3568719. PMID 23253613.
- [48] Fransen M, McConnell S, Hernandez-Molina G, Reichenbach S (2014). "Exercise for osteoarthritis of the hip". *Cochrane Database Syst Rev* **4** (4): CD007912. doi:10.1002/14651858.CD007912.pub2. PMID 24756895.
- [49] Juhl C, Christensen R, Roos EM, Zhang W, Lund H (Mar 2014). "Impact of exercise type and dose on pain and disability in knee osteoarthritis: a systematic review and meta-regression analysis of randomized controlled trials". *Arthritis & rheumatology (Hoboken, N.J.)* **66** (3): 622–36. doi:10.1002/art.38290. PMID 24574223.
- [50] Wang SY, Olson-Kellogg B, Shamliyan TA, Choi JY, Ramakrishnan R, Kane RL (November 2012). "Physical therapy interventions for knee pain secondary to osteoarthritis: a systematic review". *Annals of Internal Medicine* **157** (9): 632–44. doi:10.7326/0003-4819-157-9-201211060-00007. PMID 23128863.
- [51] De Luigi AJ (May 2012). "Complementary and alternative medicine in osteoarthritis". *PM R* **4** (5 Suppl): S122–33. doi:10.1016/j.pmrj.2012.01.012. PMID 22632691.
- [52] French HP, Brennan A, White B, Cusack T (2011). "Manual therapy for osteoarthritis of the hip or knee — a systematic review". *Man Ther* **16** (2): 109–117. doi:10.1016/j.math.2010.10.011. PMID 21146444.
- [53] Sturme DL, Tiedemann A, Chapman K, Munro B, Murray SM, Lord SR (November 2004). "Physiological risk factors for falls in older people with lower limb arthritis". *J. Rheumatol.* **31** (11): 2272–9. PMID 15517643.
- [54] Penny P, Geere J, Smith TO (October 2013). "A systematic review investigating the efficacy of laterally wedged insoles for medial knee osteoarthritis". *Rheumatol. Int.* **33** (10): 2529–38. doi:10.1007/s00296-013-2760-x. PMID 23612781.
- [55] Parkes MJ, Maricar N, Lunt M, LaValley MP, Jones RK, Segal NA, Takahashi-Narita K, Felson DT (August 2013). "Lateral wedge insoles as a conservative treatment for pain in patients with medial knee osteoarthritis: a meta-analysis". *JAMA* **310** (7): 722–30. doi:10.1001/jama.2013.243229. PMID 23989797.
- [56] Page CJ, Hinman RS, Bennell KL (2011). "Physiotherapy management of knee osteoarthritis". *Int J Rheum Dis* **14** (2): 145–152. doi:10.1111/j.1756-185X.2011.01612.x. PMID 21518313.
- [57] "Osteoarthritis Lifestyle and home remedies". *Diseases and Conditions*. Mayo Clinic.
- [58] Consumer Reports Health Best Buy Drugs (July 2013), "The Nonsteroidal Anti-Inflammatory Drugs: Treating Osteoarthritis and Pain. Comparing effectiveness, safety, and price." (PDF), *NSAIDs*, Yonkers, New York: Consumer Reports, retrieved 12 February 2014
- [59] Zhang W, Moskowitz RW, Nuki G, Abramson S, Altman RD, Arden N, Bierma-Zeinstra S, Brandt KD, Croft P, Doherty M, Dougados M, Hochberg M, Hunter DJ, Kwoh K, Lohmander LS, Tugwell P (September 2007). "OARSI recommendations for the management of hip and knee osteoarthritis, part I: critical appraisal of existing treatment guidelines and systematic review of current research evidence". *Osteoarthr. Cartil.* **15** (9): 981–1000. doi:10.1016/j.joca.2007.06.014. PMID 17719803.
- [60] Machado, GC; Maher, CG; Ferreira, PH; Pinheiro, MB; Lin, CW; Day, RO; McLachlan, AJ; Ferreira, ML (31 March 2015). "Efficacy and safety of paracetamol for spinal pain and osteoarthritis: systematic review and meta-analysis of randomised placebo controlled trials.". *BMJ (Clinical research ed.)* **350**: h1225. doi:10.1136/bmj.h1225. PMID 25828856.
- [61] Chen YF, Jobanputra P, Barton P, Bryan S, Fry-Smith A, Harris G, Taylor RS (April 2008). "Cyclooxygenase-2 selective non-steroidal anti-inflammatory drugs (etodolac, meloxicam, celecoxib, rofecoxib, etoricoxib, valdecoxib and lumiracoxib) for osteoarthritis and rheumatoid arthritis: a systematic review and economic evaluation". *Health Technol Assess* **12** (11): 1–278, iii. doi:10.3310/hta12110. PMID 18405470.
- [62] Wielage, RC; Myers, JA; Klein, RW; Hapich, M (December 2013). "Cost-effectiveness analyses of osteoarthritis oral therapies: a systematic review.". *Applied Health Economics and Health Policy* **11** (6): 593–618. doi:10.1007/s40258-013-0061-x. PMID 24214160.
- [63] McAlindon, TE; Bannuru, RR; Sullivan, MC; Arden, NK; Berenbaum, F; Bierma-Zeinstra, SM; Hawker, GA; Henrotin, Y; Hunter, DJ; Kawaguchi, H; Kwoh, K; Lohmander, S; Rannou, F; Roos, EM; Underwood, M (March 2014). "OARSI guidelines for the non-surgical management of knee osteoarthritis.". *Osteoarthritis and cartilage / OARS, Osteoarthritis Research Society* **22** (3): 363–88. doi:10.1016/j.joca.2014.01.003. PMID 24462672.
- [64] Hochberg, MC; Altman, RD; April, KT; Benkhalti, M; Guyatt, G; McGowan, J; Towheed, T; Welch, V; Wells, G; Tugwell, P; American College of, Rheumatology (April 2012). "American College of Rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee.". *Arthritis care & research* **64** (4): 465–74. doi:10.1002/acr.21596. PMID 22563589.
- [65] da Costa, BR; Nüesch, E; Kasteler, R; Husni, E; Welch, V; Rutjes, AW; Jüni, P (17 September 2014). "Oral or transdermal opioids for osteoarthritis of the knee or hip.". *The Cochrane database of systematic reviews* **9**: CD003115. doi:10.1002/14651858.CD003115.pub4. PMID 25229835.

- [66] Derry S, Moore RA, Rabbie R (2012). "Topical NSAIDs for chronic musculoskeletal pain in adults". *The Cochrane Library* **9**: CD007400. doi:10.1002/14651858.CD007400.pub2. PMID 22972108.
- [67] da Costa BR, Nüesch E, Kasteler R, Husni E, Welch V, Rutjes AW, Jüni P (2014). "Oral or transdermal opioids for osteoarthritis of the knee or hip.". *The Cochrane database of systematic reviews* **9**: CD003115. doi:10.1002/14651858.CD003115.pub4. PMID 25229835.
- [68] De Silva V, El-Metwally A, Ernst E, Lewith G, Macfarlane GJ (May 2011). "Evidence for the efficacy of complementary and alternative medicines in the management of osteoarthritis: a systematic review". *Rheumatology (Oxford)* **50** (5): 911–20. doi:10.1093/rheumatology/keq379. PMID 21169345.
- [69] Cameron M, Gagnier JJ, Little CV, Parsons TJ, Blümle A, Chrubasik S (November 2009). "Evidence of effectiveness of herbal medicinal products in the treatment of arthritis. Part I: Osteoarthritis". *Phytother Res* **23** (11): 1497–515. doi:10.1002/ptr.3007. PMID 19856319.
- [70] Altman R, Barkin RL (March 2009). "Topical therapy for osteoarthritis: clinical and pharmacologic perspectives". *Postgrad Med* **121** (2): 139–47. doi:10.3810/pgm.2009.03.1986. PMID 19332972.
- [71] Arroll B, Goodyear-Smith F (April 2004). "Corticosteroid injections for osteoarthritis of the knee: meta-analysis". *BMJ* **328** (7444): 869. doi:10.1136/bmj.38039.573970.7C. PMC 387479. PMID 15039276.
- [72] Colen S; van den Bekerom, MP; Mulier, M; Haverkamp, D (1 August 2012). "Hyaluronic acid in the treatment of knee osteoarthritis: a systematic review and meta-analysis with emphasis on the efficacy of different products.". *BioDrugs : clinical immunotherapeutics, biopharmaceuticals and gene therapy* **26** (4): 257–68. doi:10.2165/11632580-000000000-00000. PMID 22734561.
- [73] Rutjes AW, Jüni P, da Costa BR, Trelle S, Nüesch E, Reichenbach S (August 2012). "Viscosupplementation for osteoarthritis of the knee: a systematic review and meta-analysis". *Annals of Internal Medicine* **157** (3): 180–91. doi:10.7326/0003-4819-157-3-201208070-00473. PMID 22868835.
- [74] Khoshbin A, Leroux T, Wasserstein D, Marks P, Theodoropoulos J, Ogilvie-Harris D, Gandhi R, Takhar K, Lum G, Chahal J (December 2013). "The efficacy of platelet-rich plasma in the treatment of symptomatic knee osteoarthritis: a systematic review with quantitative synthesis". *Arthroscopy* **29** (12): 2037–48. doi:10.1016/j.arthro.2013.09.006. PMID 24286802.
- [75] Rodriguez-Merchan, EC (September 2013). "Intraarticular Injections of Platelet-rich Plasma (PRP) in the Management of Knee Osteoarthritis". *Archives of bone and joint surgery* **1** (1): 5–8. PMC 4151401. PMID 25207275.
- [76] Santaguida PL, Hawker GA, Hudak PL, Glazier R, Mahomed NN, Kreder HJ, Coyte PC, Wright JG (December 2008). "Patient characteristics affecting the prognosis of total hip and knee joint arthroplasty: a systematic review". *Can J Surg* **51** (6): 428–36. PMC 2592576. PMID 19057730.
- [77] Carr AJ, Robertsson O, Graves S, Price AJ, Arden NK, Judge A, Beard DJ (April 2012). "Knee replacement". *Lancet* **379** (9823): 1331–40. doi:10.1016/S0140-6736(11)60752-6. PMID 22398175.
- [78] Jenkins PJ, Clement ND, Hamilton DF, Gaston P, Patton JT, Howie CR (2013). "Predicting the cost-effectiveness of total hip and knee replacement: A health economic analysis". *The bone & joint journal* **95-B** (1): 115–21. doi:10.1302/0301-620X.95B1.29835. PMID 23307684.
- [79] Daigle ME, Weinstein AM, Katz JN, Losina E (2012). "The cost-effectiveness of total joint arthroplasty: A systematic review of published literature". *Best practice & research. Clinical rheumatology* **26** (5): 649–58. doi:10.1016/j.berh.2012.07.013. PMC 3879923. PMID 23218429.
- [80] Hunziker EB, Lippuner K, Keel MJ, Shintani N (2015). "An educational review of cartilage repair: precepts & practice – myths & misconceptions – progress & prospects". *Osteoarthritis Cartilage* **23** (3): 334–50. doi:10.1016/j.joca.2014.12.011. PMID 25534362.
- [81] Brouwer, RW; Huizinga, MR; Duivenvoorden, T; van Raaij, TM; Verhagen, AP; Bierma-Zeinstra, SM; Verhaar, JA (13 December 2014). "Osteotomy for treating knee osteoarthritis.". *The Cochrane database of systematic reviews* **12**: CD004019. doi:10.1002/14651858.CD004019.pub4. PMID 25503775.
- [82] Nelson AE, Allen KD, Golightly YM, Goode AP, Jordan JM (2014). "A systematic review of recommendations and guidelines for the management of osteoarthritis: The Chronic Osteoarthritis Management Initiative of the U.S. Bone and Joint Initiative". *Seminars in arthritis and rheumatism* **43** (6): 701–12. doi:10.1016/j.semarthrit.2013.11.012. PMID 24387819.
- [83] Katz, JN; Brownlee, SA; Jones, MH (February 2014). "The role of arthroscopy in the management of knee osteoarthritis.". *Best practice & research. Clinical rheumatology* **28** (1): 143–56. doi:10.1016/j.berh.2014.01.008. PMID 24792949.
- [84] Thorlund, JB.; Juhl, CB.; Roos, EM.; Lohmander, LS. (2015). "Arthroscopic surgery for degenerative knee: systematic review and meta-analysis of benefits and harms". *BMJ* **350**: h2747. doi:10.1136/bmj.h2747. PMC 4469973. PMID 26080045.
- [85] The effects of Glucosamine Sulphate on OA of the knee joint. BestBets.
- [86] Burdett N, McNeil JD (Sep 2012). "Difficulties with assessing the benefit of glucosamine sulphate as a treatment for osteoarthritis.". *International journal of evidence-based healthcare* **10** (3): 222–6. doi:10.1111/j.1744-1609.2012.00279.x. PMID 22925619.

- [87] Wandel S, Jüni P, Tendal B, Nüesch E, Villiger PM, Welton NJ, Reichenbach S, Trelle S (Sep 16, 2010). "Effects of glucosamine, chondroitin, or placebo in patients with osteoarthritis of hip or knee: network meta-analysis.". *BMJ (Clinical research ed.)* **341**: c4675. doi:10.1136/bmj.c4675. PMC 2941572. PMID 20847017.
- [88] Wu D, Huang Y, Gu Y, Fan W (Jun 2013). "Efficacies of different preparations of glucosamine for the treatment of osteoarthritis: a meta-analysis of randomised, double-blind, placebo-controlled trials.". *International journal of clinical practice* **67** (6): 585–94. doi:10.1111/ijcp.12115. PMID 23679910.
- [89] Chou R, McDonagh MS, Nakamoto E, Griffin J (Oct 2011). "Analgesics for Osteoarthritis: An Update of the 2006 Comparative Effectiveness Review". PMID 22091473.
- [90] Miller KL, Clegg DO (February 2011). "Glucosamine and chondroitin sulfate". *Rheum. Dis. Clin. North Am.* **37** (1): 103–18. doi:10.1016/j.rdc.2010.11.007. PMID 21220090. The best current evidence suggests that the effect of these supplements, alone or in combination, on OA pain, function, and radiographic change is marginal at best.
- [91] Rovati LC, Girolami F, Persiani S (Jun 2012). "Crystalline glucosamine sulfate in the management of knee osteoarthritis: efficacy, safety, and pharmacokinetic properties.". *Therapeutic advances in musculoskeletal disease* **4** (3): 167–80. doi:10.1177/1759720X12437753. PMC 3400104. PMID 22850875.
- [92] Gregory, PJ; Fellner, C (June 2014). "Dietary supplements as disease-modifying treatments in osteoarthritis: a critical appraisal.". *P & T : a peer-reviewed journal for formulary management* **39** (6): 436–52. PMID 25050057.
- [93] Zhang W, Moskowitz RW, Nuki G, Abramson S, Altman RD, Arden N, Bierma-Zeinstra S, Brandt KD, Croft P, Doherty M, Dougados M, Hochberg M, Hunter DJ, Kwoh K, Lohmander LS, Tugwell P (February 2008). "OARSI recommendations for the management of hip and knee osteoarthritis, Part II: OARSI evidence-based, expert consensus guidelines" (PDF). *Osteoarthr. Cartil.* **16** (2): 137–62. doi:10.1016/j.joca.2007.12.013. PMID 18279766. Archived from the original (PDF) on July 21, 2011.
- [94] Henrotin Y, Mobasheri A, Marty M (Jan 30, 2012). "Is there any scientific evidence for the use of glucosamine in the management of human osteoarthritis?". *Arthritis Research & Therapy* **14** (1): 201. doi:10.1186/ar3657. PMC 3392795. PMID 22293240.
- [95] Lopez HL (May 2012). "Nutritional interventions to prevent and treat osteoarthritis. Part II: focus on micronutrients and supportive nutraceuticals". *PM & R : the journal of injury, function, and rehabilitation* **4** (5 Suppl): S155–68. doi:10.1016/j.pmrj.2012.02.023. PMID 22632695.
- [96] Rosenbaum CC, O'Mathúna DP, Chavez M, Shields K (2010). "Antioxidants and antiinflammatory dietary supplements for osteoarthritis and rheumatoid arthritis". *Altern Ther Health Med* **16** (2): 32–40. PMID 20232616.
- [97] Pirotta M (September 2010). "Arthritis disease - the use of complementary therapies". *Aust Fam Physician* **39** (9): 638–40. PMID 20877766.
- [98] Cameron M, Chrubasik S (May 22, 2014). "Oral herbal therapies for treating osteoarthritis". *The Cochrane database of systematic reviews* **5**: CD002947. doi:10.1002/14651858.cd002947.pub2. PMC 4494689. PMID 24848732. Several other medicinal plant products, including extracts of *Boswellia serrata*, show trends of benefits that warrant further investigation in light of the fact that the risk of adverse events appear low.
- [99] Manheimer E, Cheng K, Linde K, Lao L, Yoo J, Wieland S, van der Windt DA, Berman BM, Bouter LM (2010). Manheimer, Eric, ed. "Acupuncture for peripheral joint osteoarthritis". *Cochrane Database of Systematic Reviews* (1): CD001977. doi:10.1002/14651858.CD001977.pub2. PMC 3169099. PMID 20091527.
- [100] Wang SM, Kain ZN, White PF (February 2008). "Acupuncture analgesia: II. Clinical considerations" (PDF). *Anesth. Analg.* **106** (2): 611–21, table of contents. doi:10.1213/ane.0b013e318160644d. PMID 18227323.
- [101] Rutjes AW, Nüesch E, Sterchi R, Kalichman L, Hendriks E, Osiri M, Brosseau L, Reichenbach S, Jüni P (2009). Rutjes, Anne WS, ed. "Transcutaneous electrostimulation for osteoarthritis of the knee". *Cochrane Database of Systematic Reviews* (4): CD002823. doi:10.1002/14651858.CD002823.pub2. PMID 19821296.
- [102] Brosseau, L; Welch, V; Wells, G; DeBie, R; Gam, A; Harman, K; Morin, M; Shea, B; Tugwell, P (2004). "Low level laser therapy (Classes I, II and III) for treating osteoarthritis.". *The Cochrane database of systematic reviews* (3): CD002046. doi:10.1002/14651858.CD002046.pub2. PMID 15266461.
- [103] Bjordal, J; Johnson, M; Lopes-Martins, R; Bogen, B; Chow, R; Ljunggren, A (2007). "Short-term efficacy of physical interventions in osteoarthritic knee pain. A systematic review and meta-analysis of randomised placebo-controlled trials.". *BMC Musculoskeletal Disorders* **8** (8): 51. doi:10.1186/1471-2474-8-51.
- [104] "WHO Disease and injury country estimates". *World Health Organization*. 2009. Retrieved Nov 11, 2009.
- [105] Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, et al. (December 2012). "Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010". *Lancet* **380** (9859): 2163–96. doi:10.1016/S0140-6736(12)61729-2. PMID 23245607.
- [106] Green GA (2001). "Understanding NSAIDs: from aspirin to COX-2". *Clin Cornerstone* **3** (5): 50–60. doi:10.1016/S1098-3597(01)90069-9. PMID 11464731.

- [107] “Table 9: Estimated prevalence of moderate and severe disability (millions) for leading disabling conditions by age, for high-income and low- and middle-income countries, 2004”. *The Global Burden of Disease: 2004 Update*. Geneva: World Health Organization. 2008. p. 35. ISBN 978-92-4-156371-0.
- [108] Pfuntner A., Wier L.M., Stocks C. Most Frequent Conditions in U.S. Hospitals, 2011. HCUP Statistical Brief #162. September 2013. Agency for Healthcare Research and Quality, Rockville, Maryland.
- [109] Torio CM, Andrews RM (August 2013). “National Inpatient Hospital Costs: The Most Expensive Conditions by Payer, 2011.”. *HCUP Statistical Brief #160*. Rockville, Maryland: Agency for Healthcare Research and Quality.
- [110] Pfuntner, A; Wier, L. M.; Steiner, C (December 2013). “Costs for Hospital Stays in the United States, 2011: Statistical Brief #168”. PMID 24455786.
- [111] Molnar, R. E. (2001). “Therapod Paleopathology: A Literature Survey”. In Tanke, Darren H.; Carpenter, Kenneth; Skrepnick, Michael William. *Mesozoic Vertebrate Life*. Indiana University Press. pp. 337–63. ISBN 978-0-253-33907-2.
- [112] Devaraj TL (2011). “Chapter 41: Nature cure yoga for osteoarthritis”. *Nature Cure for Common Diseases*. New Delhi: Arya Publication. p. 368. ISBN 978-8189093747.
- [113] Civjan N (2012). *Chemical Biology: Approaches to Drug Discovery and Development to Targeting Disease*. John Wiley & Sons. p. 313. ISBN 9781118437674.
- [114] Bruyère O, Burlet N, Delmas PD, Rizzoli R, Cooper C, Reginster JY (2008). “Evaluation of symptomatic slow-acting drugs in osteoarthritis using the GRADE system”. *BMC Musculoskelet Disord* **9**: 165. doi:10.1186/1471-2474-9-165. PMC 2627841. PMID 19087296.
- [115] Chu, CR; Andriacchi, TP (2015). “Dance between biology, mechanics, and structure: a systems-based approach to developing osteoarthritis prevention strategies”. *J Orthop Res* **33** (7): 939–947. doi:10.1002/jor.22817. PMID 25639920.
- [116] T. Pap, J. Schedel, G. Pap, U. Moller-Ladner, R.E. Gay, S. Gay C. Guincamp (2000). “Gene therapy in osteoarthritis”. *Joint Bone Spine*.
- [117] Kraus, VB; Blanco, FJ; Englund, M; Henrotin, Y; Lohmander, LS; Losina, E; Onnerfjord, P; Persiani, S (2015). “OARSI Clinical Trials Recommendations: Soluble biomarker assessments in clinical trials in osteoarthritis”. *Osteoarthritis Cartilage* **23** (5): 686–697. doi:10.1016/j.joca.2015.03.002. PMC 4430113. PMID 25952342.
- [118] Saberi Hosnijeh, F; Runhaar, J; van Meurs, JB; Bierma-Zeinstra, SM (2015). “Biomarkers for osteoarthritis: Can they be used for risk assessment? A systematic review”. *Maturitas* **82** (1): 36–49. doi:10.1016/j.maturitas.2015.04.004. PMID 25963100.
- [119] Nepple, JJ; Thomason, KM; An, TW; Harris-Hayes, M; Clohisy, JC (2015). “What is the utility of biomarkers for assessing the pathophysiology of hip osteoarthritis? A systematic review”. *Clin Orthop Relat Res* **473** (5): 1683–1701. doi:10.1007/s11999-015-4148-6. PMID 25623593.
- [120] Kumahashi, N; Swärd, P; Larsson, S; Lohmander, LS; Frobell, R; Struglics, A (2015). “Type II collagen C2C epitope in human synovial fluid and serum after knee injury - associations with molecular and structural markers of injury”. *Osteoarthritis Cartilage* **23** (9): 1506–12. doi:10.1016/j.joca.2015.04.022. PMID 25937025.

10 External links

- American College of Rheumatology Factsheet on OA
- Osteoarthritis The Arthritis Foundation
- National Institute of Arthritis and Musculoskeletal and Skin Diseases - US National Institute of Arthritis and Musculoskeletal and Skin Diseases

11 Text and image sources, contributors, and licenses

11.1 Text

- **Osteoarthritis** *Source:* <https://en.wikipedia.org/wiki/Osteoarthritis?oldid=698151159> *Contributors:* Damian Yerrick, Alex.tan, Edward, Madvenu, Gabbe, Sannse, Arpingstone, Ellywa, Andrewman327, IceKarma, Tpbradbury, Robert2957, Robbot, Iaen, Sunray, Mushroom, Lupo, Techelf, Barbara Shack, Bradeos Graphon, Michael Devore, Jfdwolff, Guanaco, Adenosine, Quadell, Scottperry, Bodnotbod, Ropers, TheObtuseAngleOfDoom, Shiftchange, DanielCD, Discospinster, Rich Farmbrough, Petersam, Sfahey, Mwanner, Bobo192, Reinyday, Dreish, Davidruben, Arcadian, Giraffedata, Nsaa, Rernst, Alansohn, Eric Kvaalen, Wouterstomp, Gilgamesfuel, ClockworkSoul, Cromwellt, Cburnett, Scieurinae, Ceyockey, Abanima, Alkarex, Garrison Roo, Woohookitty, Skyraider, Pol098, Nick Drake, Stixpir, Mandarax, Sjakkalle, Rjwilmsi, BirdbrainedPhoenix, Ramalho, RexNL, Stevenfruitsmaak, Travis.Thurston, DVdm, WriterHound, Wavelength, RobotE, Cooke, RussBot, Chris Capoccia, Stephenb, CambridgeBayWeather, Eleassar, Draeco, NawlinWiki, Irishguy, Raven4x4x, PM Poon, Epipelagic, Dbfirs, S. Neuman, Bdel1555, PTSE, Imaninjapirate, Zakksez, Badgettrg, Garion96, Katieh5584, SmackBot, Kast-enwiki, DTM, Edgar181, Cazort, Unforgettableid, PeterSymonds, Gilliam, Ohnoitsjamie, Indium, Marc Kupper, Vvarkey, Kurykh, RDBrown, SchiftyThree, Delink, Uthbrian, Colonies Chris, Coffeeholic!, Coopermd, Salmar, ArthritisCritic, Egsan Bacon, Abyssal, Roseave, Cybercobra, Gregnz, Nakon, James McNally, Salamurai, FelisLeo, Cyberevil, JorisvS, IronGargoyle, 16@r, Beetstra, Uuhwulj, SandyGeorgia, RBJ, Hu12, Paleolith, Iridescent, Frank Lofaro Jr., ChrisCork, Harold f, Slann, Prithason, CmdrObot, Pmyteh, Pajast, Lentower, Maxxicum, Cydebot, Corpx, Hopping, XcepticZP, Evets snivelb, Thijs!bot, Hildre, Mojo Hand, Sartaj beary, Headbomb, Bobblehead, Topnife, Transhumanist, Cyclonenim, Seaphoto, Marokwitz, Vendettax, ChristianBk, Lfstevens, JAnDbot, Wiki0709, MER-C, ByronRigby, Antelan, VoABot II, JamesBWatson, Bwhack, Caesarjbsquitti, Lenihanlover, Npos, Charlenelieu, JaGa, WLU, TimidGuy, FisherQueen, Yobol, MartinBot, Ladarzak, CliffC, Vigyani, Steveblevins, Drwhapcaplet, Dr. St-Amant, J.delanoy, CFCF, Trusilver, Nbaum, Boghog, Surfunc, Ghoshkgb, Lama010101, Michael Daly, Mschamberlain, Lystrablue, Tagus, Jewzip, LoopTel, Lrunge, Jhashemi, Idioma-bot, VolkovBot, CaliforniaLyme, Macedonian, Torswin, Jadevt, Anonymous Dissident, Una Smith, Guest9999, Latulla, Rysbr333, Falcon8765, RGB2, Derekawesome, Doc James, Darxus, Smuggy4, SieBot, Brenont, SheepNotGoats, Gerakibot, BService, LeadSong-Dog, SignalWaveLenght00, Nummer29, PatientAdvocateBJD, Mbgreenblue, Acweb, Macy, OKBot, Tomtchek, Bogwhistle, Wlbowden, Osteoimm, Drgarden, Beemer69, Lekshumi, ClueBot, JamesStewart7, Sthubbar, Rebelyell1916, Benolam, ImperfectlyInformed, RPainter, Sanholding, Mild Bill Hiccup, Celiakozlowski, Beauhanks, Telekenesis, NuclearWarfare, Dekisugi, IForTheMoney, Kendra.gilbert, Hoverflysmiles, Porkman Pork, Arcinfo, Drpawansharma, Homerstrykercenterlibrary, Attaboy, XLinkBot, EastTN, SilvonenBot, Arazai, Fire-Brandon, Klemonehead1001, Addbot, DOI bot, Dr nick s, RCPWilson, Lisa4edit, SnapeLady, Diptanshu.D, MrOllie, Download, Debresser, Ahmad.ghamdi.24, LinkFA-Bot, Craigsjones, Tide rolls, Lightbot, Apteva, ماني, DR66, Shalvata, Noumenon, Luckas-bot, Yobot, DoctorDW, Neilandkathgriff, AnomieBOT, Rubinbot, Ssmithm, Piano non troppo, Tom87020, Sz-iwbot, Blueraspberry, MaterialsScientist, Skygod7, Melaniemalone, Citation bot, Redsoxcool, Jmarchn, ArthurBot, Carturo222, Xqbot, Zad68, Happyrabbit, Flavonoid, Almbot, PRfection, J04n, GrouchoBot, DVMresearcher, Twirligig, Addingrefs, Prunesqualer, Jgalt23, Ajax151, Jbs sumc, Mirubuffy, San Fernando Valley, Afromayun, Dannysfeir, Flatronwide, Contentmaven, Thugmed, Holyangel104, Citation bot 1, Sunlit Forest, Pinethicket, I dream of horses, Fuzbaby, Sctechlaw, Jonesey95, Hiah1974, RedBot, Vicmahajan, Florescent, Full-date unlinking bot, Corinne68, Tim1357, Orenburg1, Trappist the monk, Forge999, Jenday, Fastlysock, Suffusion of Yellow, Onel5969, Mean as custard, RjwilmsiBot, TjBot, WildBot, Elmodude5, EmausBot, Immunize, BillyPreset, RYagihara, Ankush8, StanStun, Amr071291, ZéroBot, MithrandirAgain, KuduIO, Millsysj, Puldis, JoeSperrazza, Targad, Cyberdog958, Ssmithmfr, Whoop whoop pull up, ClueBot NG, DPG1993, Puhlaa, Jameswatts1990, Qwerasdzxcv1234, Rezabot, Widr, JTT Young, Doomguy, Llogan02, Helpful Pixie Bot, Ruskipotter, KLBot2, BG19bot, Krenair, AndrewSiminoff, MusikAnimal, Elucches, Carystus, StevinSimon, Jaysn1, Achowat, LindaFisher7, Fuse809, Ryttyho usa, BattyBot, Biosthorms, Arr4, Abbyquack, Cyberbot II, ChrisGualtieri, TylerDurden8823, JYBot, Dexbot, FoCuSandLeArN, Hmainsbot1, AbstractIllusions, 900mill, Comodny, BeachComber1972, Drway, Krgoria, DavidLeighEllis, OARSI, BruceBlaus, Alice Person, Madelynn Dudas, Anr-nusna, MsLindseyParks, Monkbob, RJM76, Gronk Oz, Lumican, Sweet Orange Juice, Sushi juju, Laundromatzat, Youniszainal, Mikeale Wallace, Sundayclose, Frogilly1, Bobkarlz, Criel44, Mysterious Gopher, KasparBot, Osteomat, Anna.morakot, Unenthusiastic, MelisaK212, Nateshnandha, Carl.martin05, Oliverhutchon, Adhithyan Arulsamy, Nuggets of Knowledge, 61sman16 and Anonymous: 412

11.2 Images

- **File:0910_Oateoarthritis_Hip_A.png** *Source:* https://upload.wikimedia.org/wikipedia/commons/8/81/0910_Oateoarthritis_Hip_A.png *License:* CC BY-SA 4.0 *Contributors:* Own work *Original artist:* CFCF
- **File:0910_Oateoarthritis_Hip_B.png** *Source:* https://upload.wikimedia.org/wikipedia/commons/3/3e/0910_Oateoarthritis_Hip_B.png *License:* CC BY-SA 4.0 *Contributors:* Own work *Original artist:* CFCF
- **File:Areas_affected_by_osteoarthritis.gif** *Source:* https://upload.wikimedia.org/wikipedia/commons/7/74/Areas_affected_by_osteoarthritis.gif *License:* Public domain *Contributors:* http://www.niams.nih.gov/health_info/Osteoarthritis/default.asp *Original artist:* US federal government
- **File:Commons-logo.svg** *Source:* <https://upload.wikimedia.org/wikipedia/en/4/4a/Commons-logo.svg> *License:* ? *Contributors:* ? *Original artist:* ?
- **File:Exercise.png** *Source:* <https://upload.wikimedia.org/wikipedia/commons/4/43/Exercise.png> *License:* Public domain *Contributors:* http://www.niams.nih.gov/health_info/Osteoarthritis/default.asp *Original artist:* US federal government
- **File:Osteoarthritis_world_map_-_DALY_-_WHO2004.svg** *Source:* https://upload.wikimedia.org/wikipedia/commons/b/be/Osteoarthritis_world_map_-_DALY_-_WHO2004.svg *License:* CC BY-SA 2.5 *Contributors:*
- Vector map from BlankMap-World6, compact.svg by Canuckguy et al. *Original artist:* Lokal_Profil
- **File:Secondary_osteoarthritis_Ankle_F.png** *Source:* https://upload.wikimedia.org/wikipedia/commons/6/6e/Secondary_osteoarthritis_Ankle_F.png *License:* CC BY-SA 3.0 *Contributors:* Own work *Original artist:* Jmarchn
- **File:Secondary_osteoarthritis_Ankle_P.png** *Source:* https://upload.wikimedia.org/wikipedia/commons/0/08/Secondary_osteoarthritis_Ankle_P.png *License:* CC BY-SA 3.0 *Contributors:* Own work *Original artist:* Jmarchn

11.3 Content license

- Creative Commons Attribution-Share Alike 3.0