

# EPFL

## Gradient Flows in Wasserstein Spaces

Variational Inference and Sampling



Student: L.Raffo

PostDoc. L.V.Santoro

# motivation

consider the Euclidean space  $\mathbb{R}^d$ .




# motivation

consider the Euclidean space  $\mathbb{R}^d$ .



look at the family of probability measures  $\mathcal{P}$  on  $\mathbb{R}^d$ .




# motivation

consider the Euclidean space  $\mathbb{R}^d$ .



look at the family of probability measures  $\mathcal{P}$  on  $\mathbb{R}^d$ .



define a distance  $\mathcal{W}$  between probability measures.


$$\mathcal{W}(\text{broad curve}, \text{narrow curve})$$

# motivation

consider the Euclidean space  $\mathbb{R}^d$ .



look at the family of probability measures  $\mathcal{P}_2$  on  $\mathbb{R}^d$ .



define a distance  $\mathcal{W}_2$  between probability measures.

$$\mathcal{W}_2(\text{~}\sim\text{~}, \text{~}\wedge\text{~})$$

# motivation

the metric space  $(\mathcal{P}_2, \mathcal{W}_2)$  has nice geometric properties.

# motivation

the metric space  $(\mathcal{P}_2, \mathcal{W}_2)$  has nice geometric properties.

despite being infinite dimensional, it resembles a Riemannian manifold.

# motivation

the metric space  $(\mathcal{P}_2, \mathcal{W}_2)$  has nice geometric properties.

despite being infinite dimensional, it resembles a Riemannian manifold.



geodesics

tangent spaces



# motivation

the metric space  $(\mathcal{P}_2, \mathcal{W}_2)$  has nice geometric properties.

despite being infinite dimensional, it resembles a Riemannian manifold.

```
graph TD; A[Riemannian manifold] --> B[geodesics]; A --> C[tangent spaces]; B --> D[midpoints]; C --> E[Riemannian metric];
```

geodesics

midpoints

tangent spaces

Riemannian metric

# motivation

we can study how measures evolve in  $(\mathcal{P}_2, \mathcal{W}_2)$ .

# motivation


we can study how measures evolve in  $(\mathcal{P}_2, \mathcal{W}_2)$ .



gradient flows in  $(\mathcal{P}_2, \mathcal{W}_2)$

# motivation

we can study how measures evolve in  $(\mathcal{P}_2, \mathcal{W}_2)$ .



# motivation

we can study how measures evolve in  $(\mathcal{P}_2, \mathcal{W}_2)$ .



gradient flows in  $(\mathcal{P}_2, \mathcal{W}_2)$



variational inference



particles variational inference

# motivation

we can give a geometrical interpretation to some SDEs.

# motivation

we can give a geometrical interpretation to some SDEs.



Langevin diffusion

# motivation

we can give a geometrical interpretation to some SDEs.



Langevin diffusion

sampling



# plan

- 1.** preliminaries. metric geometry, Monge and Kantorovich problems.

# plan

- 1. preliminaries.** metric geometry, Monge and Kantorovich problems.
- 2. Wasserstein spaces.** pseudo-Riemannian geometry, evolution of measures, first variations, Wasserstein gradient flows.

# plan

- 1. preliminaries.** metric geometry, Monge and Kantorovich problems.
- 2. Wasserstein spaces.** pseudo-Riemannian geometry, evolution of measures, first variations, Wasserstein gradient flows.
- 3. variational inference.** KL divergence, geodesic convexity, hints on the JKO scheme.

# plan

- 1. preliminaries.** metric geometry, Monge and Kantorovich problems.
- 2. Wasserstein spaces.** pseudo-Riemannian geometry, evolution of measures, first variations, Wasserstein gradient flows.
- 3. variational inference.** KL divergence, geodesic convexity, hints on the JKO scheme.
- 4. particles variational inference.** (RKHS), many particles systems, SVGD.

# plan

- 1. preliminaries.** metric geometry, Monge and Kantorovich problems.
- 2. Wasserstein spaces.** pseudo-Riemannian geometry, evolution of measures, first variations, Wasserstein gradient flows.
- 3. variational inference.** KL divergence, geodesic convexity, hints on the JKO scheme.
- 4. particles variational inference.** (RKHS), many particles systems, SVGD.
- 5. sampling.** Langevin diffusion as a gradient flow.

# **preliminaries.** metric geometry.

in short: abstraction of key ideas from differential geometry.

# **preliminaries.** metric geometry.


in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .

# preliminaries. metric geometry.

in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .



positive definiteness  
symmetry  
triangle inequality




# preliminaries. metric geometry.

in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .

we can already define *paths*:  $\omega : I \rightarrow \mathcal{S}$ ,  $\omega$  continuous.



positive definiteness  
symmetry  
triangle inequality

# preliminaries. metric geometry.

in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .

we can already define *paths*:  $\omega : I \rightarrow \mathcal{S}$ ,  $\omega$  continuous.

$I \subseteq \mathbb{R}$

positive definiteness  
symmetry  
triangle inequality

# preliminaries. metric geometry.

in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .

positive definiteness  
symmetry  
triangle inequality

we can already define *paths*:  $\omega : I \rightarrow \mathcal{S}$ ,  $\omega$  continuous.

$I \subseteq \mathbb{R}$

and lengths:  $L(\omega) := \sup \sum_{i=1}^{n-1} d(\omega_{t_i}, \omega_{t_{i+1}}) \approx \int_I \left\| \frac{d\omega}{dt} \right\|_d dt$ .

# preliminaries. metric geometry.

in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .

positive definiteness  
symmetry  
triangle inequality

we can already define *paths*:  $\omega : I \rightarrow \mathcal{S}$ ,  $\omega$  continuous.

$I \subseteq \mathbb{R}$

and lengths:  $L(\omega) := \sup \sum_{i=1}^{n-1} d(\omega_{t_i}, \omega_{t_{i+1}}) \approx \int_I \left\| \frac{d\omega}{dt} \right\|_d dt$ .

we say that  $\omega : [a, b] \rightarrow \mathcal{S}$  has constant speed if  $L(\omega_{[s,t]}) = \frac{t-s}{b-a} L(\omega)$ , for any  $a \leq s \leq t \leq b$ .

# preliminaries. metric geometry.

in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .

positive definiteness  
symmetry  
triangle inequality

we can already define *paths*:  $\omega : I \rightarrow \mathcal{S}$ ,  $\omega$  continuous.

$I \subseteq \mathbb{R}$

and lengths:  $L(\omega) := \sup \sum_{i=1}^{n-1} d(\omega_{t_i}, \omega_{t_{i+1}}) \approx \int_I \left\| \frac{d\omega}{dt} \right\|_d dt$ .

we say that  $\omega : [a, b] \rightarrow \mathcal{S}$  has constant speed if  $L(\omega_{[s, t]}) = \frac{t-s}{b-a} L(\omega)$ , for any  $a \leq s \leq t \leq b$ .

$\omega_2 : I_2 \rightarrow \mathcal{S}$  is a *reparametrixation* of  $\omega_1 : I_1 \rightarrow \mathcal{S}$  if  $\exists \phi : I_1 \rightarrow I_2$  such that  $\omega_1 = \omega_2 \circ \phi$ .

# preliminaries. metric geometry.

in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .

positive definiteness  
symmetry  
triangle inequality

we can already define *paths*:  $\omega : I \rightarrow \mathcal{S}$ ,  $\omega$  continuous.

$I \subseteq \mathbb{R}$

and lengths:  $L(\omega) := \sup \sum_{i=1}^{n-1} d(\omega_{t_i}, \omega_{t_{i+1}}) \approx \int_I \left\| \frac{d\omega}{dt} \right\|_d dt$ .

we say that  $\omega : [a, b] \rightarrow \mathcal{S}$  has constant speed if  $L(\omega_{[s, t]}) = \frac{t-s}{b-a} L(\omega)$ , for any  $a \leq s \leq t \leq b$ .

$\omega_2 : I_2 \rightarrow \mathcal{S}$  is a *reparametrixation* of  $\omega_1 : I_1 \rightarrow \mathcal{S}$  if  $\exists \phi : I_1 \rightarrow I_2$  such that  $\omega_1 = \omega_2 \circ \phi$ .

continuous  
non decreasing  
bijective

# preliminaries. metric geometry.

in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .

positive definiteness  
symmetry  
triangle inequality

we can already define *paths*:  $\omega : I \rightarrow \mathcal{S}$ ,  $\omega$  continuous.

$I \subseteq \mathbb{R}$

and lengths:  $L(\omega) := \sup \sum_{i=1}^{n-1} d(\omega_{t_i}, \omega_{t_{i+1}}) \approx \int_I \left\| \frac{d\omega}{dt} \right\|_d dt$ .

we say that  $\omega : [a, b] \rightarrow \mathcal{S}$  has constant speed if  $L(\omega_{[s, t]}) = \frac{t-s}{b-a} L(\omega)$ , for any  $a \leq s \leq t \leq b$ .

$\omega_2 : I_2 \rightarrow \mathcal{S}$  is a *reparametrixation* of  $\omega_1 : I_1 \rightarrow \mathcal{S}$  if  $\exists \phi : I_1 \rightarrow I_2$  such that  $\omega_1 = \omega_2 \circ \phi$ .

**fact:** any  $\omega_1$  can be reparametrized to have constant speed.

continuous  
non decreasing  
bijective

# preliminaries. metric geometry.

in short: abstraction of key ideas from differential geometry.

fix any metric space  $(\mathcal{S}, d)$ .

positive definiteness  
symmetry  
triangle inequality

we can already define *paths*:  $\omega : I \rightarrow \mathcal{S}$ ,  $\omega$  continuous.

$I \subseteq \mathbb{R}$

and lengths:  $L(\omega) := \sup \sum_{i=1}^{n-1} d(\omega_{t_i}, \omega_{t_{i+1}}) \approx \int_I \left\| \frac{d\omega}{dt} \right\|_d dt$ .

we say that  $\omega : [a, b] \rightarrow \mathcal{S}$  has constant speed if  $L(\omega_{[s, t]}) = \frac{t-s}{b-a} L(\omega)$ , for any  $a \leq s \leq t \leq b$ .

$\omega_2 : I_2 \rightarrow \mathcal{S}$  is a *reparametrixation* of  $\omega_1 : I_1 \rightarrow \mathcal{S}$  if  $\exists \phi : I_1 \rightarrow I_2$  such that  $\omega_1 = \omega_2 \circ \phi$ .

**fact:** any  $\omega_1$  can be reparametrized to have constant speed.

moreover, we can choose  $I_2 = [0, 1]$ .

continuous  
non decreasing  
bijective



# **preliminaries.** metric geometry.

fix  $x_0, x_1 \in S$ .

# preliminaries. metric geometry.

fix  $x_0, x_1 \in S$ .

a path  $\omega : [0, 1] \rightarrow S$ , with  $\omega(0) = x_0$  and  $\omega(1) = x_1$  is a *geodesic* if  $d(x_0, x_1) = L(\omega)$ .

# preliminaries. metric geometry.

fix  $x_0, x_1 \in S$ .

a path  $\omega : [0, 1] \rightarrow S$ , with  $\omega(0) = x_0$  and  $\omega(1) = x_1$  is a *geodesic* if  $d(x_0, x_1) = L(\omega)$ .

$\omega$  can be reparametrized to be a *constant speed geodesic*.

# preliminaries. metric geometry.

fix  $x_0, x_1 \in S$ .

a path  $\omega : [0, 1] \rightarrow S$ , with  $\omega(0) = x_0$  and  $\omega(1) = x_1$  is a *geodesic* if  $d(x_0, x_1) = L(\omega)$ .

$\omega$  can be reparametrized to be a *constant speed geodesic*.

$(S, d)$  is said to be a *geodesic space* if for any given  $x_0, x_1$  we can exhibit a geodesic.

# preliminaries. metric geometry.

fix  $x_0, x_1 \in S$ .

a path  $\omega : [0, 1] \rightarrow S$ , with  $\omega(0) = x_0$  and  $\omega(1) = x_1$  is a *geodesic* if  $d(x_0, x_1) = L(\omega)$ .

$\omega$  can be reparametrized to be a *constant speed geodesic*.

$(S, d)$  is said to be a *geodesic space* if for any given  $x_0, x_1$  we can exhibit a geodesic.

↓  
for us, constant speed geodesic.

# preliminaries. metric geometry.

fix  $x_0, x_1 \in S$ .

a path  $\omega : [0, 1] \rightarrow S$ , with  $\omega(0) = x_0$  and  $\omega(1) = x_1$  is a *geodesic* if  $d(x_0, x_1) = L(\omega)$ .

$\omega$  can be reparametrized to be a *constant speed geodesic*.


$(S, d)$  is said to be a *geodesic space* if for any given  $x_0, x_1$  we can exhibit a geodesic.

↓  
for us, constant speed geodesic.

if  $(S, d)$  is a geodesic space, for any  $x_0, x_1 \in S$  we can define the *midpoint* as  $\omega_{\frac{1}{2}}$ , where  $\omega_{\frac{1}{2}} = \omega(\frac{1}{2})$  and  $\omega$  is a constant speed geodesic between  $x_0$  and  $x_1$ .


# preliminaries. metric geometry.

$(\mathcal{S}, d) = (\mathbb{R}^2, \|\cdot\|_2)$  is a geodesic space.




# preliminaries. metric geometry.

$(\mathcal{S}, d) = (\mathbb{R}^2, \|\cdot\|_2)$  is a geodesic space.



$(\mathcal{S}, d) = (\mathbb{S}^2, \|\cdot\|_2)$  is a geodesic space.



$\|\cdot\|_2$  is induced by the inner product on tangent spaces of  $\mathbb{S}^2$  induced by the Euclidean one.