

Monte Carlo Methods **ETH**



**Nicholas Constantine
Metropolis**



Stanislaw Ulam

1

Books about Monte Carlo **ETH**

- **M.H. Kalos and P.A. Whitlock: „Monte Carlo Methods“ (Wiley-VCH, Berlin, 2008)**
- **J.M. Hamersley and D.C. Handscomb: „Monte Carlo Methods“ (Wiley & Sons, N.Y., 1964)**
- **K. Binder and D. Heermann: „Monte Carlo Simulations in Statistical Physics“ (Springer, Berlin, 1992)**
- **R.Y. Rubinstein: „Simulation and the Monte Carlo Method“ (Wiley & Sons, N.Y., 1981)**

2

**Simulates an experimental measuring process
with sampling and averaging.**

Big advantage:

**Systematic improvement by increasing
the number of samples N .**


Error goes like:

$$\Delta \propto \frac{1}{\sqrt{N}}$$

3

- **Statistical partition functions,
i.e. averages at constant
temperature**
- **High dimensional integrals**

4



$$\pi = 4 \int_0^1 \sqrt{1-x^2} dx$$

Choose N pairs
 (x_i, y_i) , $i = 1, \dots, N$
of random numbers
 $x_i, y_i \in \{0, 1\}$

5

$$c = 0$$

$$\text{if } y_i^2 < 1 - x_i^2 \Rightarrow c = c + 1$$

c is number of points that fall inside circle.


$$\text{area} \propto c / N$$

$$\pi(N) = 4 \frac{c}{N}$$

[applet](#)

$$\Delta = \pi - \pi(N) \propto \frac{1}{\sqrt{N}}$$

6



$$\int_a^b g(x) dx \approx (b-a) \left[\frac{1}{N} \sum_{i=1}^N g(x_i) \right]$$

7

random numbers $x_i \in [a, b]$
homogeneously distributed:

„simple sampling“

Good if $g(x)$ smooth.


8

conventional method:

choose N equidistant points \Rightarrow distance

$$h = \frac{b-a}{N}$$

one dimension



$$\text{area} \propto h^2 \propto \frac{1}{N^2} \propto \frac{1}{T^2}$$

**where T is the
computer time.**

error:

$$\Delta \propto (N \text{ area})^2 \propto T^{-2}$$

9

Error in d dimensions

in d dimensions:

$$h \propto \frac{1}{L}, \quad T \propto N = L^d \quad \Rightarrow \quad h \propto T^{-\frac{1}{d}}$$

error with conventional method:

$$\Delta \propto (N h h^d)^2 \propto T^2 h^{2(d+1)} \propto T^{-\frac{2}{d}}$$

error with Monte Carlo:

$$\Delta \propto \frac{1}{\sqrt{N}} \propto T^{-\frac{1}{2}}$$


\Rightarrow Monte Carlo better for $d > 4$

Consider n hard spheres of radius R in a $3d$ box.
Phase space is $3n$ dimensional: $(x_i, y_i, z_i), i = 1, \dots, n$.

$$r_{ij} = \sqrt{(x_i - x_j)^2 + (y_i - y_j)^2 + (z_i - z_j)^2}$$

with hard sphere constraint

$$\sqrt{(x_i - x_j)^2 + (y_i - y_j)^2 + (z_i - z_j)^2} > 2R$$



Calculate average distance:


$$\langle r_{ij} \rangle = \frac{1}{Z} \int \frac{2}{n(n-1)} \sum_{i < j} r_{ij} dx_1, \dots, dx_n dy_1, \dots, dy_n dz_1, \dots, dz_n$$

11

MC strategy

- Choose particle position.
- If the excluded volume condition is not fulfilled then reject.
- Once the n particles are placed, calculate the average of r_{ij} over all pairs (i, j) .

12



13

Calculation of integrals

$$\int_a^b g(x) dx = \int_a^b \frac{g(x)}{p(x)} p(x) dx \approx (b-a) \frac{1}{N} \sum_{i=1}^N \frac{g(x_i)}{p(x_i)}$$

if x_i randomly distributed according to $p(x)$.

Good convergence when function $\frac{g(x)}{p(x)}$ smooth.

„importance sampling“

14

$E(X)$ = energy of configuration X

Probability for system to be in X is given by

Boltzmann factor:

$$p_{eq}(X) = \frac{1}{Z_T} e^{-\frac{E(X)}{kT}}$$

Z_T is the partition function:

$$Z_T = \sum_X e^{-\frac{E(X)}{kT}}$$


$$\sum_X p_{eq}(X) = 1$$

15

Problem of sampling

$$\langle Q(T) \rangle = \sum_X Q(X) p_{eq}(X)$$

The distribution of energy E around the average $\langle E \rangle_T$ gets sharper with increasing size.



Choosing configurations equally distributed over energy would be very ineffective.

16

N.C. Metropolis, A.W. Rosenbluth,
M.N. Rosenbluth, A.H. Teller and E. Teller (1953)

importance sampling through a Markov chain:

$$X_1 \rightarrow X_2 \rightarrow \dots$$

where the probability for a configuration is $p_{eq}(X)$

Markov chain: X_t only depends on X_{t-1}

17

Properties of Markov chain

Start in configuration X and **propose** a new configuration Y with probability $T(X \rightarrow Y)$.

1. **Ergodicity**: One must be able to reach any configuration Y after a finite number of steps.

2. **Normalization**:

$$\sum_Y T(X \rightarrow Y) = 1$$

3. **Reversibility**:

$$T(X \rightarrow Y) = T(Y \rightarrow X)$$

18

The proposed configuration Y will be
accepted with probability $A(X \rightarrow Y)$.

Total probability of Markov chain is:

$$W(X \rightarrow Y) = T(X \rightarrow Y) \cdot A(X \rightarrow Y)$$

Master equation:

$$\frac{dp(X, t)}{dt} = \sum_Y p(Y) W(Y \rightarrow X) - \sum_Y p(X) W(X \rightarrow Y)$$

19

Properties of $W(X \rightarrow Y)$

• **Ergodicity:**

$$\forall X, Y : W(X \rightarrow Y) > 0$$

• **Normality:**

$$\sum_Y W(X \rightarrow Y) = 1$$

• **Homogeneity:**

$$\sum_Y p_{st}(Y) W(Y \rightarrow X) = p_{st}(X)$$

20

$$\frac{dp(X,t)}{dt} = \sum_Y p(Y) W(Y \rightarrow X) - \sum_Y p(X) W(X \rightarrow Y)$$

In stationary state one should have
equilibrium distribution (Boltzmann)

$$\frac{dp_{st}(X,t)}{dt} = 0 \quad , \quad p_{st}(X) = p_{eq}(X)$$

$$\Rightarrow \sum_Y p_{eq}(Y) W(Y \rightarrow X) = \sum_Y p_{eq}(X) W(X \rightarrow Y)$$

sufficient condition is **detailed balance**:

$$p_{eq}(Y) W(Y \rightarrow X) = p_{eq}(X) W(X \rightarrow Y)$$

21

Metropolis (M(RT)²)

$$A(X \rightarrow Y) = \min \left(1, \frac{p_{eq}(Y)}{p_{eq}(X)} \right)$$

Boltzmann:

$$p_{eq}(X) = \frac{1}{Z_T} e^{-\frac{E(X)}{kT}}$$

$$A(X \rightarrow Y) = \min \left(1, e^{-\frac{E(Y)-E(X)}{kT}} \right) = \min \left(1, e^{-\frac{\Delta E}{kT}} \right)$$

If energy decreases always accept
increases accept with probability $e^{-\frac{\Delta E}{kT}}$.

22

Roy C. Glauber (1963)
(Nobel prize 2005)




$$A(X \rightarrow Y) = \frac{e^{-\frac{\Delta E}{kT}}}{1 + e^{-\frac{\Delta E}{kT}}}$$


**also
fulfills
detailed
balance**

23

The Ising Model



**Spins
on a
lattice**



**Ernst Ising
(1900-1998)**

- **Magnetic Systems**
- **Opinion models**
- **Binary mixtures**

24

Binary variables:

$$\sigma_i = \pm 1, \quad i = 1, \dots, N$$

on a graph of N sites

interacting via the Hamiltonian:


$$\mathcal{H} = E = -J \sum_{i,j:nn}^N \sigma_i \sigma_j - H \sum_{i=1}^N \sigma_i$$

25

Order parameter

spontaneous magnetization:

$$M(T) = \lim_{H \rightarrow 0} \left\langle \frac{1}{N} \sum_{i=1}^N \sigma_i \right\rangle$$



$$M \propto (T - T_c)^\beta$$

$$\beta = 1/8 \quad (2d)$$


$$\beta \approx 0.326 \quad (3d)$$

26

$$\chi(T) \propto |T - T_c|^{-\gamma}$$

$$\gamma = 7/4 \text{ (2d)}$$

$$\gamma \approx 1.24 \text{ (3d)}$$



numerical data from a finite system

27

MC of the Ising Model

Single flip Metropolis:

- Choose one site i (having spin σ_i).
- Calculate $\Delta E = E(Y) - E(X) = 2J\sigma_i h_i$.
- If $\Delta E < 0$ then flip spin: $\sigma_i \rightarrow -\sigma_i$.
- If $\Delta E > 0$ flip with probability $\exp(-\Delta E/kT)$.

where h_i is the local field at site i

$$h_i = \sum_{nn \text{ of } i} \sigma_j$$

[applet](#)

28

Consider two species A and B distributed with given concentrations on the sites of a lattice.

E_{AA} is energy of A-A bond.

E_{BB} is energy of B-B bond.

E_{AB} is energy of A-B bond.


Set $E_{AA} = E_{BB} = 0$ and $E_{AB} = 1$.

Number of each species is **constant**.

29

Kawasaki dynamics

- Choose any A-B bond.
- Calculate ΔE for A-B \rightarrow B-A.
- **Metropolis:** If $\Delta E \leq 0$ flip, else flip with $p = \exp(-\beta\Delta E)$.
- **Glauber:** Flip with probability $p = \exp(-\beta\Delta E)/(1 + \exp(-\beta\Delta E))$.



Kyozi Kawasaki

$$\beta = 1/kT$$


30

Adsorption
an Einzelplätzen

Mono-
schicht



Mehrfach-
schichten

Porenkondensation



31

Interfaces




surface tension

$$\gamma = f_{A+B} - f_A$$


32

$$\mathcal{H} = E = -J \sum_{i,j:nn}^N \sigma_i \sigma_j$$

Fixed boundary conditions:
upper half „+“, lower half „-“
Simulate with Kawasaki
dynamics at temperature T .




33



interface width W :

$$W = \sqrt{\frac{1}{N} \sum_i (h_i - \bar{h})^2}$$



roughening transition

34




Figure 1. Interface width W versus time t for the RSOS model (Ref. [11]) in 1 + 1 dimensions, in two different lattice lengths L .

Family-Vicsek scaling (1985):

$$W(L, t) = L^{\xi} f(t / L^z)$$

ξ is the roughening exponent, z the dynamic exponent.

35

Self-affine scaling

$$W(L, t) = L^{\xi} f(t / L^z)$$

$$u = t / L^z$$

$$t \rightarrow \infty : W \propto L^{\xi} \Rightarrow f(u \rightarrow \infty) = \text{const}$$

$$L \rightarrow \infty : W \propto t^{\beta} \Rightarrow f(u \rightarrow 0) \propto u^{\beta}$$

$$W \propto L^{\xi} u^{\beta} = L^{\xi} (t / L^z)^{\beta} = L^{\xi - \beta z} t^{\beta}$$

\Rightarrow

$$\beta = \frac{\xi}{z}$$

β is the growth exponent.


Numerically these laws are verified by data collapse.

36

Add next-nearest neighbor interaction:

$$\mathcal{H} = E = -J \sum_{i,j:nn} \sigma_i \sigma_j - K \sum_{i,j:nnn} \sigma_i \sigma_j$$

punishes curvature \leftrightarrow introduces stiffness



37

Shape of drop

- Start with a block of +1 sites attached to wall of an $L \times L$ system filled with -1.
- Hamiltonian including gravity g :


$$\mathcal{H} = E = -J \sum_{i,j:nn} \sigma_i \sigma_j - K \sum_{i,j:nnn} \sigma_i \sigma_j - \sum_j h_j \sum_{\text{line } j} \sigma_i$$

with $h_j = h_1 + \frac{(j-1)(h_L - h_1)}{L-1}$ and $g = \frac{h_L - h_1}{L}$


- Use Kawasaki dynamics and further do not allow for disconnected clusters of +1.

38

$L = 257, V = 6613, g = 0.001$
after 5×10^7 MC updates
averaged over 20 samples.




Contact angle θ is function
of temperature and vanishes
when approaching T_c .



39

Solid on solid model (SOS) ETH

adatoms and
surface growth



interface without islands and **without overhangs**

$$\mathcal{H} = E = -\varepsilon \sum_{i,j:nn}^N |h_i - h_j|$$

40

- T. Vicsek, „Fractal Growth Phenomena“ (World Scientific, Singapore, 1989)
- A.-L. Barabasi and H.E. Stanley, „Fractal Concepts in Surface Growth“ (Cambridge Univ. Press, 1995)
- H.J. Herrmann, „Geometric Cluster Growth Models and Kinetic Gelation“, Phys. Rep. 136, 153 (1986)

41

Irreversible growth



No thermal equilibrium

- Deposition and aggregation patterns
- Fluid instabilities
- Electric breakdown
- Biological morphogenesis
- Fracture and Fragmentation



42



43


Random deposition

simplest possible growth model

Pick a random column.
Add a particle on top of
that column


$$\langle h \rangle \sim t$$

$$W \sim t^{1/2}$$



$$\beta = 1/2$$

$$\xi = 1/2$$



44


Random deposition with surface diffusion **ETH**

Particle can move a short distance to find a more stable configuration.


Pick a random column i .
Compare $h(i)$ and $h(i+1)$.
Particle is added onto whichever is lower. If they are equal, add to column i or $i+1$ with equal probability.

$$\langle h \rangle \sim t$$

$$W \sim t^{1/4}$$



$$\beta = 1/4$$
$$\xi = 1/2$$




45


Restricted Solid On Solid Model (RSOS) **ETH**

Neighbouring sites may not have a height difference greater than 1.

Pick a random column i .
Add a particle only if $h(i) \leq h(i-1)$ and $h(i) \leq h(i+1)$.
Otherwise pick a new column.



$$\beta = 1/3$$
$$\xi = 1/2$$





46

simple model for tumor growth or epidemic spread

Each site that is a neighbour of an occupied site has an equal probability of being filled.


Make a list of neighbour sites (red).
Pick one at random. Add it to cluster (green).
Neighbours of this site then become red....

There can be more than one growth site in a column. Can get overhangs.



47

Eden cluster




$$\beta = 1/3$$
$$\xi = 1/2$$

48


Particles fall vertically from above and stick when they touch a particle below or a neighbour on one side.

Pick a column i . Let particle fall till it touches a neighbour. Possible growth sites are indicated in red. Only one possible site per column.



$$\beta = 1/3$$

$$\xi = 1/2$$



49

Growth equations

Edwards Wilkinson equation:

$$\frac{\partial h(x,t)}{\partial t} = \nu \Delta h + \eta(x,t)$$

S.F. Edwards and D.R. Wilkinson (1982)

$$\beta = 1/4 \text{ and } \xi = 1/2$$

noise

KPZ equation:

$$\frac{\partial h(x,t)}{\partial t} = \nu \Delta h + \frac{\lambda}{2} (\nabla h)^2 + \eta(x,t)$$

M. Kardar, G. Parisi and Y.-C. Zhang (1986)



$$\beta = 1/3 \text{ and } \xi = 1/2$$

50

Diffusion Limited Aggregation (DLA)


A particles starts a long way from surface.

It diffuses until it first touches the surface. If it moves too far away, another particle is started.




51

Electrodeposition



52

DLA clusters




fractal
dimension

$$d_f = 1.7$$

(in 2d)

53

Viscous Fingering



54


DLA -- N=2500



DLA -- N=5000



DLA -- N=10000




anisotropy on square lattice:




55

Scale Invariance



56



57

Dielectric breakdown model (DBM)

Solve Laplacian field φ :


$$\Delta\phi = 0$$

and occupy site
at boundary
with probability:



$$p \propto (\nabla\phi)^\eta$$


same as DLA

$\eta = 1$





58





same as Eden model





Fractal dimension is $d_f \approx 1.42$ in 2d, $d_f \approx 1.7$ in 3d.

dynamical scaling: $n_s = s^{-2} f(s / t^z)$

with dynamical exponent z .

63

Simulated Annealing

- P.J.M. Van Laarhoven and E.H. Aarts, „Simulated Annealing: Theory and Applications“ (Kluwer, 1987)
- A. Das and B.K. Chakrabarti (eds.), „Quantum Annealing and Related Optimization Methods“ Lecture Notes No. 679 (Springer, 1990)

64

S. Kirkpatrick, C.D. Gelatt and M.P. Vecchi, 1983

SA is a stochastic optimization technique.

**Given a finite set S of solutions
and a cost function $F: S \rightarrow \mathbb{R}$.**

Search a global minimum:

$$s^* \in S^* := \{ s \in S : F(s) \leq F(t) \quad \forall t \in S \}.$$

Difficult when S is very big (like $|S| = n!$).

65

Travelling Salesman

Given n cities σ_i and the travelling costs $c(\sigma_i, \sigma_j)$.

We search for the cheapest trip through all cities.

- $S = \{ \text{permutations of } \{1, \dots, n\} \}$
- $F(\sigma) = \sum_{i=1..n} c(\sigma_i, \sigma_j)$ für $\sigma \in S$


Finding the best trajectory is a **NP-complex problem,
i.e. time to solve grows faster than any polynomial of n .**

66

Make local changes:

⇒ Define close configuration on S

$N : S \rightarrow 2^S$ with $i \in N(j) \Leftrightarrow j \in N(i)$



Traditional optimization algorithm:

Improve systematically the costs by exploring close solutions.

If $F(\sigma') < F(\sigma)$ replace $\sigma := \sigma'$ until $F(\sigma) \leq F(t)$ for all $t \in N(\sigma)$.

Problem: One gets stuck in local minima.

67

Travelling Salesman

Simulated annealing optimization algorithm:

If $F(\sigma') < F(\sigma)$ replace $\sigma := \sigma'$

If $F(\sigma') > F(\sigma)$ replace $\sigma := \sigma'$

with probability $\exp(-\Delta F / T)$


with $\Delta F = F(\sigma') - F(\sigma) > 0$

T is a constant (like a temperature)

Go slowly to $T \rightarrow 0$ in order to find global minimum.

Applet

68



Different cooling protocols are possible.


Asymptotic convergence is guarantied and leads to an exponential convergence time.

69

- D.H. Rothman and S. Zaleski, „Lattice-Gas Cellular Automata“ (Cambridge Univ. Press, 1997)
- J.-P. Rivet and J.P. Boon, „Lattice Gas Hydrodynamics“ (Cambridge Univ. Press, 2001)
- D.A. Wolf-Gladrow, „Lattice-Gas Cellular Automata and Lattice Boltzmann Models“ (Lecture Notes, Springer, 2000)

70

Particles move on a triangular lattice and follow the following collision rules:




Momentum is conserved at each collision.
It can be proven (Chapman-Enskog) that its continuum limit is the Navier Stokes eq.

71

von Karman street

velocity field of a fluid behind an obstacle



Each vector is an average over time and space.

72

Finding the solution (root) of an equation:

$$f(x) = 0$$

is equivalent to the optimization problem of finding the minimum (or maximum) of $F(x)$ given by:

$$\frac{d}{dx} F(x) = 0$$


73

Newton method

Be x_0 a first guess, then linearize around x_0 :

$$f(x_1) \approx f(x_0) + (x_1 - x_0)f'(x_0) = 0$$

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$




74

If derivative of f is not known analytically:

$$f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}}$$

$$x_{n+1} = x_n - (x_n - x_{n-1}) \frac{f(x_n)}{f(x_n) - f(x_{n-1})}$$

75




76

Take two starting values x_1 and x_2
with $f(x_1) < 0$ and $f(x_2) > 0$.

Define mid-point x_m as $x_m = (x_1 + x_2) / 2$.

If $\text{sign}(f(x_m)) = \text{sign}(f(x_1))$
then replace x_1 by x_m
otherwise replace x_2 by x_m .

77



78


Take two starting values x_1 and x_2
with $f(x_1) < 0$ and $f(x_2) > 0$.

Approximate f by a straight line between
 $f(x_1)$ and $f(x_2)$ and calculate its root as:

$$x_m = (f(x_1) x_2 - f(x_2) x_1) / (f(x_1) - f(x_2)).$$

If $\text{sign}(f(x_m)) = \text{sign}(f(x_1))$, then
replace x_1 by x_m otherwise replace x_2 by x_m .

79



80

Be \vec{x} a *N*-dimensional vector.

System of *N* coupled equations:

$$\vec{f}(\vec{x}) = 0$$

corresponding to the *N*-dimensional optimization problem:

$$\vec{\nabla} F(\vec{x}) = 0$$

81

N-dimensional Newton method

Define the Jacobi matrix:

$$J_{i,j}(\vec{x}) = \frac{\partial f_i(\vec{x})}{\partial x_j}$$

Must be non-singular and
also well-conditioned
for numerical inversion.

$$\vec{x}_{n+1} = \vec{x}_n - \vec{J}^{-1} \vec{f}(\vec{x}_n)$$

82

$$b_{11}x_1 + \dots + b_{1N}x_N = c_1$$

$$\begin{matrix} \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{matrix} \Leftrightarrow \vec{B}\vec{x} = \vec{c}$$

$$b_{N1}x_1 + \dots + b_{NN}x_N = c_N$$

solution:

$$\vec{x} = \vec{B}^{-1}\vec{c}$$

83

$$\vec{f}(\vec{x}) = \vec{B}\vec{x} - \vec{c} = 0 \Rightarrow \vec{J} = \vec{B}$$

apply Newton method: $\vec{x}_{n+1} = \vec{x}_n - \vec{J}^{-1}\vec{f}(\vec{x}_n)$

$$\vec{x}_{n+1} = \vec{x}_n - \vec{B}^{-1}(\vec{B}\vec{x}_n - \vec{c}) = \vec{B}^{-1}\vec{c}$$

\Rightarrow exact solution in one step

84

If the derivatives are not known analytically:

$$J_{i,j}(\vec{x}) = \frac{f_i(\vec{x} + h_j \vec{e}_j) - f_i(\vec{x})}{h_j}$$

where h_j should be chosen as:
being ε the machine precision,
i.e. $\approx 10^{-16}$ for a 64 bit computer.

$$h_j \approx x_j \sqrt{\varepsilon}$$

85

Other techniques

Relaxation method:

$$\vec{f}(\vec{x}) = 0 \quad \rightarrow \quad x_i = g_i(x_j, j \neq i), \quad i = 1, \dots, N$$

Start with $x_i(0)$ and iterate: $x_i(t+1) = g_i(x_j(t))$.

Gradient methods:

1. Steepest descent
2. Conjugate gradient

86

First order ODE, initial value problem:

$$\boxed{\frac{dy}{dt} = f(y, t)} \quad \text{with } y(t_0) = y_0$$

examples:

radioactive decay

$$\boxed{\frac{dN}{dt} = -\lambda N}$$

coffee cooling

$$\boxed{\frac{dT}{dt} = -\gamma(T - T_{room})}$$

87

Euler method

explicit forward integration algorithm

choose small Δt , Taylor expansion:

$$\begin{aligned} y(t_0 + \Delta t) &= y(t_0) + \Delta t \frac{dy}{dt}(t_0) + O(\Delta^2 t) \\ &= y_0 + \Delta t \cdot f(y_0, t_0) + O(\Delta^2 t) \equiv y(t_1) \equiv y_1 \end{aligned}$$

convention: $t_n = t_0 + n\Delta t$, $y_n = y(t_n)$

88


Start with y_0 and iterate:

$$y_{n+1} = y_n + \Delta t \cdot f(y_n, t_n) + O(\Delta^2 t)$$

This is the simplest **finite difference** method.

Since the error goes with $\Delta^2 t$ one needs a very small Δt and that is numerically very expensive.

89



90

$$m \frac{d^2 x}{dt^2} = F(x)$$

2nd order ODE

Transform in a system of two coupled ODEs of first order.

$$\begin{aligned} \frac{dx}{dt} &= v \\ \frac{dv}{dt} &= \frac{F(x)}{m} \end{aligned}$$

91

Euler method for N coupled ODEs

N coupled ODEs of first order:

$$\frac{dy_i}{dt} = f_i(y_1, \dots, y_N, t), \quad i = 1, \dots, N$$

iterate with a small Δt :

$$y_i(t_{n+1}) = y_i(t_n) + \Delta t \cdot f_i(y_1(t_n), \dots, y_N(t_n), t_n) + O(\Delta^2 t)$$

with

$$t_n = t_0 + n \cdot \Delta t$$

92

If the error at one time-step is $O(\Delta^n t)$ the method is „**locally of order n** “. To consider a fixed time interval T one needs $T / \Delta t$ time-steps so that the total error is:

$$\frac{T}{\Delta t} O(\Delta^n t) = O(\Delta^{n-1} t)$$

and therefore the method is „**globally of order $n-1$** “.

The Euler method is globally of first order.

93

Runge - Kutta method



Carl David Tolmé Runge
(1856-1927)



Martin Wilhelm Kutta
(1867-1944)

94

- G.E. Forsythe, M.A. Malcolm and C.B. Moler, „Computer Methods for Mathematical Computations“ (Prentice Hall, Englewood Cliffs, NJ, 1977), Chapter 6
- E. Hairer, S.P. Nørsett and G. Wanner, “Solving Ordinary Differential Equations I” (Springer, Berlin, 1993)
- W.H. Press, B.P. Flannery, S.A. Teukolsky and W.T. Vetterling, “Numerical Recipes” (Cambridge University Press, Cambridge, 1988), Sect. 16.1 and 16.2
- J.C. Butcher, „The Numerical Analysis of Ordinary Differential Equations“ (Wiley, New York, 1987)
- J.D. Lambert, „Numerical Methods for Ordinary Differential Equations“ (John Wiley & Sons, New York, 1991)
- L.F. Shampine, „Numerical Solution of Ordinary Differential Equations“ (Chapman and Hall, London, 1994)

2nd order Runge - Kutta method

Extrapolate using Euler method

to the value halfway, i.e. at $t + \Delta t / 2$:

$$y_i(t + \frac{1}{2} \Delta t) = y_i(t) + \frac{1}{2} \Delta t \cdot f(y_i(t), t)$$

Evaluate derivative in Euler method at this value:

$$y_i(t + \Delta t) = y_i(t) + \Delta t \cdot f(y_i(t + \frac{1}{2} \Delta t), t + \frac{1}{2} \Delta t) + O(\Delta^3 t)$$

$$k_1 = \Delta t \cdot f(y_n, t_n)$$

$$k_2 = \Delta t \cdot f(y_n + k_1 / 2, t_n + \Delta t / 2)$$

$$k_3 = \Delta t \cdot f(y_n + k_2 / 2, t_n + \Delta t / 2)$$

$$k_4 = \Delta t \cdot f(y_n + k_3, t_n + \Delta t)$$

RK4

$$y_{n+1} = y_n + \frac{k_1}{6} + \frac{k_2}{3} + \frac{k_3}{3} + \frac{k_4}{6} + O(\Delta^5 t)$$

97

RK4

- k_1 is the slope at the beginning of the interval.
- k_2 is the slope at the midpoint of the interval, using slope k_1 to determine the value of y at the point $t_n + \Delta t / 2$ using Euler method.
- k_3 is again the slope at the midpoint, but now using the slope k_2 to determine the y -value.
- k_4 is the slope at the end of the interval, with its y -value determined using slope k_3 .

Then use Euler method with

$$\text{slope} = \frac{k_1 + 2k_2 + 2k_3 + k_4}{6}$$

98

iteration:

$$y_{n+1} = y_n + \Delta t \cdot \sum_{i=1}^q \omega_i k_i$$

with

$$k_i = f(y_n + \Delta t \sum_{j=1}^{i-1} \beta_{ij} k_j, t_n + \Delta t \alpha_i)$$

and $\alpha_1 = 0$

explicit

implicit:

$$k_i = f(y_n + \Delta t \sum_{j=1}^q \beta_{ij} k_j, t_n + \Delta t \alpha_i)$$

99

$q = 1$ is the Euler method (unique)

Butcher array or Runge-Kutta tableau:

α_2	β_{21}
α_3	$\beta_{31} \quad \beta_{32}$
\vdots	\vdots
α_q	$\beta_{q1} \quad \beta_{q2} \quad \dots \quad \beta_{q,q-1}$
	$\omega_1 \quad \omega_2 \quad \dots \quad \omega_{q-1} \quad \omega_q$

**resumes all
parameters
of the general
Runge-Kutta**

Butcher array:

its stage is: $q = 4$

0				
$\frac{1}{2}$	$\frac{1}{2}$			
$\frac{1}{2}$	0	$\frac{1}{2}$		
1	0	0	1	
	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{6}$

RK4 is of order $p = 4$

[applet](#)

101

Order of Runge - Kutta method

The R-K method is of order p if:

$$y(t + \Delta t) - y(t) - \Delta t \cdot \sum_{i=1}^q \omega_i k_i = O(\Delta t^{p+1})$$

One must choose the α_i , β_{ij} and ω_i for $i, j \in [1, p]$ such that the left hand side = 0 in $O(\Delta t^m)$ for all $m \leq p$.

Taylor expansion:

$$y(t + \Delta t) - y(t) = \sum_{m=1}^p \frac{1}{m!} \Delta t^m \cdot \left[\frac{d^{m-1} f}{dt^{m-1}} \right]_{y(t), t} + O(\Delta t^{p+1})$$

102

$$\Rightarrow \sum_{i=1}^q \omega_i k_i = \sum_{m=1}^p \frac{1}{m!} \Delta t^{m-1} \cdot \left[\frac{d^{m-1} f}{dt^{m-1}} \right]_{y(t), t} \quad \text{up to } O(\Delta t^{p+1})$$

Example $q = p = 1$: $\omega_1 \cdot f(y_n, t_n) = f(y_n, t_n) \Rightarrow \omega_1 = 1$

\Rightarrow gives Euler method.

Example $q = p = 2$: $\omega_1 \cdot k_1 + \omega_2 \cdot k_2 = f_n + \frac{1}{2} \Delta t \left[\frac{df}{dt} \right]_n$

where index „ n “ means „at (y_n, t_n) “.

103

Example $q = p = 2$

$$\omega_1 \cdot k_1 + \omega_2 \cdot k_2 = f_n + \frac{1}{2} \Delta t \left[\frac{df}{dt} \right]_n = f_n + \frac{1}{2} \Delta t \left(\left[\frac{\partial f}{\partial t} \right]_n + \left[\frac{\partial f}{\partial y} \right]_n \cdot f_n \right)$$

insert

$$\begin{aligned} k_1 &= f(y_n, t_n) \equiv f_n \\ k_2 &= f(y_n + \Delta t \cdot \beta_{21} \cdot k_1, t_n + \Delta t \cdot \alpha_2) \\ &= f_n + \Delta t \cdot \beta_{21} \cdot \left[\frac{\partial f}{\partial y} \right]_n \cdot f_n + \Delta t \cdot \alpha_2 \cdot \left[\frac{\partial f}{\partial t} \right]_n + O(\Delta t^2) \end{aligned}$$

$$\Rightarrow \omega_1 + \omega_2 = 1, \quad \omega_2 \cdot \alpha_2 = \frac{1}{2}, \quad \omega_2 \cdot \beta_{21} = \frac{1}{2}$$

3 equations for 4 parameters \Rightarrow one-parameter family

104

$$y_{n+1} = y_n + \Delta t \cdot \left[(1 - \omega_2) \cdot k_1 + \omega_2 k_2 \right]$$

$$k_1 = f(y_n, t_n)$$

$$k_2 = f\left(y_n + \frac{\Delta t}{2\omega_2}, t_n + \frac{\Delta t}{2\omega_2}\right)$$

105

Order of Runge - Kutta method

To obtain a Runge-Kutta method of given order p one needs a minimum stage of q_{\min} .

p	1	2	3	4	5	6	7	8	9	10
q_{\min}	1	2	3	4	6	7	9	11	12 – 17	13 - 17



John Butcher

106

Example: Lorenz equation

Is a simplified system of equations describing the 2-dimensional flow of fluid of uniform depth in the presence of an imposed temperature difference taking into account gravity, buoyancy, thermal diffusivity, and kinematic viscosity (friction).

σ Prandtl number
 ρ Rayleigh number
 $\sigma = 10$, $\beta = 8/3$
 ρ is varied.
Chaos for $\rho = 28$.

$$\begin{aligned}y_1' &= \sigma(y_2 - y_1) \\y_2' &= y_1(\rho - y_3) - y_2 \\y_3' &= y_1 y_2 - \beta y_3\end{aligned}$$



Edward Norton Lorenz
(1963)

[applet](#)

C.Sparrow: „The Lorenz Equations: Bifurcations, Chaos and Strange Attractors“ (Springer Verlag, N.Y., 1982)

107

Example: Lorenz equation

Chaotic solutions of the
Lorenz equation exist and
are not numerical artefacts
(14th math problem of Smale).



Warwick Tucker
(2002)

108

Define the error $\delta_{expected}$ you want to accept.
Then measure the real error $\delta_{measured}$ and
define a new
time-step through:

$$\Delta t_{new} = \Delta t_{old} \left(\frac{\delta_{expected}}{\delta_{measured}} \right)^{\frac{1}{p+1}}$$

because $\delta \propto \Delta t^{p+1}$.

109

Error estimation

Let us consider a method of order p , (error = $\Phi \Delta t^{p+1}$).
Be y_1 the predicted value for $2\Delta t$
and y_2 the predicted value making two steps Δt .

Define error as $\delta = y_1 - y_2$

$$y(t + 2\Delta t) = \begin{cases} y_1 + (2\Delta t)^{p+1} \Phi + O(\Delta t^{p+2}) \\ y_2 + 2(\Delta t)^{p+1} \Phi + O(\Delta t^{p+2}) \end{cases}$$
$$\Rightarrow \delta = (2^{p+1} - 2) \Delta t^{p+1} \Phi + O(\Delta t^{p+2})$$

110

improve systematically:

$$y(t + \Delta t) = y_2 + \frac{2\delta}{2^{p+1} - 2} + O(\Delta t^{p+2})$$

example RK4:

$$y(t + \Delta t) = y_2 + \frac{\delta}{15} + O(\Delta t^{p+2})$$

111


General for two different time steps Δt_1 and Δt_2

$$\begin{aligned} y &= y_{\Delta t_i} + \Phi \Delta t_i^p + O(\Delta t_i^{p+1}) \quad , \quad i = 1, 2 \\ (\Delta t_2^p - \Delta t_1^p) y &= \Delta t_2^p y_{\Delta t_1^p} - \Delta t_1^p y_{\Delta t_2^p} + O(\Delta t_i^{p+1}) \\ y &= \frac{\Delta t_2^p y_{\Delta t_1^p} - \Delta t_1^p y_{\Delta t_2^p}}{\Delta t_2^p - \Delta t_1^p} + O(\Delta t_i^{p+1}) \end{aligned}$$

112

Calculate for
various Δt_i
and extrapolate

$$y = \lim_{\Delta t_i \rightarrow 0} y_{\Delta t_i}$$



113

Bulirsch - Stoer Method

Extrapolate with rational function:

$$y_{\Delta t_n} = \frac{p_0 + p_1 \Delta t_n + \dots + p_k (\Delta t_n)^k}{q_0 + q_1 \Delta t_n + \dots + q_m (\Delta t_n)^m} \xrightarrow{\Delta t_n \rightarrow 0} y$$
$$\Delta t_n = \frac{\Delta t}{n}, \quad n = 2, 4, 6, 8, 12, 16, 24, 32, 48, 64, 96, \dots$$

Choose k and m appropriately.

**R. Bulirsch and J. Stoer, „Introduction to
Numerical Analysis“ (Springer, NY, 1992)**

114

idea:
$$y(t + \Delta t) \approx y(t) + \Delta t \cdot \frac{f(y(t)) + f(y(t + \Delta t))}{2}$$

implicit equation

make prediction

using Taylor:

$$y^P(t + \Delta t) = y(t) + \Delta t \cdot \frac{dy}{dt}(t) + O(\Delta t^2)$$

correct by

inserting:
$$y^C(t + \Delta t) = y(t) + \frac{\Delta t}{2} [f(y(t)) + f(y^P(t + \Delta t))] + O(\Delta t^3)$$

Can be iterated by again inserting corrected value.

115

3th order predictor-corrector

Predict through 3rd order Taylor expansion:

$$y^P(t + \Delta t) = y(t) + \Delta t \cdot \frac{dy}{dt}(t) + \frac{\Delta t^2}{2} \cdot \frac{d^2 y}{dt^2}(t) + \frac{\Delta t^3}{6} \cdot \frac{d^3 y}{dt^3}(t) + O(\Delta t^4)$$

$$\left(\frac{dy}{dt} \right)^P(t + \Delta t) = \frac{dy}{dt}(t) + \Delta t \cdot \frac{d^2 y}{dt^2}(t) + \frac{\Delta t^2}{2} \cdot \frac{d^3 y}{dt^3}(t) + O(\Delta t^3)$$

$$\left(\frac{d^2 y}{dt^2} \right)^P(t + \Delta t) = \frac{d^2 y}{dt^2}(t) + \Delta t \cdot \frac{d^3 y}{dt^3}(t) + O(\Delta t^2)$$

$$\left(\frac{d^3 y}{dt^3} \right)^P(t + \Delta t) = \frac{d^3 y}{dt^3}(t) + O(\Delta t)$$

116

3th order predictor- corrector

use equation:

$$\left(\frac{dy}{dt}\right)^C(t + \Delta t) = f(y^P(t + \Delta t))$$

define error:

$$\delta = \left(\frac{dy}{dt}\right)^C(t + \Delta t) - \left(\frac{dy}{dt}\right)^P(t + \Delta t)$$

correct:

$$\begin{aligned} y^C(t + \Delta t) &= y^P + c_0 \cdot \delta \\ \left(\frac{d^2 y}{dt^2}\right)^C(t + \Delta t) &= \left(\frac{d^2 y}{dt^2}\right)^P + c_2 \cdot \delta \\ \left(\frac{d^3 y}{dt^3}\right)^C(t + \Delta t) &= \left(\frac{d^3 y}{dt^3}\right)^P + c_3 \cdot \delta \end{aligned}$$

Procedure
can be
repeated.

Gear coefficients:

$$c_0 = 3/8$$

$$c_2 = 3/4$$

$$c_3 = 1/6$$

117

5th order predictor- corrector

Be $\mathbf{r}_0 \equiv \mathbf{y}$. Then one can define the time derivatives:

$$\mathbf{r}_1 = \partial t(d\mathbf{r}_0/dt), \mathbf{r}_2 = 1/2\partial t^2(d^2\mathbf{r}_0/dt^2), \mathbf{r}_3 = 1/6\partial t^3(d^3\mathbf{r}_0/dt^3), \text{ etc.}$$

Predictor:

$$\begin{pmatrix} \mathbf{r}_0^p(t + \partial t) \\ \mathbf{r}_1^p(t + \partial t) \\ \mathbf{r}_2^p(t + \partial t) \\ \mathbf{r}_3^p(t + \partial t) \\ \mathbf{r}_4^p(t + \partial t) \\ \mathbf{r}_5^p(t + \partial t) \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 & 4 & 5 \\ 0 & 0 & 1 & 3 & 6 & 10 \\ 0 & 0 & 0 & 1 & 4 & 10 \\ 0 & 0 & 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \mathbf{r}_0(t) \\ \mathbf{r}_1(t) \\ \mathbf{r}_2(t) \\ \mathbf{r}_3(t) \\ \mathbf{r}_4(t) \\ \mathbf{r}_5(t) \end{pmatrix}$$

118

5th order predictor- corrector

1st order eq.:

$$\frac{dr}{dt} = f(r) \Rightarrow r_1^C = f(r_0^P) \Rightarrow \Delta r = r_1^C - r_1^P$$

2nd order eq.:

$$\frac{d^2r}{dt^2} = f(r) \Rightarrow r_2^C = 2f(r_0^P) \Rightarrow \Delta r = r_2^C - r_2^P$$

Corrector:

$$\begin{pmatrix} r_0^C(t + \partial t) \\ r_1^C(t + \partial t) \\ r_2^C(t + \partial t) \\ r_3^C(t + \partial t) \\ r_4^C(t + \partial t) \\ r_5^C(t + \partial t) \end{pmatrix} = \begin{pmatrix} r_0^P(t + \partial t) \\ r_1^P(t + \partial t) \\ r_2^P(t + \partial t) \\ r_3^P(t + \partial t) \\ r_4^P(t + \partial t) \\ r_5^P(t + \partial t) \end{pmatrix} + \begin{pmatrix} c_0 \\ c_1 \\ c_2 \\ c_3 \\ c_4 \\ c_5 \end{pmatrix} \cdot \Delta r$$

119

Gear coefficients (1971)

1st order equation:

Value	c_0	c_1	c_2	c_3	c_4	c_5
3	5/12	1	1/2			
4	3/8	1	3/4	1/6		
5	251/720	1	11/12	1/3	1/24	
6	95/288	1	25/24	35/72	5/48	1/120


2nd order equation:

Value	c_0	c_1	c_2	c_3	c_4	c_5
3	0	1	1			
4	1/6	5/6	1	1/3		
5	19/120	3/4	1	1/2	1/12	
6	3/20	251/360	1	11/18	1/6	1/60

Function	c_0	c_1	c_2	c_3	c_4	c_5
$\dot{r} = f(r)$	95/288	1	25/24	35/72	5/48	1/120
$\ddot{r} = f(r)$	3/20	251/360	1	11/18	1/6	1/60
$\ddot{r} = f(r, \dot{r})$	3/16	251/360	1	11/18	1/6	1/60

120

for fixed number of iterations n



121

Sets of coupled ODEs

Explicit Runge Kutta and Predictor Corrector methods can be straightforwardly generalized to a set of coupled 1st order ODEs:

$$\frac{dy_i}{dt} = f_i(y_1, \dots, y_N, t), \quad i = 1, \dots, N$$

by inserting simultaneously all the values of the previous iteration.

122


example:

$$y'(t) = -15y(t) \quad , \quad t \geq 0 \quad , \quad y(0) = 1$$

Euler:

$$y(t + \Delta t) = y(t) - \Delta t \, 15 \, y(t) + O(\Delta t^2)$$

Becomes unstable if Δt not small enough:



Use implicit method (Adams – Moulton):

$$y(t + \Delta t) = y(t) + \frac{1}{2} \Delta t \left(f(y(t), t) - f(y(t + \Delta t), t + \Delta t) \right) + O(\Delta t^2)$$

123

Stiff sets of equations

$$\vec{y}'(t) = \vec{K} \cdot \vec{y}(t) + \vec{f}(t)$$

This system is called „stiff“ if matrix K has at least one very large eigenvalue.

$$\vec{y}'(t) = \vec{f}(\vec{y}(t), t)$$

This system is called „stiff“ if Jacobi matrix has at least one very large eigenvalue.

example:

$$\begin{aligned} y_1' &= 998y_1 + 1998y_2 \\ y_2' &= 999y_1 + 1999y_2 \\ y_1(0) &= 1 \quad , \quad y_2(0) = 1 \end{aligned}$$

Solve with with implicit method.

124