Lensing Galaxies in the CFHT Legacy Survey

Rafael Küng¹ Prasenjit Saha¹ Elisabeth Baeten² Jonathan Coles³ Claude Cornen² Christine Macmillan² Phil Marshall⁴ Anupreeta More⁵ Surhud More⁵ Aprajita Verma⁶ Julianne K. Wilcox²

¹Physik-Institut, University of Zurich, Zurich, Switzerland ²Zooniverse, c/o Astrophysics Department, University of Oxford, Oxford, UK ³Exascale Research Computing Lab, Bruyeres-le-Chatel, France ⁴Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, Stanford, USA ⁵Kavli Institute for the Physics and Mathematics of the Universe, University of Tokyo, Kashiwa-shi, Japan ⁶Sub-department of Astrophysics, University of Oxford, Oxford, UK

MG14 - 17. July 2015



Motivation

- Strong lensing analyzes wide field surveys
- Robots are not very good at finding and modelling lenses
- Human intervention is needed!
- \Rightarrow SpaceWarps citizen science project (50'000 volunteers; 11 mio classifications; 51 candidates)

SpaceWarps Results and Outlook

- CHFT Legacy Survey: 150 deg²
- 59 candidates found (29 promising)¹
- $\Rightarrow \approx 1$ lens every few deg²
 - DES, PanStarrs; later LSST, Euclid
 - more area, better resolution
- \Rightarrow 10'000 lenses over 10 years (pprox one per hour)

¹[A. More et al; arXiv:1504.05587]

Outlook

A lot of computational and manpower needed

Detecting lenses:

- Robots making progress (RingFinder; ArcFinder) in combination with
- Future SpaceWarps runs

Post processing?

- Robots not there (yet?)
- ⇒ Citizen science: SpaghettiLens

_ Theory

Theory

Fermat's Principle

Fermat's Principle²

Rays of light traverse the path of stationary optical length with respect to variations of the path.

Fermat's Principle


Time t for path X:

$$t\left[X\right] = \frac{1}{c} \int_{t_1}^{t_2} n\left(\vec{x}\left(t\right)\right) \sqrt{1 + \left(\frac{\mathsf{d}\vec{x}\left(t'\right)}{\mathsf{d}t'}\right)^2} \mathsf{d}t'$$

Path X where t stationary.

²Ghatak, Ajoy (2009), Optics

Setup



Light Travel Time

Light travel time

$$t(x,y) = t_{geom} + t_{grav} \tag{1}$$

$$t_{\text{geom}} \propto (x - x_s)^2 + (y - y_s)^2$$
 (2)

$$t_{\text{grav}} = \langle t_{\text{grav}}(x_{\circ}, y_{\circ}) \rangle + (1 + z_{L}) \frac{2G}{c^{3}} M(x_{\bullet}, y_{\bullet})$$
 (3)

L Theory

$$A_t = A_{\text{geom}} + A_{\text{grav}} \tag{4}$$

$$A_{\text{geom}} = \frac{1}{2} \left(x^2 + y^2 \right) \tag{5}$$

$$\nabla^2 A_{\text{grav}}(x, y) = -2\kappa(x, y) \tag{6}$$

$$A = \frac{cD_L}{(1+z_L)^2} \frac{D_{LS}}{D_S} \times t \tag{7}$$

$$\kappa(x,y) = \frac{4\pi G}{c^2} \frac{D_L}{1+z_L} \frac{D_{LS}}{D_S} \times \Sigma(x,y)$$
 (8)

Alternative explanation





















































































Lensing Ga











































































Simulation program: gravlens by C. Huwiler $\,$


Arrival Time Surface
















Figure: ASW0004q9e (SpaceWarps)









Figure: ASW0004q9e (SpaceWarps)

SpaghettiLens¹

- Extremal Points (Images)
- Self Intersecting Contour Lines

http:

//labs.spacewarps.org/spaghetti/



L Theory

SpaghettiLens Results: Test of Performance

- Use simulated lenses
- Let volunteers model them
- Recover Einstein Radii
- Volunteers perform well!

arxiv:1502.00008

Mon. Not. R. Astron. Soc. 447, 2170-2180 (2015)

Printed February 3, 2015

(MN IATEX style file v2.2)

Gravitational Lens Modelling in a Citizen Science Context

Rafael Küng,¹ Prasenjit Saha,¹ Anupreeta More,² Elisabeth Baeten,³ Jonathan Coles,⁴ Claude Cornen,³ Christine Macmillan,³ Phil Marshall,⁵

Surhud More, ² Jonas Odermatt, ⁶ Aprajita Verma⁷ and Julianne K. Wilcox³

¹ Physik-Institut, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

³ Zooniverse, c/o Astrophysics Department, University of Oxford, Oxford OX1 3RH, UK

² Kavli Institute for the Physics and Mathematics of the Universe, University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa-shi 277-8583, Japan


 $^{^4\}textit{Exascale Research Computing Lab, Campus Teratec, 2 Rue de la Piquetterie, 91680 Bruyeres-le-Chatel, France}$

⁵ Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, 452 Lomita Mall, Stanford, CA 94035, USA 6 Kantonsschule Zug, Lüssiweg 24, 6300 Zug, Switzerland

⁷ Sub-department of Astrophysics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford, OX1 3RH, UK

SpaghettiLens Results: Stellar vs Lensing Mass

- Lensing mass against the stellar mass of the candidate lens galaxies
- Stellar mass fraction of order 20 percent
- With decreasing trend for the most massive galaxies
- Expected for early type galaxies
- Outliers? Maybe non-lenses (not yet spectroscopically confirmed)



Outlook

We are currently working on:

- Fit parametrized models to the free-form mass distributions (Lucy Oswald)
- Determination of photometric red shifts
- Estimate stellar populations (using galfit, SExtractor)

Questions?

Questions? rafael.kueng@uzh.ch

Appendix