

# Numerical Algorithms

## Fall 2020

---

### Assignment 6 (only for students of the 6 ECTS course) October 15, 2020

---

#### Exercise 1 [10 points]

Consider the  $n + 1$  interpolation points  $P_0, \dots, P_n$  from Exercise 2 of Assignment 4 and the uniform knot vector  $T = (t_0, \dots, t_{m+7})$  for  $m = n - 1$  with quadruple end knots, that is,

$$t_0 = t_1 = t_2 = t_3 = 0, \quad t_{i+3} = i, \quad i = 1, \dots, m, \quad t_{m+4} = t_{m+5} = t_{m+6} = t_{m+7} = m + 1.$$

Compute the cubic B-spline curve  $F_n: [t_0, t_{m+7}] \rightarrow \mathbb{R}^2$  over  $T$  that interpolates the  $P_i$  at  $t_{i+3}$ , that is,

$$F_n(t_{i+3}) = P_i, \quad i = 0, \dots, m + 1$$

and satisfies the natural end conditions

$$F''(t_3) = F''(t_{m+4}) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Plot the three curves  $F_{10}$ ,  $F_{20}$ , and  $F_{40}$  into three different pictures, together with the curve  $f$  from which the interpolation points were sampled (cf. Exercise 2 of Assignment 4), using  $M + 1$  sample points for each curve, where  $M = 10000$ .

*Note:* for plotting the curves, you can use the code of the de Boor algorithm provided on *iCorsi*.

*Hint:* use the derivative formula for B-spline curves (twice) and the de Boor algorithm to figure out how to express  $F''(t_3)$  and  $F''(t_{m+4})$  in terms of the  $D_i$ .

Make sure that your code is efficient and uses only  $O(n)$  operations, by taking advantage of the fact that it needs to solve a tridiagonal system

→ see [https://en.wikipedia.org/wiki/Tridiagonal\\_matrix\\_algorithm](https://en.wikipedia.org/wiki/Tridiagonal_matrix_algorithm)

Describe how you derived your code, and hand in your code and the three pictures with the plots.

#### Bonus Exercise [5 points]

Given the uniform knot vector  $T = (t_0, \dots, t_8)$  with  $t_i = i$  and the control points

$$D_0 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad D_1 = \begin{pmatrix} 6 \\ 0 \end{pmatrix}, \quad D_2 = \begin{pmatrix} 0 \\ 12 \end{pmatrix}, \quad D_3 = \begin{pmatrix} 12 \\ 12 \end{pmatrix}, \quad D_4 = \begin{pmatrix} 12 \\ 12 \end{pmatrix},$$

consider the cubic B-spline curve  $F(t) = \sum_{i=0}^4 D_i N_i^3(t)$ . This curve has two cubic pieces:  $F_0: [3, 4] \rightarrow \mathbb{R}^2$  and  $F_1: [4, 5] \rightarrow \mathbb{R}^2$ . Compute the Bézier control points  $C_{0,0}, \dots, C_{0,3}$  and  $C_{1,0}, \dots, C_{1,3}$  for the curves  $F_0$  and  $F_1$  and verify, using the A-frame condition, that they join indeed with  $C^2$ -continuity.

---

**Solutions must be returned online or in class on October 22, 2020**