Nonlinear Dynamics in Computer Graphics: Reaction–Diffusion Systems in Procedural Generation

Ragav Srinivasan

November 11, 2026

Abstract

We start by ti

1 Introduction

1.1 Structure of this report

Section 2 gives the background on reaction–diffusion systems and the phenomena they model. We then discuss a famous result of Alan Turing on reaction–diffusion systems, and finally focus on a simple reaction–diffusion system called the Gray–Scott model. Section 3 is my original contribution: a stability analysis of the Gray–Scott model and a numerical simulation of it. Section 4 gives the conclusion and possible extensions of this project.

2 Background

2.1 Procedural Generation

Procedural generation is a broad set of techniques in computer graphics that is used to create visuals, such as pseudo-realistic textures, terrains and surfaces on the fly. It eliminates the need for artists to manually prototype and create their own worlds. Instead, a deterministic function called *noise* is used to produce the desired visuals, on the fly as the player explores. This makes it possible to build large, *lifelike* landscapes with mountains, rivers and vegetation with little impact on memory (Lagae et al. 2010).

Procedural generation is based on many algorithms. In this section we study one of the most widely used algorithms: *Perlin noise*.

Ken Perlin devised his noise function after working on the visual effects for Disney's 1982 film *Tron* at Mathematical Applications Group, Inc. (MAGI), and published it in 1985 (Perlin 1985). His technique was so pivotal to the industry that it earned him an Academy Award for Technical Achievement (Perlin 2026), and it is still widely used in modern-day films and video games (Lagae et al. 2010).

How Perlin noise works. Need to write how perlin noise works next.

Building textures from noise. A single layer of noise is smooth but bland. Natural surfaces have detail at many scales, so copies of the noise at doubled frequency and halved amplitude are added together:

$$\mathrm{fBm}(\mathbf{x}) = \sum_{j=0}^{n-1} \frac{1}{2^j} \,\mathrm{noise}\big(2^j \mathbf{x}\big)\,, \tag{2.1}$$

which is known as fractal Brownian motion. Replacing noise by |noise| in Equation (2.1) gives a "turbulence" function, and feeding it into a simple formula produces a material. Perlin's classic marble, for example, is $\sin\big(x_1 + \text{turbulence}(\mathbf{x})\big)$ (Perlin 1985).

Limitations. Textures created with Perlin noise are random and have no structure (Lagae et al. 2010). Although this approach is ideal for terrain generation, it cannot be used to generate patterns found in biology such as coral reefs, zebra stripes, giraffe spots, etc. For this we use a different technique found in Reaction Diffusion Systems.

2.2 Reaction Diffusion Systems

This brings us to the heart of this report: *reaction-diffusion systems*. These are nonlinear dynamical systems, originally from chemistry, and model changes in concentrations of several chemical substances as they react and diffuse over space and time (Turing 1952).

Diffusion refers to the behaviour of chemical substances *spreading* from regions of high concentration to low concentration and as a result the substance is uniformly distributed across a medium (Encyclopædia Britannica 2026). It turns out that mathematically modelling this behaviour leads to an interesting variety of pattern formation called Turing patterns, which will be the focus of Section 2.3.

The general form of a reaction–diffusion system is

$$\frac{\partial \mathbf{q}}{\partial t} = D\nabla^2\mathbf{q} + R(\mathbf{q}), \tag{2.2}$$

Above is a dimensionless equation wherein all the physical quantities are stripped and all the terms scaled (Strogatz 2015).

2.3 Turing patterns

In 1952, towards the end of Alan Turing's life, Turing publishes one of his most seminal contributions in the field of mathematical biology. In his 1952 paper titled "The Chemical Basis of Morphogenesis", Turing discovers a mechanism which describes the formation of some patterns found in nature (Turing 1952). For instance, the patterns found in animal skin such as the stripes of a zebra, spots of a jaguar, the structure of our brains and the formation of coral reefs are considered to be Turing patterns (University of Sheffield 2022).

In the paper, Turing studies the development of an embryo and describes a system of signalling molecules called *morphogens*. The reaction and diffusion shown by these morphogens can destabilise the system leading to an interesting case of pattern formation (Turing 1952).

In the following section, we will be studying Turing patterns formed by a simple model called the Gray–Scott model.

2.4 The Gray–Scott model

The setup for the Gray–Scott model supposes that we have a *well-mixed tank* of chemical U. Then the concentration of U. We are interested in knowing the concentration of the chemical in space and time. For this, we define a vector function $u(x,t)$ that is currently unknown. We assume that chemical U has the same concentration in the spatial domain except for a small patch in the center of the tank where chemical V is introduced into the system.

In chemistry, the relationship between U and V is defined as *autocatalysis* where the resulting product is a reactant, as is the case with V in the reaction shown below. The relationship is akin to that of *prey and predator* models.

$$\underbrace{U + 2V \to 3V}_{V \text{ reacts with } U \text{ to make more of itself}} \qquad \underbrace{V \to P}_{\text{decay}}$$

The second rule suggests decay for V by turning into an inert product P which means that it becomes a non-reactive substance.


$$\frac{\partial U}{\partial t} = D_u \nabla^2 U - UV^2 + F(1 - U)$$
$$\frac{\partial V}{\partial t} = D_v \nabla^2 V + UV^2 - (F + k)V$$

- D_u and D_v are the diffusivities of chemicals U and V.
- U and V are the unknown functions representing the concentrations of the respective chemicals in space and time.
- F is the feed rate of chemical U.
- k is the decay rate for chemical V.

3 Original Contribution

3.1 Linear Stability Analysis of the Gray–Scott Model

Steady states of a dynamical system are states after which point the system shows no change. We consider a small disturbance around a steady state and say the state is *stable* if the disturbance dampens and the system loops back to its steady state. While an *unstable* steady state is when the disturbance grows and thus the system never returns back to its steady state (Teschl 2012). To give an analogy, Figure 1 compares a ball on a hilltop versus in a valley. Panel (a) is a stable state as the ball returns to its steady state after a small nudge. Panel (b) is an example of an unstable state, the ball never returns to a steady state after a small push and finally panel (c) shows a bistable state where the ball returns to a steady state but at a different point.



(a) Stable (b) Unstable (c) Bistable

Figure 1: An analogy for Stability Analysis of Steady States

Nonlinear systems such as the Gray-Scott model are difficult to solve analytically. Therefore, I will be doing a stability analysis to find out the behaviour of the system near its steady states. To do this, we will first linearise the system by its Jacobian matrix and classify by its trace and determinant.

In this analysis I will show how diffusion can *destabilise* a steady state and result in pattern formation. The analysis will be done twice, once by dropping the coefficients of diffusion D_u and D_v and then we will add them back to the system and check whether it destabilises the system.

$$\begin{aligned}\frac{\partial U}{\partial t} &= D_u \nabla^2 U + \underbrace{-UV^2 + F(1-U)}_{f(U,V)},\\ \frac{\partial V}{\partial t} &= D_v \nabla^2 V + \underbrace{UV^2 - (F+k)V}_{g(U,V)}.\end{aligned} \tag{3.1}$$

Stability Analysis without Diffusion

Setting $D_u = D_v = 0$ in Equation (3.1), every point in space will behave in the same way. We define $\dot{U}$ and $\dot{V}$:

$$\begin{aligned}\dot{U} &= f(U,V) = -UV^2 + F(1-U),\\ \dot{V} &= g(U,V) = UV^2 - (F+k)V.\end{aligned} \tag{3.2}$$

A steady state is when $f = g = 0$. Setting $g = 0$ and factorizing by V gives us $g = V(UV + (F+k)) = 0$. Here, we have two cases for V. If $V = 0$, then $U = 1$ which is our *trivial steady state*.
Let's first classify stability of the trivial steady state by the Jacobian of the system given by:

$$J = \begin{pmatrix} \frac{\partial f}{\partial U} & \frac{\partial f}{\partial V} \\ \frac{\partial g}{\partial U} & \frac{\partial g}{\partial V} \end{pmatrix} = \begin{pmatrix} -V^2 - F & -2UV \\ V^2 & 2UV - (F+k) \end{pmatrix}. \tag{3.3}$$

To classify the steady state, we first substituted the point into the above Jacobian. Then calculate, $p = trJ$ and $q = detJ$. The state is considered to be stable if $p < 0$ and $q > 0$.

$$J(1,0) = \begin{pmatrix} -F & 0 \\ 0 & -(F+k) \end{pmatrix}, \qquad p = -(2F+k), \qquad q = F(F+k), \qquad p^2 - 4q = k^2. \tag{3.4}$$

Since $F, k > 0$, we have $p < 0$, $q > 0$ and $p^2 - 4q \geq 0$, so the trivial state is a *stable node for every* F and k. This suggests that when a small concentration of chemical V is added to the tank mixed by U, the V dies out and the system is back to its original state.

Next, we consider the second case where $UV + (F+k) = 0 \rightarrow UV = (F+k)$. Substituting $UV = (F+k)$ in $f(U,V)$ gives the set of non-trivial steady states:

$$\begin{aligned} V &= \frac{F(1-U)}{F+k}, \qquad U^2 - U + c = 0 \quad \text{with } c = \frac{(F+k)^2}{F},\\ U_\pm &= \frac{1}{2}\left(1 \pm \sqrt{1 - \frac{4(F+k)^2}{F}}\right), \qquad V_\pm = \frac{F+k}{U_\pm}. \end{aligned} \tag{3.5}$$

The steady states only exist under the following condition:

$$F \geq 4(F+k)^2 \Rightarrow k \leq \tfrac{1}{2}\sqrt{F} - F. \tag{3.6}$$

Classifying the non-trivial states. Using $UV = F + k$,

$$J(U_\pm, V_\pm) = \begin{pmatrix} -V^2 - F & -2(F+k) \\ V^2 & F+k \end{pmatrix}, \qquad p = k - V^2, \qquad q = (F+k)(V^2 - F). \tag{3.7}$$

From the quadratic, $(F+k)^2 = FU(1-U)$, so $V^2 = (F+k)^2/U^2 = F(1-U)/U$ and $V^2 - F = F(1-2U)/U$. Since $U_+ > \frac{1}{2} > U_-$:

- (U_+, V_+) has $q < 0$, so it is a *saddle* (always unstable);
- (U_-, V_-) has $q > 0$, so it is stable when $p < 0$, i.e. $V_-^2 < k$. It loses stability when $p = 0$ with $q > 0$, a *Hopf bifurcation*, on the curve $V_-^2 = k$.

Adding back diffusion

3.2 Simulation of the Gray–Scott Model

3.3 Texture Mapping in Blender

[TODO: I will do this section if I have time, for now I will just do the parameter map and stability analysis.]

4 Conclusion

To conclude the patterns formed by Reaction Diffusion Systems and specifically the Gray Scott Model are very interesting

References

Encyclopædia Britannica (2026). *Diffusion*. URL: https://www.britannica.com/science/diffusion (visited on 11/11/2026).

Lagae, Ares et al. (2010). "A Survey of Procedural Noise Functions". In: *Computer Graphics Forum* 29.8, pp. 2579–2600. DOI: 10.1111/j.1467-8659.2010.01827.x.

Perlin, Ken (1985). "An Image Synthesizer". In: *Computer Graphics (Proceedings of SIGGRAPH '85)* 19.3, pp. 287–296. DOI: 10.1145/325165.325247.

— (2026). *Noise and Turbulence*. URL: https://mrl.cs.nyu.edu/~perlin/doc/oscar.html (visited on 11/11/2026).

Strogatz, Steven H. (2015). *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemistry, and Engineering*. 2nd ed. Boulder, CO: Westview Press.

Teschl, Gerald (2012). *Ordinary Differential Equations and Dynamical Systems*. Vol. 140. Graduate Studies in Mathematics. Providence, RI: American Mathematical Society. URL: https://www.mat.univie.ac.at/~gerald/ftp/book-ode/.

Turing, Alan M. (1952). "The Chemical Basis of Morphogenesis". In: *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 237.641, pp. 37–72. DOI: 10.1098/rstb.1952.0012.

University of Sheffield (2022). *Turing Patterns*. Turing Pattern Project. URL: https://turing-pattern-project.sites.sheffield.ac.uk/turing-patterns (visited on 11/11/2026).

5 Project Plan

6 Reflection

When I started working on this project, I had set myself some high expectations on this project. In the end, the plan I gave myself turned out to be unrealistic.

I chose this topic because of my interest in Computer Graphics enjoy its use for mathematics and found Reaction Diffusion Systems to be link between nonlinear dynamics and computer graphics. In hindsight, I'm not sure if it turned out to be a good idea do something that interested as it created a lot of bias in the research.

I planned the report to be a study on a specific Reaction Diffusion model called the Gray Scott Model. For the original contribution, I wanted to do a stability analysis and my own simulation

of the Grey Scott Model using a method described by J.E Pearson in his paper (insert paper title.). However, when i started doing the stability analysis for the Gray Scott Model with Diffusion I personally found it to be very difficult. So in my report, I've only been able to do a stability analysis of a single aspect of the model.

I planned my report to be a study on Turing Patterns formed by Reaction Diffusion Systems. There is a fair amount of literature on the topic of Turing Patterns including Turing's discovery of these patterns in 1952, J.E Pearson's numerical simulation on the Gray Scott Model.

I'm not very proud with the current state of my report and feel like there is a lot left to be desired. Although, I have stopped working on this report. I find the topic fascinating so I've made the commitment to keep learning more about this topic. Specifically, I want to fill the gap in my stability analysis in which I couldn't complete the part of adding back the diffusion coefficients.

On a positive note, Doing this research project has given me a better grasp of Nonlinear Dynamics and the numerical methods for solving a system of Partial Differential Equations that I learned last semester.

7 Appendix