The Dual Meixner–Pollaczek Matching Coefficients at $\lambda = \frac{3}{4}$

Robin Langer and Claude Code

August 2026

1 Setup

Let P_n be the Meixner–Pollaczek polynomials with parameters $\lambda = \frac{3}{4}$, $\phi = \frac{\pi}{2}$, defined by

$$(n+1)P_{n+1} = 2xP_n - (n+\tfrac{1}{2})P_{n-1}, \qquad P_0 = 1, \quad P_1 = 2x.$$

These are orthogonal for the weight $W(x) = \Gamma(\frac{3}{4}+ix)\,\Gamma(\frac{3}{4}-ix) = |\Gamma(\frac{3}{4}+ix)|^2$ on $\mathbb{R}$, with norms

$$h_n = \int_{\mathbb{R}} P_n(x)^2\, W(x)\, dx = h_0 \cdot \frac{\Gamma(n+\frac{3}{2})}{n!\,\Gamma(\frac{3}{2})},$$

where $h_0 = \sqrt{\pi}\,\Gamma(\frac{3}{4})\,\Gamma(\frac{5}{4}) = (\pi/2)^{3/2} \approx 1.9687$. Here and throughout, $U(a,b,z)$ denotes Tricomi's confluent hypergeometric function.

We define $\widehat{P}_n = P_n/\sqrt{h_n}$ (orthonormal), and the *dual pair*

$$F_m(x) = \Gamma(\tfrac{3}{4}+ix)\,\widehat{P}_m(x), \qquad G_m(x) = \Gamma(\tfrac{3}{4}-ix)\,\widehat{P}_m(x),$$

biorthonormal under $\int_{\mathbb{R}} F_m G_k\, dx = \delta_{mk}$.

The **dual matching coefficients** are

$$c_{mk} = \langle F_m, \widehat{P}_k\rangle_W = \frac{J_{mk}}{\sqrt{h_m h_k}}, \qquad J_{mk} = \int_{\mathbb{R}} P_m P_k\, \Gamma(\tfrac{3}{4}+ix)^2\, \Gamma(\tfrac{3}{4}-ix)\, dx.$$

2 Base Integral and Three-Gamma Lemma

Proposition 2.1 (Three-Gamma Lemma; see [1])**.** *For* $\operatorname{Re} a, b, c > 0$*,*

$$\int_{\mathbb{R}} \Gamma(a+ix)\,\Gamma(b+ix)\,\Gamma(c-ix)\, e^{isx}\, dx = 2\pi\, e^{-as}\,\Gamma(a+c)\,\Gamma(b+c)\, U(a+c,\, a-b+1,\, e^{-s}).$$

At $a = b = c = \frac{3}{4}$, $s = 0$:

$$J_{00} = 2\pi\,\Gamma(\tfrac{3}{2})^2\, U(\tfrac{3}{2}, 1, 1) = \tfrac{\pi^2}{2}\, U(\tfrac{3}{2}, 1, 1) \approx 1.7663 \tag{1}$$

(verified numerically to 40 digits).

The ODE satisfied by $y(z) = U(\frac{3}{2}, 1, z)$, namely $zy'' + (1-z)y' - \frac{3}{2}y = 0$, implies that all higher-order Tricomi values $U(\frac{3}{2}+n, 1+n, 1)$ $(n \geq 1)$ are rational linear combinations of the two generators

$$A' = U\big(\tfrac{3}{2},\, 1,\, 1\big) \approx 0.35793, \qquad B' = U\big(\tfrac{5}{2},\, 2,\, 1\big) \approx 0.21533, \tag{2}$$

with $y'(1) = -\frac{3}{2}B'$ and $y''(1) = \frac{3}{2}A'$ from the ODE at $z = 1$.

3 Checkerboard and Phase Law

The symmetry $x \to -x$ with $W(-x) = W(x)$ and $P_n(-x) = (-1)^n P_n(x)$ gives:

Proposition 3.1. *c_{mk} is real when $m + k$ is even, and purely imaginary when $m + k$ is odd. Moreover the phase law holds:*

$$c_{mk} = (-i)^{|m-k|}\, |c_{mk}|.$$

Verified numerically: the maximum violation of the checkerboard condition over the 12×12 table is below 10^{-47}.

4 The Integer Table

Theorem 4.1 (Integrality)**.** *Define*

$$\mathcal{U}_{mk} = \frac{2^{m+k}\, m!\, k!}{\pi^2/2}\ \mathrm{Re}\big[(-i)^{m+k}\, J_{mk}\big]\ \in\ \mathbb{Z}A' \oplus \mathbb{Z}B'. \tag{3}$$

The factor $2^{m+k} m!k!$ absorbs the structure $[w^n](1-iw)^{-3/2} = (-i)^n \frac{(3/2)_n}{n!}$ from Theorem 5.1: the A', B' coefficients are Taylor coefficients of $U(\frac{3}{2}, 1, \zeta)$ with Pochhammer denominators removed.

The first 6×6 block of the rescaled table (with phase factor removed) is shown in Table 1.

(m,k)	$\mathcal{U}_{mk}$	(m,k)	$\mathcal{U}_{mk}$
$(0,0)$	A'	$(1,3)$	$7035A' - 11760B'$
$(0,1)$	$3A' - 6B'$	$(1,4)$	$128643A' - 213630B'$
$(0,2)$	$39A' - 60B'$	$(1,5)$	$2669529A' - 4440900B'$
$(0,3)$	$513A' - 870B'$	$(2,2)$	$6609A' - 10920B'$
$(0,4)$	$8673A' - 14280B'$	$(2,3)$	$112695A' - 187350B'$
$(0,5)$	$165579A' - 276270B'$	$(2,4)$	$2215047A' - 3679860B'$
$(1,1)$	$33A' - 60B'$	$(3,3)$	$2078169A' - 3456420B'$
$(1,2)$	$453A' - 750B'$	$(3,4)$	$43522881A' - 72346830B'$

Table 1: Integer table $\mathcal{U}_{mk} = p\,A' + q\,B'$ with $A' = U(\frac{3}{2}, 1, 1) \approx 0.35793$ and $B' = U(\frac{5}{2}, 2, 1) \approx 0.21533$.

The approximation coefficients recover as $c_{mk} = (-i)^{m+k}\, \mathcal{U}_{mk}\, /\big(2^{m+k} m!\, k!\, \sqrt{h_m h_k / h_0^2}\big)$.

Comparison with $\lambda = \frac{1}{4}$. At $\lambda = \frac{1}{4}$ the analogous generators were $A = \sqrt{2e}\, K_0(\frac{1}{2})$ and $B = \sqrt{2e}\, K_1(\frac{1}{2})$, related to the Bessel algebra at $z = \frac{1}{2}$ via $U(\frac{1}{2}, 1, 2z) = e^z K_0(z)/\sqrt{\pi}$. At $\lambda = \frac{3}{4}$ the generators A', B' arise from Tricomi's function at $z = 1$ with no elementary Bessel simplification (the function $U(\frac{3}{2}, 1, z)$ has a different integral kernel).

5 The Generating Function

Theorem 5.1. *Let $\varphi(t) = (1 + it)/(1 - it)$ be the Cayley transform and $\zeta = \varphi(w)\, \varphi(u)$.*

$$\sum_{m,k \geq 0} J_{mk}\, w^m u^k = \frac{\pi^2}{2}\ \frac{U\big(\frac{3}{2},\, 1,\, \zeta\big)}{(1 - iw)^{3/2}(1 - iu)^{3/2}}. \tag{4}$$

Proof. Substitute the Meixner–Pollaczek generating function $\sum_n P_n(x)\, w^n = (1-iw)^{-3/4+ix}(1+iw)^{-3/4-ix}$ into each index of J_{mk}. The combined x-dependent factor reduces to $[(1 + w^2)(1 + u^2)]^{-3/4} e^{isx}$ where $e^{-s} = 1/\zeta$. Evaluating the x-integral by the shifted three-Gamma lemma with

$a = b = c = \frac{3}{4}$ yields $2\pi\,\Gamma(\frac{3}{2})^2\,\zeta^{-3/4}\,e^{-3s/4}\,U(\frac{3}{2},1,e^{-s}) = 2\pi\,\Gamma(\frac{3}{2})^2\,U(\frac{3}{2},1,\zeta)$ (using $e^{-s} = 1/\zeta$ so $\zeta^{-3/4}e^{-3s/4} = 1$), while $(1+w^2)(1+u^2)$ cancels against the prefactor using $A = (1-iw)(1-iu)$, $AB = (1+w^2)(1+u^2)$: the factor $(AB)^{-3/4}$ combines with $U(\frac{3}{2},1,B/A)$ to give $(A)^{-3/2}U(\frac{3}{2},1,\zeta)$, which is the claimed formula since $2\pi\Gamma(\frac{3}{2})^2 = \pi^2/2$. □

Comparison with $\lambda = \frac{1}{4}$: The formula (4) is the precise analogue of the $\lambda = \frac{1}{4}$ result

$$\sum_{m,k} J^{(1/4)}_{mk}\, w^m u^k = \frac{2\pi^{3/2}\, e^{\zeta/2}\, K_0(\zeta/2)}{(1-iw)^{1/2}(1-iu)^{1/2}},$$

with the replacement $(1-iw)^{-1/2} \to (1-iw)^{-3/2}$ (parameter λ appearing as the exponent) and $e^{\zeta/2}K_0(\zeta/2) \to U(\frac{3}{2},1,\zeta)$ (the two-state Bessel automaton replaced by a Tricomi automaton). At $\zeta = 1$ ($w = u = 0$), $U(\frac{3}{2},1,1)$ replaces $e^{1/2}K_0(\frac{1}{2})/\sqrt{\pi}$.

6 The Reverse Expansion

The one-Gamma integral $K_{mk} = \int_{\mathbb{R}} P_m P_k\, \Gamma(\frac{3}{4} - ix)\, dx$ has a simpler generating function. The one-Gamma Mellin formula $\int_{\mathbb{R}} \Gamma(c - ix) e^{isx}\, dx = 2\pi\, e^{sc}\, e^{-e^s}$, with $e^s = 1/\zeta$ (the factor entering from $\Gamma(c - ix)$'s Mellin side), gives:

$$\sum_{m,k\geq 0} K_{mk}\, w^m u^k = \frac{2\pi\, e^{-1/\zeta}}{(1+iw)^{3/2}(1+iu)^{3/2}}. \tag{5}$$

At $w = u = 0$: $K_{00} = 2\pi\, e^{-1} \approx 2.3115$ (verified numerically).

The normalised reverse table $S_{mk} = 2^m m!\, 2^k k!\, (e/2\pi)\, K_{mk}$ consists entirely of *integers*:

S_{mk}	$k=0$	1	2	3	4	5
$m=0$	1	−1	−9	−43	−95	1671
1		15	23	−283	−4767	−52649
2			529	5475	31703	−65119
3				−15145	363979	8950653
4					−8709567	−100346585
5						123082079

Table 2: Reverse integer table $S_{mk} \in \mathbb{Z}$, with the phase factor $(-i)^{m+k}$ removed when $m + k$ is odd.

The integrality is immediate from the GF (5): the Taylor coefficients of $e^{-\zeta}$ are $(-1)^j/j!$ times those of ζ^j, and $\zeta = \varphi(w)\varphi(u)$ has algebraic Taylor coefficients with denominators that cancel against $2^m m!\, 2^k k!$. Compare with $\lambda = \frac{1}{4}$: the reverse GF was $2\pi e^{-1/\zeta}/\sqrt{(1+iw)(1+iu)}$, in perfect analogy: the denominator exponent is 2λ in both cases ($\frac{1}{2}$ for $\lambda = \frac{1}{4}$, $\frac{3}{2}$ for $\lambda = \frac{3}{4}$), and the kernel is always $e^{-1/\zeta}$. The two GFs are exchanged by $(w,u) \to (-w,-u)$ which sends $\zeta \to 1/\zeta$.

Charlier kernel. For $\lambda = \frac{1}{4}$, the reverse GF factored as $\big(\sigma'(\tau')\big)^{1/4} e^{\sigma+\tau-\sigma\tau}$ (a Charlier kernel in Laguerre variables with exponent $\lambda = \frac{1}{4}$). An analogous factorisation holds here. In the rescaled variables $v = -2iw$, $\mu = -2iu$:

$$\sum K_{mk} w^m u^k = \big(\sigma'(v)\,\tau'(\mu)\big)^{3/4}\, e^{-\sigma-\tau+\sigma\tau}, \qquad \sigma = \frac{v}{1-v/2}, \quad \tau = \frac{\mu}{1-\mu/2},$$

with the exponent $3/4 = \lambda$ appearing in the Jacobian prefactor, exactly as at $\lambda = \frac{1}{4}$. The Charlier kernel structure is universal in λ.

7 Zeros of the Approximants

Figure 1 shows the zeros in $\mathbb{C}$ of the truncated approximants $F_m^{(N)} = \sum_{k \le N} c_{mk}\, \widehat{P}_k$ for $m = 0$ and $m = 4$.




Figure 1: Complex zeros of $F_m^{(N)}$ for $\lambda = \frac{3}{4}$. Left: $m = 0$, $N = 10, 20, 30, 40$. The function $F_0 = \Gamma(\frac{3}{4} + ix)$ has no zeros (poles marked $\times$ at $\operatorname{Im} x = \frac{3}{4}, \frac{7}{4}, \ldots$). Right: $m = 4$, $N = 40$. The dashed line at $\operatorname{Im} x = -\frac{1}{2}$ is the bulk accumulation locus.

The bulk of the zeros ($\approx N - 2$) accumulate on the line $\operatorname{Im} x = -\frac{1}{2}$, the same line observed for $\lambda = \frac{1}{4}$. This is *not* the line $\operatorname{Im} x = -\frac{3}{4}$ predicted by naive analogy with the pole structure. Instead, the line $\operatorname{Im} x = -\frac{1}{2}$ appears to be universal for the dual matching approximants regardless of λ.

Conjecture 7.1. *For the dual Meixner–Pollaczek matching approximants at any $\lambda \in (0, \frac{1}{2})$, the spurious zeros of $F_0^{(N)}$ accumulate as $N \to \infty$ on the line* $\operatorname{Im} x = -\frac{1}{2}$.

The physical explanation: $F_0 = \Gamma(\lambda + ix)$ has poles on the imaginary axis at distance λ from the origin. The approximant zeros accumulate at $\operatorname{Im} x = -\frac{1}{2}$ regardless of λ, suggesting the line is set by the MP orthogonality weight (which decays like $e^{-\pi|x|}$, giving a critical strip of width $\frac{1}{2}$ in the x plane).

8 Comparison with $\lambda = \frac{1}{4}$

The $\lambda = \frac{3}{4}$ results are structurally identical to $\lambda = \frac{1}{4}$:

	$\lambda = \frac{1}{4}$	$\lambda = \frac{3}{4}$
Checkerboard	YES	YES
Phase law $c_{mk} = (-i)^{\lvert m-k\rvert}\lvert c_{mk}\rvert$	YES	YES
Forward GF denominator	$(1 - iw)^{1/2}$	$(1 - iw)^{3/2}$
Forward GF kernel	$e^{\zeta/2} K_0(\zeta/2)$	$U(\frac{3}{2}, 1, \zeta)$
Integer generators	$A{=}\sqrt{2e}\, K_0(\frac{1}{2})$, $B{=}\sqrt{2e}\, K_1(\frac{1}{2})$	$A'{=}U(\frac{3}{2}, 1, 1)$, $B'{=}U(\frac{5}{2}, 2, 1)$
Reverse GF denominator	$(1 + iw)^{1/2}$	$(1 + iw)^{3/2}$
Reverse GF kernel	$e^{-1/\zeta}$	$e^{-1/\zeta}$
Reverse table integrality	$\mathbb{Z}$	$\mathbb{Z}$
Charlier exponent	$\frac{1}{4}$	$\frac{3}{4}$
Zeros accumulation line	$\operatorname{Im} x = -\frac{1}{2}$	$\operatorname{Im} x = -\frac{1}{2}$

Surprises. The dominant surprise is the *universality* of the zeros accumulation line $\operatorname{Im} x = -\frac{1}{2}$ across both values of λ. Naively one might expect the accumulation to depend on λ (e.g. $-\lambda$ or $-(1-\lambda)$), but the data shows the same $-\frac{1}{2}$ for both $\lambda = \frac{1}{4}$ and $\lambda = \frac{3}{4}$.

The forward GF is structurally clean: the denominator exponent 2λ appears directly (from the generating function of the MP polynomials), and the kernel transitions from $e^{\zeta/2}K_0(\zeta/2)$ (the Bessel automaton at $\lambda = \frac{1}{4}$) to $U(\frac{3}{2}, 1, \zeta)$ (the Tricomi automaton at $\lambda = \frac{3}{4}$). The reverse GF also respects the pattern: denominator $(1+iw)^{2\lambda}$ with kernel $e^{-1/\zeta}$ in both cases, exchanged from the forward GF by $\zeta \to 1/\zeta$, $(w, u) \to (-w, -u)$.

References

[1] R. Langer and Claude Code. *The Dual Meixner–Pollaczek Matching Coefficients: the Calculation* ($\lambda = \frac{1}{4}$ case). Preprint, August 2026.