Sturmian sequences of $\sqrt{n}$: the proofs

Robin Langer

This is the proof-dense companion to `sturmian.tex`. Every claim made there is either proved here or listed in §10 as unproved. Two claims that `sturmian.tex` states for $n = 2$ and $n = 7$ alone are proved for every n in §8, and one statement there is false as written; §6 gives the counterexample and the repair.

1 Codings

Fix an irrational $\alpha \in (0,1)$. Write $\{y\}$ for $y - \lfloor y \rfloor$ and, for $x \in [0,1)$,

$$x_k \;=\; \{x + k\alpha\}, \qquad k \geq 0.$$

Definition 1. The *lower* and *upper codings* of x are the sequences

$$\mathcal{C}(\alpha,x)_k \;=\; \begin{cases} 1 & x_k \in [1-\alpha,\,1) \\ 0 & \text{otherwise,} \end{cases} \qquad\qquad \mathcal{C}^+(\alpha,x)_k \;=\; \begin{cases} 1 & x_k \in (1-\alpha,\,1) \text{ or } x_k = 0 \\ 0 & \text{otherwise.} \end{cases}$$

The two codings read the same circle with the same rotation and differ only in which endpoint of the arc of length α carries the letter 1.

Lemma 2. *For $x \in [0,1)$,*

$$\mathcal{C}(\alpha,x)_k \;=\; \lfloor (k+1)\alpha + x \rfloor - \lfloor k\alpha + x \rfloor, \qquad \mathcal{C}^+(\alpha,x)_k \;=\; \lceil (k+1)\alpha + x \rceil - \lceil k\alpha + x \rceil.$$

Proof. Put $m = \lfloor k\alpha + x \rfloor$ and $f = x_k$, so $k\alpha + x = m + f$ and $(k+1)\alpha + x = m + f + \alpha$ with $f + \alpha \in [0,2)$. Hence $\lfloor (k+1)\alpha + x \rfloor - m = \lfloor f + \alpha \rfloor$, which is 1 exactly when $f \geq 1 - \alpha$.

For the second, $\lceil m + f \rceil = m$ when $f = 0$ and $m + 1$ otherwise, so $\lceil m + f + \alpha \rceil - \lceil m + f \rceil = 1$ exactly when $f = 0$ or $f + \alpha > 1$. □

So $\mathcal{C}(\alpha,x)$ is the sequence of `sturmian.tex` §2 with $\rho = x$.

Lemma 3. *$\mathcal{C}(\alpha,x)$ and $\mathcal{C}^+(\alpha,x)$ are equal unless $x_{k_0} = 0$ for some $k_0 \geq 0$, which happens for at most one k_0. In that case they differ exactly at the positions $k_0 - 1$ and k_0, where they read `10` and `01`.*

Proof. The letter rules differ only at $x_k \in \{0,\, 1-\alpha\}$. If $x_k = 1-\alpha$ then $x_{k+1} = 0$, and if $x_{k_0} = 0$ with $k_0 \geq 1$ then $x_{k_0-1} = 1 - \alpha$. Two indices with $x_k = 0$ would give $(j-k)\alpha \in \mathbb{Z}$. □

Lemma 4. *For $x \in [0,1)$ and $N \geq 1$ the number of 1s among the first N letters of $\mathcal{C}(\alpha,x)$ is $\lfloor N\alpha + x \rfloor$.*

Proof. Sum Lemma 2 over $k < N$ and use $\lfloor x \rfloor = 0$. □

Corollary 5. *The density of 1s in $\mathcal{C}(\alpha,x)$ is α. The same holds for $\mathcal{C}^+(\alpha,x)$.*

Proof. $\lfloor N\alpha + x \rfloor / N \to \alpha$. The upper coding differs in at most two positions. □

The density is called the *slope*. It is a property of the sequence alone, so two codings that are equal as sequences have equal slopes. Every use of "solve for α" below rests on this corollary.

Lemma 6. *$\mathcal{C}(\alpha, x)_{k+1} = \mathcal{C}(\alpha, \{x + \alpha\})_k$, and likewise for $\mathcal{C}^+$.*

Proof. $\{x + (k+1)\alpha\} = \{\{x + \alpha\} + k\alpha\}$. □

2 The orbit is dense

Lemma 7. *For irrational α and any $x \in [0, 1)$, the set $\{x_k : k \geq 0\}$ is dense in $[0, 1)$.*

Proof. Let $\varepsilon > 0$. By Dirichlet there are integers $q \geq 1$ and p with $|q\alpha - p| < 1/q < \varepsilon$; put $\delta = |q\alpha - p| > 0$. The points x_0, x_q, x_{2q}, … differ consecutively by $\pm\delta$ modulo 1, with a fixed sign, so the first $\lceil 1/\delta \rceil + 1$ of them meet every interval of length δ. Take $\delta < \varepsilon$. □

Corollary 8. *$x \mapsto \mathcal{C}(\alpha, x)$ is injective on $[0, 1)$.*

Proof. Let $x < y$ and suppose the codings agree. By Lemma 4, $\lfloor N\alpha + x \rfloor = \lfloor N\alpha + y \rfloor$ for every $N \geq 1$. Choose N with $\{N\alpha + x\} > 1 - (y - x)$; then $N\alpha + x < \lceil N\alpha + x \rceil \leq N\alpha + y$. □

3 Complexity

Definition 9. A *factor* of length m of a sequence ℓ is a block $\ell_k \ell_{k+1} \cdots \ell_{k+m-1}$. Let $F_\ell(m)$ be the number of distinct factors of length m. The sequence is *Sturmian* when $F_\ell(m) = m + 1$ for every $m \geq 1$.

Theorem 10. *$F_{\mathcal{C}(\alpha,x)}(m) = m + 1$ for every $m \geq 1$.*

Proof. For $j \geq 1$ and $t \in [0, 1)$,

$$\lfloor t + j\alpha \rfloor \;=\; \lfloor j\alpha \rfloor + \begin{cases} 1 & t \geq \{-j\alpha\} \\ 0 & t < \{-j\alpha\}, \end{cases}$$

because $t + j\alpha = \lfloor j\alpha \rfloor + (\{j\alpha\} + t)$ with $\{j\alpha\} + t \in [0, 2)$, and $\{j\alpha\} + t \geq 1$ exactly when $t \geq 1 - \{j\alpha\} = \{-j\alpha\}$.

Let w be the factor of length m at position k. By Lemma 6 its letters are $w_j = \mathcal{C}(\alpha, x_k)_j$ for $j < m$, so by Lemma 4 the partial sums of w give $\lfloor x_k + j\alpha \rfloor$ for every $j \leq m$. By the display, w therefore determines, for each $j \in \{1, \ldots, m\}$, whether $x_k \geq \{-j\alpha\}$; and conversely those m comparisons determine w.

The m points $\{-j\alpha\}$, $1 \leq j \leq m$, are distinct and lie in $(0, 1)$, so they cut $[0, 1)$ into $m + 1$ intervals of positive length. Factors of length m are in bijection with those intervals that contain some x_k, and by Lemma 7 every one of them does. □

Corollary 11. *$F_{\mathcal{C}^+(\alpha,x)}(m) = m + 1$ for every $m \geq 1$.*

Proof. One can rerun the proof above with $\lceil \cdot \rceil$ for $\lfloor \cdot \rfloor$: the comparisons become $x_k > \{-j\alpha\}$ and the intervals are half-open at the other end. But that is more work than is needed, because an upper coding is the bitwise complement of a lower coding:

$$\mathcal{C}^+(\alpha, x)_k + \mathcal{C}(1 - \alpha, -x)_k = 1 \qquad (k \geq 0).$$

Indeed $\mathcal{C}^+(\alpha,x)_k = 1$ exactly when $\{x+k\alpha\} \in (1-\alpha,1)\cup\{0\}$, whose complement in $[0,1)$ is $(0,1-\alpha]$; and $\{-x+k(1-\alpha)\} = \{-(x+k\alpha)\}$, so the reflection $v \mapsto 1-v$ sends the orbit of x at slope α to the orbit of $-x$ at slope $1-\alpha$, carrying $(0,1-\alpha]$ onto $[\alpha,1) = [1-(1-\alpha),1)$ — the arc of the lower coding of slope $1-\alpha$. The exceptional point $v=0$ is fixed by the reflection and lies outside $[\alpha,1)$, as it must. Complementation is a bijection on words, and $1-\alpha$ is again irrational and in $(0,1)$, so Theorem 10 gives the count. □

The second proof is the one formalised. In `lean/Sturmian.lean` the displayed identity is `codeUp_add_code_reflect`, and the corollary is `ncard_factors_codeUp`. The first proof would have cost a parallel development: the natural range of the base point for an upper coding is $(0,1]$ rather than $[0,1)$, since $\mathcal{C}^+$ reads $\{x+k\alpha\}=0$ as the letter 1.

For the Fibonacci sequence of `sturmian.tex` §1 and $m=3$ the four intervals are cut by $\{-\alpha\},\{-2\alpha\},\{-3\alpha\}$ with $\alpha = 1/\varphi$, and the four factors are `101`, `011`, `110`, `010`. That is the table in §1.

4 Gaps

Write

$$\frac{1}{\alpha} \;=\; a_1+\beta, \qquad a_1 = \lfloor 1/\alpha \rfloor \geq 1, \qquad \beta = \{1/\alpha\} \in (0,1).$$

Then $a_1\alpha < 1 < (a_1+1)\alpha$, which is the inequality of `sturmian.tex` §3.

Let $I = [1-\alpha,1)$, the arc carrying the letter 1, and coordinatise it by

$$x \;=\; (1-\alpha)+\alpha u, \qquad u \in [0,1).$$

Theorem 12. *Let $x \in I$ with coordinate u. The least $t \geq 1$ with $\{x+t\alpha\} \in I$ is*

$$t \;=\; \begin{cases} a_1+1 & u < \beta \\ a_1 & u \geq \beta, \end{cases}$$

and the coordinate of $\{x+t\alpha\}$ is $\{u+1-\beta\}$.

Proof. $x+\alpha \in [1,1+\alpha)$, so $\{x+\alpha\} = x+\alpha-1 = \alpha u$, and for $s \geq 1$,

$$\{x+s\alpha\} \;=\; \alpha(u+s-1) \qquad \text{as long as } \alpha(u+s-1) < 1.$$

Now $\alpha(u+s-1) < 1 \iff s-1 < 1/\alpha - u = a_1+\beta-u$, and $\alpha(u+s-1) \geq 1-\alpha \iff u+s \geq a_1+\beta$. So $\{x+s\alpha\} \in I$ exactly when

$$a_1+\beta-u \;\leq\; s \;<\; a_1+\beta-u+1,$$

whose unique integer solution is $s = \lceil a_1+\beta-u \rceil$. For $u < \beta$ that is a_1+1 and for $u \geq \beta$ it is a_1; in both cases s is the least such t because the displayed formula for $\{x+s\alpha\}$ is valid and below $1-\alpha$ for all smaller s.

The new coordinate is $(\alpha(u+t-1)-(1-\alpha))/\alpha = u+t-1/\alpha$, which is $u+1-\beta$ when $t = a_1+1$ and $u-\beta$ when $t = a_1$. □

Corollary 13. *In $\mathcal{C}(\alpha,x)$ the gap between consecutive 1s is a_1 or a_1+1, so the number of 0s between them is a_1-1 or a_1.*

Definition 14. Let $k_0 < k_1 < k_2 < \cdots$ be the positions of the 1s of ℓ. The *derived sequence* ℓ' is

$$\ell'_i \;=\; \begin{cases} 1 & a_1 \text{ zeros lie between the } i\text{th and } (i{+}1)\text{th 1} \\ 0 & a_1 - 1 \text{ do.} \end{cases}$$

By Theorem 12, $\ell'_i = 1$ exactly when the coordinate u_i of x_{k_i} satisfies $u_i < \beta$, and

$$u_{i+1} \;=\; \{u_i + 1 - \beta\}.$$

Lemma 15 (block decomposition)**.** *Define* $\tau_a(1) = \mathtt{1}\,\mathtt{0}^a$ *and* $\tau_a(0) = \mathtt{1}\,\mathtt{0}^{a-1}$*. Then*

$$\ell \;=\; \mathtt{0}^{k_0} \cdot \tau_{a_1}(\ell').$$

Proof. The ith block of ℓ starting at k_i is a 1 followed by the zeros before the next 1, of which there are a_1 if $\ell'_i = 1$ and $a_1 - 1$ if $\ell'_i = 0$. □

Deriving and τ_{a_1} are inverse to each other up to the prefix $\mathtt{0}^{k_0}$. That prefix is the only reason a shift appears anywhere below.

5 The derivation theorem

Theorem 16. *Let* u_0 *be the coordinate of* x_{k_0}*, the first* 1*. Then*

$$\mathcal{C}(\alpha, x)' \;=\; \mathcal{C}^+(\beta,\, \{-u_0\}), \qquad\qquad \mathcal{C}^+(\alpha, x)' \;=\; \mathcal{C}(\beta,\, 1 - u_0).$$

Proof. For the lower coding put $v_i = \{-u_i\}$. By Theorem 12, $v_{i+1} = \{v_i + \beta\}$, so $v_i = \{v_0 + i\beta\}$. And

$$\ell'_i = 1 \iff u_i \in [0, \beta) \iff -u_i \in (-\beta, 0] \iff v_i \in (1 - \beta, 1) \text{ or } v_i = 0,$$

which is the letter rule of $\mathcal{C}^+(\beta, v_0)$.

For the upper coding the arc is $I^+ = (1 - \alpha, 1]$, coordinatised by $u \in (0, 1]$ with $u = 1$ meaning $x \equiv 0$. The computation of Theorem 12 is unchanged except that $\{x + s\alpha\} \in I^+$ reads $u + s > a_1 + \beta$ and $u + s - 1 \le a_1 + \beta$, so $t = a_1 + 1$ exactly when $u \le \beta$. Put $v_i = 1 - u_i \in [0, 1)$; then $v_{i+1} = \{v_i + \beta\}$ and

$$\ell'_i = 1 \iff u_i \le \beta \iff v_i \ge 1 - \beta,$$

which is the letter rule of $\mathcal{C}(\beta, v_0)$. □

Deriving sends the slope α to $\beta = \{1/\alpha\}$ and exchanges the two codings. The exchange is invisible for all but countably many x (Lemma 3), and it is exactly what makes the statement of Theorem 16 an equality rather than an equality off a small set.

Corollary 17. *The derived sequence of a coding is a coding, hence Sturmian, and its slope is* $\{1/\alpha\}$*.*

Corollary 18. *The slopes of* $\ell, \ell', \ell'', \ldots$ *are the orbit of* α *under the Gauss map* $\alpha \mapsto \{1/\alpha\}$*, so the block widths* a_1 *encountered are the partial quotients of* $\alpha = [\,0;\, a_1, a_2, a_3, \ldots\,]$*.*

6 Morse and Hedlund

One direction is Theorem 10: every coding with irrational slope is Sturmian. The statement in `sturmian.tex` §2 asserts the converse in the form

$$\ell_k \;=\; \lfloor (k+1)\alpha + \rho \rfloor - \lfloor k\alpha + \rho \rfloor, \qquad \rho \in [0,1),$$

that is, $\ell = \mathcal{C}(\alpha, \rho)$. In that form it is false.

Let $\alpha = 1/\varphi = (\sqrt{5}-1)/2$ and let $c = \mathcal{C}(\alpha, \alpha)$, the Fibonacci sequence. Let $\mathtt{1}c$ be c with a 1 prefixed.

Theorem 19. *$\mathtt{1}c = \mathcal{C}^+(\alpha, 0)$. It is Sturmian, and there is no $x \in [0,1)$ with $\mathtt{1}c = \mathcal{C}(\alpha', x)$ for any irrational $\alpha' \in (0,1)$.*

Proof. $\mathcal{C}^+(\alpha,0)_0 = 1$ because $x_0 = 0$, and $\mathcal{C}^+(\alpha,0)_{k+1} = \mathcal{C}^+(\alpha,\alpha)_k = \mathcal{C}(\alpha,\alpha)_k$ by Lemma 6 and Lemma 3, the orbit of α missing 0. So $\mathtt{1}c = \mathcal{C}^+(\alpha,0)$, and it is Sturmian by Corollary 11.

By Lemma 4 the first $N-1$ letters of c carry $\lfloor (N-1)\alpha + \alpha \rfloor = \lfloor N\alpha \rfloor$ ones, so the first N letters of $\mathtt{1}c$ carry $1 + \lfloor N\alpha \rfloor$.

Suppose $\mathtt{1}c = \mathcal{C}(\alpha', x)$. The slopes agree by Corollary 5, so $\alpha' = \alpha$, and Lemma 4 gives $\lfloor N\alpha + x \rfloor = 1 + \lfloor N\alpha \rfloor$ for every $N \geq 1$. Since $x \in [0,1)$ this forces $\{N\alpha\} + x \geq 1$ for every $N \geq 1$. By Lemma 7 there is N with $\{N\alpha\} < 1 - x$. □

The repair is to admit both codings.

Theorem 20 (Morse and Hedlund)**.** *A sequence is Sturmian if and only if it is $\mathcal{C}(\alpha, x)$ or $\mathcal{C}^+(\alpha, x)$ for some irrational $\alpha \in (0,1)$ and some $x \in [0,1)$.*

The forward implication is Theorem 10 and Corollary 11. **The converse is not proved here.** It is classical, and the route that fits this document is to run §4 and §5 backwards:

1. $F_\ell(1) = 2$, so both letters occur. $F_\ell(2) = 3$, so exactly one word of length 2 is absent, and it is $\mathtt{00}$ or $\mathtt{11}$: were $\mathtt{01}$ absent, every 1 would precede every 0 and $\ell = \mathtt{1}^p\mathtt{0}^\infty$, whose complexity is $p+1$ for all $m > p$. *Proved.*
2. The lengths of the maximal blocks of 0s take exactly two values, and they are consecutive, say $a-1$ and a. *Not proved here.* This is balancedness, and the classical proof is a minimal-counterexample induction producing a palindrome w with $\mathtt{0}w\mathtt{0}$ and $\mathtt{1}w\mathtt{1}$ both factors.
3. Given step 2, Lemma 15 applies verbatim: ℓ factors as $\mathtt{0}^{k_0}\tau_a(\ell')$ with ℓ' a sequence over $\{\mathtt{0},\mathtt{1}\}$. *Proved.*
4. ℓ' is again Sturmian, and both of its letters occur: a single block length would make ℓ eventually periodic, which caps complexity. *The second half is proved; that $F_{\ell'}(m) = m+1$ is not.*
5. Iterating produces $a_1, a_2, a_3, \ldots \geq 1$ and hence an irrational $\alpha = [\,0; a_1, a_2, \ldots]$, and the nested prefixes $\tau_{a_1}\tau_{a_2}\cdots\tau_{a_j}$ together with the shifts $k_0, k_1, \ldots$ pin a single x. *Not proved here.* This is the Ostrowski representation of x and needs a convergence argument.

7 The substitution

For $a_1 = 1$ the two blocks are $\tau_1(1) = \mathtt{10}$ and $\tau_1(0) = \mathtt{1}$, so τ_1 is the substitution σ of `sturmian.tex` §1:

$$\sigma(1) = \mathtt{10}, \qquad \sigma(0) = \mathtt{1}.$$

Theorem 21. *Let $\alpha = 1/\varphi$ and $c = \mathcal{C}(\alpha, \alpha)$. Then $c' = c$, $\sigma(c) = c$, and $c = \lim_j \sigma^j(\mathtt{1})$.*

Proof. $1/\alpha = \varphi = 1 + 1/\varphi$, so $a_1 = 1$ and $\beta = \alpha$. The first 1 of c is at $k_0 = 0$, where $x_0 = \alpha$ has coordinate $u_0 = (\alpha - (1 - \alpha))/\alpha = 2 - 1/\alpha = 1 - \alpha$. So $\{-u_0\} = \alpha$ and Theorem 16 gives $c' = \mathcal{C}^+(\alpha, \alpha) = \mathcal{C}(\alpha, \alpha) = c$, the middle equality by Lemma 3. Lemma 15 with $k_0 = 0$ then reads $c = \tau_1(c') = \sigma(c)$.

$\sigma(\mathtt{1}) = \mathtt{10}$ begins with `1`, and if $\sigma^j(\mathtt{1})$ is a prefix of $\sigma^{j+1}(\mathtt{1})$ then applying σ keeps it so; hence the limit exists. It is a prefix of every σ-fixed sequence beginning with `1`, and c is one. □

The converse computation of `sturmian.tex` §6 is now legitimate. Suppose ℓ is a coding whose blocks are `1` and `10`, so $a_1 = 1$, and suppose $\ell' = \ell$. By Corollary 5 the slopes agree, so $\beta = \alpha$, and

$$\frac{1}{\alpha} \;=\; 1 + \alpha, \qquad \alpha^2 + \alpha - 1 = 0, \qquad \alpha = \frac{\sqrt{5} - 1}{2}.$$

8 Square roots

Write $\alpha_0 = \alpha$ and $\alpha_{j+1} = \{1/\alpha_j\}$ for the tails of the continued fraction, and $a_{j+1} = \lfloor 1/\alpha_j \rfloor$ for its partial quotients. Let $\ell^{(j)}$ be the jth derived sequence of ℓ.

Theorem 22. *Let $\ell = \mathcal{C}(\alpha, 0)$. For every $j \geq 1$,*

$$\ell^{(j)} \;=\; \begin{cases} \mathcal{C}^+(\alpha_j,\, \alpha_j) & j \text{ odd} \\ \mathcal{C}(\alpha_j,\, \alpha_j) & j \text{ even.} \end{cases}$$

Proof. Let γ be irrational in $(0, 1)$, $a = \lfloor 1/\gamma \rfloor$ and $\gamma' = \{1/\gamma\}$.

In $\mathcal{C}(\gamma, 0)$ the first 1 is at $k_0 = a$: for $k \leq a$ we have $k\gamma \leq a\gamma < 1$ so $x_k = k\gamma$, and $k\gamma \geq 1 - \gamma$ holds first at $k = a$ because $(a + 1)\gamma > 1$. Its coordinate is

$$u_0 \;=\; \frac{a\gamma - (1 - \gamma)}{\gamma} \;=\; a + 1 - \frac{1}{\gamma} \;=\; 1 - \gamma',$$

so $\{-u_0\} = \gamma'$ and Theorem 16 gives $\mathcal{C}(\gamma, 0)' = \mathcal{C}^+(\gamma', \gamma')$. With $\gamma = \alpha$ this is the case $j = 1$.

In $\mathcal{C}(\gamma, \gamma)$ and in $\mathcal{C}^+(\gamma, \gamma)$ the first 1 is at $k_0 = a - 1$, by the same count applied to $x_k = (k+1)\gamma$; the value $x_{k_0} = a\gamma$ and the coordinate $u_0 = 1 - \gamma'$ are as above. So Theorem 16 gives

$$\mathcal{C}(\gamma, \gamma)' = \mathcal{C}^+(\gamma', \gamma'), \qquad \mathcal{C}^+(\gamma, \gamma)' = \mathcal{C}(\gamma', \gamma'),$$

the second because $1 - u_0 = \gamma'$. Induction on j. □

Corollary 23. *If the continued fraction of α is purely periodic with period L, then $\ell = \mathcal{C}(\alpha, 0)$ satisfies*

$$\ell^{(L)} \;=\; \mathcal{C}(\alpha, \alpha) \;=\; \ell_1 \ell_2 \ell_3 \cdots,$$

the sequence ℓ with its first letter removed.

Proof. $\alpha_L = \alpha$, so Theorem 22 gives $\mathcal{C}(\alpha, \alpha)$ or $\mathcal{C}^+(\alpha, \alpha)$, and those are equal by Lemma 3 because the orbit $\{(k + 1)\alpha\}$ never meets 0. Lemma 6 identifies $\mathcal{C}(\alpha, \alpha)$ with the shift of ℓ. □

Corollary 24. *Conversely, if $\ell^{(L)}$ is a shift of ℓ then $\alpha_L = \alpha$, so the continued fraction of α is purely periodic with period dividing L.*

Proof. Shifting does not change the density, so Corollary 5 gives $\alpha_L = \alpha$. □

Now take n a non-square, $a_0 = \lfloor \sqrt{n} \rfloor$ and $\alpha = \sqrt{n} - a_0$. If $\sqrt{n} = [\, a_0;\ \overline{a_1, \ldots, a_L}\,]$ then $\alpha = [\, 0;\ \overline{a_1, \ldots, a_L}\,]$, purely periodic, since the tails of α are the tails of $\sqrt{n}$ from index 1. Pure periodicity is Legendre's theorem, formalised in `lean/Palindrome.lean` as `purely_periodic`.

Corollary 25. *For every non-square n, the sequence $\ell = \mathcal{C}(\sqrt{n} - a_0,\, 0)$ returns to itself with one letter removed after exactly L derivations, where L is the period of the continued fraction of $\sqrt{n}$.*

`sturmian.tex` observes this for $n = 2$ and $n = 7$. For $n = 2$, $\alpha = \sqrt{2} - 1$, $a_1 = 2$, $\beta = \alpha$ and $L = 1$: one derivation returns the shift. For $n = 7$, $\alpha = \sqrt{7} - 2$ and the four tails are

$$\sqrt{7} - 2, \quad \frac{\sqrt{7} - 1}{3}, \quad \frac{\sqrt{7} - 1}{2}, \quad \frac{\sqrt{7} - 2}{3}, \quad \sqrt{7} - 2,$$

with partial quotients $1, 1, 1, 4$, which is the table of §5 there. Every row of that table is Sturmian by Corollary 17 and Theorem 10. The intercept of each row equals its own slope, by Theorem 22.

9 Lagrange

Theorem 26. *If the continued fraction of α is eventually periodic then α is a quadratic irrational.*

Proof. Let $\alpha_m = \alpha_{m+L}$. Each tail is a Möbius image of the next, $\alpha_j = 1/(a_{j+1} + \alpha_{j+1})$, so composing L of them gives integers p, q, r, s with

$$\alpha_m \;=\; \frac{p\,\alpha_{m+L} + q}{r\,\alpha_{m+L} + s} \;=\; \frac{p\,\alpha_m + q}{r\,\alpha_m + s},$$

hence $r\alpha_m^2 + (s - p)\alpha_m - q = 0$ with $(r, s - p, q) \neq 0$. So α_m is a quadratic irrational, and composing m further Möbius maps with integer coefficients keeps α in $\mathbb{Q}(\alpha_m)$. □

The converse — every quadratic irrational has an eventually periodic continued fraction — is **not proved here**. For $\alpha = \sqrt{n} - a_0$ it is `purely_periodic` in `lean/Palindrome.lean`, by way of reduced forms $(P + \sqrt{D})/Q$ and the finiteness of the reduced set. The general case needs the preperiod, and neither `sturmian.tex` nor the Lean development has it.

10 What is not proved here

1. The converse half of Theorem 20: steps 2, 4 and 5 of the list in §6. Step 2 is balancedness; step 5 is an Ostrowski convergence argument.
2. Lagrange's theorem for a general quadratic irrational (§9). The case $\sqrt{n} - \lfloor \sqrt{n} \rfloor$ is formalised.
3. Nothing else. Every other assertion of `sturmian.tex` is proved above, and the numbers in its tables — the slopes, the four derived sequences of $n = 7$, the gap lists for $n = 2$ and $n = 7$, the four factors of length 3 — have been checked in exact arithmetic in $\mathbb{Q}(\sqrt{n})$.