Ellipticity

1 Definition

Ellipticity is a property of differential operators that ensures certain regularity and solvability conditions for partial differential equations (PDEs). A linear differential operator is said to be elliptic if its principal symbol is invertible. Formally, consider a linear differential operator of the form:

$$L(u) = \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha u, \tag{1}$$

where D^α denotes the multi-index derivative operator, $a_\alpha(x)$ are smooth coefficient functions, and m is the order of the operator. The operator L is elliptic if the principal symbol $p(x, \xi)$ defined by:

$$p(x, \xi) = \sum_{|\alpha| = m} a_\alpha(x) \xi^\alpha \tag{2}$$

is non-vanishing for all (x, ξ) in the cotangent bundle, i.e., $p(x, \xi) \neq 0$ for all $\xi \neq 0$.

2 Examples

A classic example of an elliptic operator is the Laplace operator given by:

$$\Delta u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}. \tag{3}$$

The principal symbol of Δ is:

$$p(x, \xi) = \xi_1^2 + \xi_2^2, \tag{4}$$

which is clearly positive for all $(\xi_1, \xi_2) \neq (0, 0)$, confirming that Δ is elliptic.
Another example is the operator:

$$L(u) = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \lambda u, \tag{5}$$

where $\lambda > 0$. The principal symbol is:

$$p(x, \xi) = \xi_1^2 + \xi_2^2 + \lambda, \tag{6}$$

which is also non-vanishing for all (ξ_1, ξ_2), thus confirming ellipticity.

3 Key Results

One of the fundamental results regarding elliptic operators is the existence and uniqueness of solutions to boundary value problems. For instance, the following theorem holds:

[Schauder Estimates] Let L be a linear elliptic operator defined on a bounded domain $\Omega \subset \mathbb{R}^n$ with smooth boundary. If $f \in C^{0,\alpha}(\overline{\Omega})$ for some $0 < \alpha < 1$, then the Dirichlet problem for $Lu = f$ has a unique solution $u \in C^{2,\alpha}(\overline{\Omega})$ [**?**].

This theorem illustrates the regularity properties of solutions to elliptic equations, which are crucial in both theoretical and applied mathematics.

4 Diagrams / Visual Description

To visualize ellipticity, consider the principal symbol $p(x, \xi)$ as a surface in the (x, ξ) space. The condition for ellipticity can be represented as a non-vanishing condition on this surface, indicating that the operator behaves well in terms of invertibility. A diagram could show the region where $p(x, \xi) = 0$ and highlight that for elliptic operators, this region must not intersect with the non-zero ξ axis.

5 Connections

Ellipticity is closely related to several other concepts in the field of PDEs and differential geometry. For instance, it is connected to the notion of regularity theory, where elliptic operators yield smooth solutions under appropriate conditions. Additionally, ellipticity is a key condition for the Fredholm alternative, which states that for elliptic operators, the kernel and cokernel are finite-dimensional. Furthermore, elliptic operators are often contrasted with hyperbolic and parabolic operators, which exhibit different properties regarding the propagation of solutions.

6 References

- [**?**]
- [**?**]