

reference priors

Dr. Merlise Clyde

objective prior distributions

can you be Bayesian and not use “prior” information?

objective prior distributions

can you be Bayesian and not use “prior” information?

let prior sample size n_0 go to zero, s_0^2 go to zero and
 $v_0 = n_0 - 1$ go to -1

objective prior distributions

can you be Bayesian and not use “prior” information?

let prior sample size n_0 go to zero, s_0^2 go to zero and $v_0 = n_0 - 1$ go to -1

$$m_n = \frac{n\bar{Y} + n_0 m_0}{n + n_0} \rightarrow \bar{Y}$$

objective prior distributions

can you be Bayesian and not use “prior” information?

let prior sample size n_0 go to zero, s_0^2 go to zero and $v_0 = n_0 - 1$ go to -1

$$m_n = \frac{n\bar{Y} + n_0 m_0}{n + n_0} \rightarrow \bar{Y}$$

$$n_n = n_0 + n \rightarrow n$$

objective prior distributions

can you be Bayesian and not use “prior” information?

let prior sample size n_0 go to zero, s_0^2 go to zero and $v_0 = n_0 - 1$ go to -1

$$m_n = \frac{n\bar{Y} + n_0 m_0}{n + n_0} \rightarrow \bar{Y}$$

$$n_n = n_0 + n \rightarrow n$$

$$v_n = v_0 + n \rightarrow n - 1$$

objective prior distributions

can you be Bayesian and not use “prior” information?

let prior sample size n_0 go to zero, s_0^2 go to zero and $v_0 = n_0 - 1$ go to -1

$$m_n = \frac{n\bar{Y} + n_0 m_0}{n + n_0} \rightarrow \bar{Y}$$

$$n_n = n_0 + n \rightarrow n$$

$$v_n = v_0 + n \rightarrow n - 1$$

$$s_n^2 = \frac{1}{v_n} \left[s_0^2 v_0 + s^2 (n - 1) + \frac{n_0 n}{n_n} (\bar{Y} - m_0)^2 \right] \rightarrow s^2$$

reference priors

- ▶ limiting NormalGamma(0, 0, 0, -I) conjugate “prior” is

$$p(\mu \mid \sigma^2) \propto 1$$

reference priors

- ▶ limiting NormalGamma(0, 0, 0, -I) conjugate “prior” is

$$p(\mu \mid \sigma^2) \propto 1$$

$$p(\sigma^2) \propto 1/\sigma^2$$

reference priors

- ▶ limiting NormalGamma(0, 0, 0, -I) conjugate “prior” is

$$p(\mu \mid \sigma^2) \propto 1$$

$$p(\sigma^2) \propto 1/\sigma^2$$

$$p(\mu, \sigma^2) \propto 1/\sigma^2$$

reference priors

- ▶ limiting NormalGamma(0, 0, 0, -I) conjugate “prior” is

$$p(\mu \mid \sigma^2) \propto 1$$

$$p(\sigma^2) \propto 1/\sigma^2$$

$$p(\mu, \sigma^2) \propto 1/\sigma^2$$

- ▶ Non-Generative or “improper prior distribution”

reference priors

- ▶ limiting NormalGamma(0, 0, 0, -1) conjugate “prior” is

$$p(\mu \mid \sigma^2) \propto 1$$

$$p(\sigma^2) \propto 1/\sigma^2$$

$$p(\mu, \sigma^2) \propto 1/\sigma^2$$

- ▶ Non-Generative or “improper prior distribution”
- ▶ “Formal” posterior distribution, proper distribution if $n \geq 2$

$$\text{NormalGamma}(\bar{Y}, n, s^2, n - 1)$$

$$\mu \mid \sigma^2, \text{data} \sim \text{N}(\bar{Y}, \sigma^2/n)$$

$$1/\sigma^2 \mid \text{data} \sim \text{Gamma}((n - 1)/2, s^2(n - 1)/2)$$

reference analysis

- ▶ under the reference prior $p(\mu, \sigma^2) \propto 1/\sigma^2$

$$\frac{\mu - \bar{Y}}{\sqrt{s^2/n}} \mid \text{data} \sim t(n-1, 0, 1)$$

reference analysis

- ▶ under the reference prior $p(\mu, \sigma^2) \propto 1/\sigma^2$

$$\frac{\mu - \bar{Y}}{\sqrt{s^2/n}} \mid \text{data} \sim t(n - 1, 0, 1)$$

- ▶ prior to seeing the data

$$\frac{\mu - \bar{Y}}{\sqrt{s^2/n}} \mid \mu, \sigma^2 \sim t(n - 1, 0, 1)$$

reference analysis

- ▶ under the reference prior $p(\mu, \sigma^2) \propto 1/\sigma^2$

$$\frac{\mu - \bar{Y}}{\sqrt{s^2/n}} \mid \text{data} \sim t(n-1, 0, 1)$$

- ▶ prior to seeing the data

$$\frac{\mu - \bar{Y}}{\sqrt{s^2/n}} \mid \mu, \sigma^2 \sim t(n-1, 0, 1)$$

- ▶ intervals estimates $(\bar{Y} - t_{1-\alpha/2}s/\sqrt{n}, \bar{Y} + t_{1-\alpha/2}s/\sqrt{n})$

reference analysis

- ▶ under the reference prior $p(\mu, \sigma^2) \propto 1/\sigma^2$

$$\frac{\mu - \bar{Y}}{\sqrt{s^2/n}} \mid \text{data} \sim t(n-1, 0, 1)$$

- ▶ prior to seeing the data

$$\frac{\mu - \bar{Y}}{\sqrt{s^2/n}} \mid \mu, \sigma^2 \sim t(n-1, 0, 1)$$

- ▶ intervals estimates $(\bar{Y} - t_{1-\alpha/2}s/\sqrt{n}, \bar{Y} + t_{1-\alpha/2}s/\sqrt{n})$

- ▶ Bayesian allows probability statements after seeing the data

$$P(\bar{Y} - t_{1-\alpha/2}s/\sqrt{n} < \mu < \bar{Y} + t_{1-\alpha/2}s/\sqrt{n}) = 1 - \alpha$$


tap water example: reference analysis

R Code


```
phi = rgamma(10000, (n-1)/2, s2*(n-1)/2)
sigma = 1/sqrt(phi)
post_mu = rnorm(10000, mean=ybar, sd=sigma/(sqrt(n)))
pred_y = rnorm(10000, post_mu, sigma)
quantile(pred_y, c(.025, .975))

##          2.5%          97.5%
## 6.692877 104.225954
```

comparison of posterior densities



comparison of posterior densities



```
sum(pred_y > 80)/length(pred_y) #  $P(Y > 80 \mid data)$ 
```

```
## [1] 0.1534
```

summary

- ▶ introduced reference priors

summary

- ▶ introduced reference priors
- ▶ sensitivity analysis

summary

- ▶ introduced reference priors
- ▶ sensitivity analysis
- ▶ check assumptions of model and prior

summary

- ▶ introduced reference priors
- ▶ sensitivity analysis
- ▶ check assumptions of model and prior

next: mixtures of conjugate priors,
robustness and MCMC