

# comparing two independent means: what to report

Dr. Merlise Clyde

# birth weight and smoking

2004 survey of 999 births on North Carolina



weight gain during pregnancy and mother's age

2004 survey of 999 births on North Carolina

weight    birth weight of baby in pounds



# birth weight and smoking

2004 survey of 999 births on North Carolina

weight birth weight of baby in pounds

habit status of the mother as a 'nonsmoker' or a 'smoker'



<https://pixabay.com/en/cigarette-smoking-tobacco-nicotine-1638135/>



# birth weight and smoking

2004 survey of 999 births on North Carolina

weight birth weight of baby in pounds

habit status of the mother as a 'nonsmoker' or  
a 'smoker'

two questions



<https://pixabay.com/en/cigarette-smoking-tobacco-nicotine-1638135/>



# birth weight and smoking

2004 survey of 999 births on North Carolina

weight birth weight of baby in pounds

habit status of the mother as a 'nonsmoker' or a 'smoker'

two questions

1. is the average birth weight of babies whose mothers did not smoked different from the average weight gain of younger others?



<https://pixabay.com/en/cigarette-smoking-tobacco-nicotine-1638135/>



# birth weight and smoking

2004 survey of 999 births on North Carolina

**weight** birth weight of baby in pounds

**habit** status of the mother as a 'nonsmoker' or a 'smoker'



<https://pixabay.com/en/cigarette-smoking-tobacco-nicotine-1638135/>

two questions

1. is the average birth weight of babies whose mothers did not smoked different from the average weight gain of younger others?
2. if there are differences how large is the effect?



# model and priors

- ▶ random sample from two populations



# model and priors

- ▶ random sample from two populations

$$Y_{N,i} \stackrel{\text{iid}}{\sim} \mathbf{N}(\mu + \alpha/2, \sigma)$$



# model and priors

- ▶ random sample from two populations

$$Y_{N,i} \stackrel{\text{iid}}{\sim} \mathbf{N}(\mu + \alpha/2, \sigma)$$

$$Y_{S,i} \stackrel{\text{iid}}{\sim} \mathbf{N}(\mu - \alpha/2, \sigma)$$



# model and priors

- ▶ random sample from two populations

$$Y_{N,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu + \alpha/2, \sigma)$$

$$Y_{S,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu - \alpha/2, \sigma)$$

- ▶ difference in population means  $\mu + \alpha/2 - (\mu - \alpha/2) = \alpha$



# model and priors

- ▶ random sample from two populations

$$Y_{N,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu + \alpha/2, \sigma)$$

$$Y_{S,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu - \alpha/2, \sigma)$$

- ▶ difference in population means  $\mu + \alpha/2 - (\mu - \alpha/2) = \alpha$
- ▶  $H_1 : \alpha = 0$  versus  $H_2 : \alpha \neq 0$



# model and priors

- ▶ random sample from two populations

$$Y_{N,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu + \alpha/2, \sigma)$$

$$Y_{S,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu - \alpha/2, \sigma)$$

- ▶ difference in population means  $\mu + \alpha/2 - (\mu - \alpha/2) = \alpha$
- ▶  $H_1 : \alpha = 0$  versus  $H_2 : \alpha \neq 0$
- ▶ use the Jeffreys-Zellner-Siow prior



# model and priors

- ▶ random sample from two populations

$$Y_{N,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu + \alpha/2, \sigma)$$

$$Y_{S,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu - \alpha/2, \sigma)$$

- ▶ difference in population means  $\mu + \alpha/2 - (\mu - \alpha/2) = \alpha$
- ▶  $H_1 : \alpha = 0$  versus  $H_2 : \alpha \neq 0$
- ▶ use the Jeffreys-Zellner-Siow prior
- ▶ under  $H_2 : \delta = \alpha/\sigma$ , smaller effects  $r = .5$



# model and priors

- ▶ random sample from two populations

$$Y_{N,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu + \alpha/2, \sigma)$$

$$Y_{S,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu - \alpha/2, \sigma)$$

- ▶ difference in population means  $\mu + \alpha/2 - (\mu - \alpha/2) = \alpha$
- ▶  $H_1 : \alpha = 0$  versus  $H_2 : \alpha \neq 0$
- ▶ use the Jeffreys-Zellner-Siow prior
- ▶ under  $H_2 : \delta = \alpha/\sigma$ , smaller effects  $r = .5$
- ▶ under both  $H_1$  and  $H_2$  use reference priors



# model and priors

- ▶ random sample from two populations

$$Y_{N,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu + \alpha/2, \sigma)$$

$$Y_{S,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu - \alpha/2, \sigma)$$

- ▶ difference in population means  $\mu + \alpha/2 - (\mu - \alpha/2) = \alpha$
- ▶  $H_1 : \alpha = 0$  versus  $H_2 : \alpha \neq 0$
- ▶ use the Jeffreys-Zellner-Siow prior
- ▶ under  $H_2 : \delta = \alpha/\sigma$ , smaller effects  $r = .5$
- ▶ under both  $H_1$  and  $H_2$  use reference priors

$$p(\mu) = 1$$



# model and priors

- ▶ random sample from two populations

$$Y_{N,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu + \alpha/2, \sigma)$$

$$Y_{S,i} \stackrel{\text{iid}}{\sim} \text{N}(\mu - \alpha/2, \sigma)$$

- ▶ difference in population means  $\mu + \alpha/2 - (\mu - \alpha/2) = \alpha$
- ▶  $H_1 : \alpha = 0$  versus  $H_2 : \alpha \neq 0$
- ▶ use the Jeffreys-Zellner-Siow prior
- ▶ under  $H_2 : \delta = \alpha/\sigma$ , smaller effects  $r = .5$
- ▶ under both  $H_1$  and  $H_2$  use reference priors

$$p(\mu) = 1$$

$$p(\sigma^2) = 1/\sigma^2$$

## R code

```
library(statsr)
data(nc)
out =bayes_inference(y=weight, x=habit, data=nc,type='ht', null=0,
                     statistic='mean', alternative='twosided',
                     prior='JZS', r=.5, method='sim', show_summ=FALSE)

## Hypotheses:
## H1: mu_nonsmoker = mu_smoker
## H2: mu_nonsmoker != mu_smoker
##
## Priors: P(H1) = 0.5   P(H2) = 0.5
##
## Results:
## BF[H2:H1] = 1.4402
## P(H1|data) = 0.4098
## P(H2|data) = 0.5902
##
```




## summaries under $H_2$

### R code

```
out.ci = bayes_inference(y=weight, x=habit, data=nc, type='ci',
                          statistic='mean', prior='JZS', mu_0=0,
                          r=.5, method='sim', verbose=FALSE)
print(out.ci$summary, digits=2)
```


##	2.5%	25%	50%	75%	97.5%
## overall mean	6.856	6.95	7.0	7.04	7.1e+00
## mu_nonsmoker - mu_smoker	0.023	0.20	0.3	0.39	5.7e-01
## sigma^2	2.072	2.19	2.3	2.33	2.5e+00
## effect size	0.015	0.14	0.2	0.26	3.8e-01
## n_0	184.732	2016.09	4729.1	9594.28	2.6e+04

estimates of effect under  $H_2$





## estimates of effect under $H_2$



under  $H_2$ , 95% chance the average birth weight of babies born to nonsmokers is 0.02 to 0.57 pounds higher than that of babies born to smokers

# overall coverage

- ▶ conditional probability  $P(0.02 < \alpha < 0.57 \mid \text{data}, H_2) = 0.95$



# overall coverage

- ▶ conditional probability  $P(0.02 < \alpha < 0.57 \mid \text{data}, H_2) = 0.95$
- ▶  $P(H_2 \mid \text{data}) = 0.59$

# overall coverage

- ▶ conditional probability  $P(0.02 < \alpha < 0.57 \mid \text{data}, H_2) = 0.95$
- ▶  $P(H_2 \mid \text{data}) = 0.59$
- ▶ law of total probability  
 $P(0.02 < \alpha < 0.57 \mid \text{data})$



# overall coverage

- ▶ conditional probability  $P(0.02 < \alpha < 0.57 \mid \text{data}, H_2) = 0.95$
- ▶  $P(H_2 \mid \text{data}) = 0.59$
- ▶ law of total probability
$$P(0.02 < \alpha < 0.57 \mid \text{data})$$
$$= P(0.02 < \alpha < 0.57 \mid \text{data}, H_1)P(H_1 \mid \text{data}) +$$

# overall coverage

- ▶ conditional probability  $P(0.02 < \alpha < 0.57 \mid \text{data}, H_2) = 0.95$
- ▶  $P(H_2 \mid \text{data}) = 0.59$
- ▶ law of total probability

$$\begin{aligned} & P(0.02 < \alpha < 0.57 \mid \text{data}) \\ &= P(0.02 < \alpha < 0.57 \mid \text{data}, H_1)P(H_1 \mid \text{data}) + \\ & \quad P(0.02 < \alpha < 0.57 \mid \text{data}, H_2)P(H_2 \mid \text{data}) \end{aligned}$$



# overall coverage

- ▶ conditional probability  $P(0.02 < \alpha < 0.57 \mid \text{data}, H_2) = 0.95$
- ▶  $P(H_2 \mid \text{data}) = 0.59$
- ▶ law of total probability

$$\begin{aligned} & P(0.02 < \alpha < 0.57 \mid \text{data}) \\ &= P(0.02 < \alpha < 0.57 \mid \text{data}, H_1)P(H_1 \mid \text{data}) + \\ & \quad P(0.02 < \alpha < 0.57 \mid \text{data}, H_2)P(H_2 \mid \text{data}) \\ &= I(0 \text{ in CI})P(H_1 \mid \text{data}) + 0.95 \times P(H_2 \mid \text{data}) \end{aligned}$$

# overall coverage

- ▶ conditional probability  $P(0.02 < \alpha < 0.57 \mid \text{data}, H_2) = 0.95$
- ▶  $P(H_2 \mid \text{data}) = 0.59$
- ▶ law of total probability

$$\begin{aligned} & P(0.02 < \alpha < 0.57 \mid \text{data}) \\ &= P(0.02 < \alpha < 0.57 \mid \text{data}, H_1)P(H_1 \mid \text{data}) + \\ & \quad P(0.02 < \alpha < 0.57 \mid \text{data}, H_2)P(H_2 \mid \text{data}) \\ &= I(0 \text{ in CI})P(H_1 \mid \text{data}) + 0.95 \times P(H_2 \mid \text{data}) \\ &= 0 \times 0.41 + 0.95 \times 0.59 \end{aligned}$$



# overall coverage

- ▶ conditional probability  $P(0.02 < \alpha < 0.57 \mid \text{data}, H_2) = 0.95$
- ▶  $P(H_2 \mid \text{data}) = 0.59$
- ▶ law of total probability

$$\begin{aligned} & P(0.02 < \alpha < 0.57 \mid \text{data}) \\ &= P(0.02 < \alpha < 0.57 \mid \text{data}, H_1)P(H_1 \mid \text{data}) + \\ & \quad P(0.02 < \alpha < 0.57 \mid \text{data}, H_2)P(H_2 \mid \text{data}) \\ &= I(0 \text{ in CI})P(H_1 \mid \text{data}) + 0.95 \times P(H_2 \mid \text{data}) \\ &= 0 \times 0.41 + 0.95 \times 0.59 \\ &= 0.5605 \end{aligned}$$

# overall coverage

- ▶ conditional probability  $P(0.02 < \alpha < 0.57 \mid \text{data}, H_2) = 0.95$
- ▶  $P(H_2 \mid \text{data}) = 0.59$
- ▶ law of total probability

$$\begin{aligned} & P(0.02 < \alpha < 0.57 \mid \text{data}) \\ &= P(0.02 < \alpha < 0.57 \mid \text{data}, H_1)P(H_1 \mid \text{data}) + \\ & \quad P(0.02 < \alpha < 0.57 \mid \text{data}, H_2)P(H_2 \mid \text{data}) \\ &= I(0 \text{ in CI})P(H_1 \mid \text{data}) + 0.95 \times P(H_2 \mid \text{data}) \\ &= 0 \times 0.41 + 0.95 \times 0.59 \\ &= 0.5605 \end{aligned}$$

the average birth weight of babies born to nonsmokers is 0.02 to 0.57 pounds higher than that of babies born to smokers with probability 0.56



# summary

- ▶ compared two independent means using Bayesian hypothesis testing

# summary

- ▶ compared two independent means using Bayesian hypothesis testing
- ▶ quantify size of effect



# summary

- ▶ compared two independent means using Bayesian hypothesis testing
- ▶ quantify size of effect
- ▶ credible intervals conditional on  $H_2$

# summary

- ▶ compared two independent means using Bayesian hypothesis testing
- ▶ quantify size of effect
- ▶ credible intervals conditional on  $H_2$
- ▶ adjusting credible level to account for uncertainty of hypotheses



# summary

- ▶ compared two independent means using Bayesian hypothesis testing
- ▶ quantify size of effect
- ▶ credible intervals conditional on  $H_2$
- ▶ adjusting credible level to account for uncertainty of hypotheses

next: regression models to adjust for continuous explanatory variables