

comparing two paired means using Bayes factors

Dr. Merlise Clyde

zinc in drinking water



location	Surface	Bottom	Difference
1	0.43	0.415	0.015
2	0.266	0.238	0.028
3	0.567	0.39	0.177
4	0.531	0.41	0.121
5	0.707	0.605	0.102
⋮	⋮	⋮	⋮
10	0.723	0.612	0.111

model and hypotheses

- ▶ random sample of $n = 10$ differences Y_1, \dots, Y_n

model and hypotheses

- ▶ random sample of $n = 10$ differences Y_1, \dots, Y_n
- ▶ normal population with mean $\mu \equiv \mu_B - \mu_S$

model and hypotheses

- ▶ random sample of $n = 10$ differences Y_1, \dots, Y_n
- ▶ normal population with mean $\mu \equiv \mu_B - \mu_S$
- ▶ variance σ^2 unknown

model and hypotheses

- ▶ random sample of $n = 10$ differences Y_1, \dots, Y_n
- ▶ normal population with mean $\mu \equiv \mu_B - \mu_S$
- ▶ variance σ^2 unknown

no differences $H_1 : \mu_B = \mu_S \Leftrightarrow \mu = 0$

model and hypotheses

- ▶ random sample of $n = 10$ differences Y_1, \dots, Y_n
- ▶ normal population with mean $\mu \equiv \mu_B - \mu_S$
- ▶ variance σ^2 unknown

no differences $H_1 : \mu_B = \mu_S \Leftrightarrow \mu = 0$

means are different $H_2 : \mu_B \neq \mu_S \Leftrightarrow \mu \neq 0$

model and hypotheses

- ▶ random sample of $n = 10$ differences Y_1, \dots, Y_n
- ▶ normal population with mean $\mu \equiv \mu_B - \mu_S$
- ▶ variance σ^2 unknown

no differences $H_1 : \mu_B = \mu_S \Leftrightarrow \mu = 0$

means are different $H_2 : \mu_B \neq \mu_S \Leftrightarrow \mu \neq 0$

Bayes factor: $BF[H_1 : H_2] = \frac{p(\text{data}|H_1)}{p(\text{data}|H_2)}$

Bayes factor

Bayes factor:

$$BF[H_1 : H_2] = \frac{\int p(\text{data}|\mu=0,\sigma^2)p(\sigma^2|H_1) d\sigma^2}{\iint p(\text{data}|\mu,\sigma^2)p(\mu|\sigma^2,H_2)p(\sigma^2|H_2) d\mu d\sigma^2}$$

Bayes factor

Bayes factor:

$$BF[H_1 : H_2] = \frac{\int p(\text{data}|\mu=0,\sigma^2)p(\sigma^2|H_1) d\sigma^2}{\iint p(\text{data}|\mu,\sigma^2)p(\mu|\sigma^2,H_2)p(\sigma^2|H_2) d\mu d\sigma^2}$$

priors:

Bayes factor

Bayes factor:

$$BF[H_1 : H_2] = \frac{\int p(\text{data}|\mu=0,\sigma^2)p(\sigma^2|H_1) d\sigma^2}{\iint p(\text{data}|\mu,\sigma^2)p(\mu|\sigma^2,H_2)p(\sigma^2|H_2) d\mu d\sigma^2}$$

priors:

- ▶ $\mu \mid \sigma^2, H_2 \sim \text{N}(0, \sigma^2/n_0)$

Bayes factor

Bayes factor:

$$BF[H_1 : H_2] = \frac{\int p(\text{data}|\mu=0,\sigma^2)p(\sigma^2|H_1) d\sigma^2}{\iint p(\text{data}|\mu,\sigma^2)p(\mu|\sigma^2,H_2)p(\sigma^2|H_2) d\mu d\sigma^2}$$

priors:

- ▶ $\mu \mid \sigma^2, H_2 \sim \text{N}(0, \sigma^2/n_0)$
- ▶ $p(\sigma^2) \propto 1/\sigma^2$ for both H_1 and H_2

Bayes factor with reference prior σ^2

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

Bayes factor with reference prior σ^2

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

► t-statistic

$$t = \frac{|\bar{Y}|}{s/\sqrt{n}}$$

Bayes factor with reference prior σ^2

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

- ▶ t-statistic $t = \frac{|\bar{Y}|}{s/\sqrt{n}}$
- ▶ sample standard deviation s

Bayes factor with reference prior σ^2

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

- ▶ t-statistic $t = \frac{|\bar{Y}|}{s/\sqrt{n}}$
- ▶ sample standard deviation s
- ▶ degrees of freedom $\nu = n - 1$

Bayes factor with reference prior σ^2

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

- ▶ t-statistic $t = \frac{|\bar{Y}|}{s/\sqrt{n}}$
 - ▶ sample standard deviation s
 - ▶ degrees of freedom $\nu = n - 1$
- $n_0 \rightarrow 0, BF[H_1 : H_2] \rightarrow \infty$

Bayes factor with reference prior σ^2

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

- ▶ t-statistic $t = \frac{|\bar{Y}|}{s/\sqrt{n}}$
- ▶ sample standard deviation s
- ▶ degrees of freedom $\nu = n - 1$
 $n_0 \rightarrow 0, BF[H_1 : H_2] \rightarrow \infty$
- ▶ no improper or vague prior on μ

information paradox

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

information paradox

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

- fixed n and n_0

information paradox

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

- ▶ fixed n and n_0
- ▶ t-statistic $|t| \rightarrow \infty$

information paradox

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

- ▶ fixed n and n_0
- ▶ t-statistic $|t| \rightarrow \infty$
- ▶ bounded Bayes factor

$$BF[H_1 : H_2] \rightarrow \left(\frac{n_0}{n_0+n} \right)^{\frac{n-1}{2}}$$

information paradox

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

- ▶ fixed n and n_0
- ▶ t-statistic $|t| \rightarrow \infty$
- ▶ bounded Bayes factor

$$BF[H_1 : H_2] \rightarrow \left(\frac{n_0}{n_0+n} \right)^{\frac{n-1}{2}}$$

- ▶ Jeffrey's recommended a Cauchy prior $C(0, r^2 \sigma^2)$

information paradox

$$BF[H_1 : H_2] = \left(\frac{n+n_0}{n_0} \right)^{1/2} \left(\frac{t^2 \frac{n_0}{n+n_0} + \nu}{t^2 + \nu} \right)^{\frac{\nu+1}{2}}$$

- ▶ fixed n and n_0
- ▶ t-statistic $|t| \rightarrow \infty$
- ▶ bounded Bayes factor

$$BF[H_1 : H_2] \rightarrow \left(\frac{n_0}{n_0+n} \right)^{\frac{n-1}{2}}$$

- ▶ Jeffrey's recommended a Cauchy prior $C(0, r^2 \sigma^2)$
- ▶ Jeffrey-Zellner-Siow or 'JZS' prior

zinc concentration example

```
bayes_inference(difference, data=zinc, statistic="mean", type="ht",
                prior="JZS", mu_0=0, method="theo", alt="twosided")

## Single numerical variable
## n = 10, y-bar = 0.0804, s = 0.0523
## (Using Zellner-Siow Cauchy prior:  $\mu \sim C(0, 1 \cdot \sigma)$ )
## (Using Jeffreys prior:  $p(\sigma^2) = 1/\sigma^2$ )
##
## Hypotheses:
## H1:  $\mu = 0$  versus H2:  $\mu \neq 0$ 
## Priors:
##  $P(H1) = 0.5$  ,  $P(H2) = 0.5$ 
## Results:
##  $BF[H2:H1] = 50.7757$ 
##  $P(H1|data) = 0.0193$     $P(H2|data) = 0.9807$ 
##
```

summary

- ▶ Cauchy prior for μ and hypothesis testing

summary

- ▶ Cauchy prior for μ and hypothesis testing
- ▶ requires numerical integration

summary

- ▶ Cauchy prior for μ and hypothesis testing
- ▶ requires numerical integration
- ▶ modify scale $r < 1$ if expect smaller effect

summary

- ▶ Cauchy prior for μ and hypothesis testing
- ▶ requires numerical integration
- ▶ modify scale $r < 1$ if expect smaller effect
- ▶ use `bayes_inference` for credible intervals

summary

- ▶ Cauchy prior for μ and hypothesis testing
- ▶ requires numerical integration
- ▶ modify scale $r < 1$ if expect smaller effect
- ▶ use `bayes_inference` for credible intervals

next: comparing two means