

1. Depositional Environment Categories: Do you have a comprehensive list of depositional environment categories? If not, have you considered extracting them from the knowledge base in a two-stage process? (Extracting an exhaustive list first, and the context for each in second stage)

This needs to be done in both approaches.

- 1) Considering extracting it from knowledge base (two stage process).
- 2) Consider the comprehensive list of depositional environment given below having a gross environment (eg: fluvial), and sub environment (eg: meandering).

Continental (Terrestrial) Environments

Fluvial (Rivers)

- Straight channels: Minimal sinuosity, high-energy, coarse sediment.
- Meandering channels: High sinuosity, point bars, oxbow lakes, fine sediment.
- Braided channels: Multiple shifting channels, bars, high sediment load.
- Anabranching/anastomosing: Multiple semi-stable channels separated by vegetated islands.

Alluvial Fans

- Proximal fan: Closest to source, coarse gravel, debris flows.
- Mid-fan: Mix of sand and gravel, transitional deposits.
- Distal fan: Finer sediments, often reworked by water/wind, higher sorting.

Aeolian (Desert Dunes)

- Dune fields: Active or stabilized dunes, various shapes (barchan, transverse, parabolic).
- Interdune areas: Flat zones between dunes, may contain ephemeral lakes.
- Sand sheets/loess: Extensive, flat deposits of wind-blown silt or sand.

Lacustrine (Lakes)

- Lake margins: Sandy to muddy, influenced by wave activity.
- Deep lake/central basin: Fine-grained sediments, low energy.
- Deltaic/fan deltas: Where rivers or fans enter the lake, may form Gilbert-type deltas.

Transitional Environments

Deltaic

- River-dominated: Channel mouth bars, distributary channels, levees.
- Wave-dominated: Smooth shorelines, beach ridges, low distributary numbers.

- Tide-dominated: Sand ridges, tidal flats, dendritic channels.

Estuarine & Tidal Flats

- Channels: Subtidal, tidal currents, mixed energy.
- Intertidal flats: Mud/sand flats, exposed at low tide, high organic activity.
- Supratidal marshes: Above high tide, vegetated, organic-rich.

Barrier Island & Lagoonal Systems

- Barrier core: Shoreface, foreshore, backshore.
- Tidal inlets: Openings allowing tidal exchange.
- Back-barrier/lagoon: Protected, low energy, fine sediments and marshes.

Marine Environments

Shallow Marine (Shelf)

- Inner shelf: Proximal, wave-action, sand and shell beds.
- Middle shelf: Mixed sediment, less energy, faunal diversity.
- Outer shelf: Muddy, low energy, bioturbation.

Carbonate Platforms & Reefs

- Fore-reef: High-energy slope, broken coral debris.
- Reef crest: Main reef, massive coral growth.
- Back-reef/lagoon: Protected, fine sediments, sometimes stromatolites.

Deep Marine (Turbidites & Submarine Fans)

- Submarine channel/levee: Coarse deposits from turbidity flows.
- Lobe/fan complexes: Finer turbidite sequences, graded bedding.
- Basin plain: Pelagic muds, low-energy fine sediments.

2. Expert Geologist's Approach: What is the standard operating procedure for expert geologists when identifying depositional environments manually? Is there a step-by-step protocol?





Figure 1

As shown in Figure 1, geologists may use multiple data types to identify depositional environments, including core data, well logs, biostratigraphic data, sedimentological data, and seismic data. Each data source provides different insights and together help build a comprehensive understanding of the depositional setting.

In our project, the initial focus will be on extracting information solely from well data. Well data, along with well tops and a suitable supporting database, will be used to identify both the gross depositional environment and its subcategories. This approach leverages well logs and associated stratigraphic markers to interpret sedimentary facies and depositional processes effectively.

3. Log File Plots: What types of plots can be generated from a log file (.las), and can you enumerate them?



All properties provided in LAS files can be plotted against depth or measured depth (MD), with the property values on the X-axis and depth increasing downward on the Y-axis (an inverted depth scale).

Among these, the gamma ray (GR) log is often the most relevant for identifying depositional environments because it effectively distinguishes between shale-rich and clean sand/shale intervals, helping to characterize lithology and sedimentary facies.




Figure 2




Figure 3

Figure 2 shows sample plots from the shared well logs, which display certain patterns. Similar patterns can also be observed in the knowledge base plots, as illustrated in Figure 3.

4. Knowledge Base Comparison: Are the graphs in the knowledge base similar in characteristics (axes, scales, units) to the plots generated from log files, allowing for direct comparison?

Direct comparison is feasible when the Y-axis depth range is matched and inverted uniformly in both the knowledge base graphs and the log file plots.

5. Inferencing Points: If direct comparison isn't possible, what inferencing points do theoretical geologists commonly use from literature to understand log file plots?

If visual comparison through plotting is challenging or ineffective, an alternative approach is to analyze and compare the log feature values numerically, which may provide quantitative insight even without graphical matching.

6. Data Size: What is the size of the knowledge base and log file collection?

Sample files are provided.

7. Golden Dataset Feasibility: Is it feasible to curate a golden dataset as mentioned in the proposal, and what would be the requirements?

For the given set of well logs the gross depositional environment is Fluvial.