


Bayesian Network

Graphical Models

- Key Idea:
  - Conditional independence assumptions useful
  - but Naïve Bayes is extreme!
  - Graphical models express sets of conditional independence assumptions via graph structure
  - Graph structure plus associated parameters define *joint probability distribution over set of variables*

Graphical Models – Why Care?

- Among most important ML developments of the decade
- Graphical models allow combining:
  - Prior knowledge in form of dependencies/independencies
  - Prior knowledge in form of priors over parameters
  - Observed training data
- Principled and ~general methods for
  - Probabilistic inference
  - Learning
- Useful in practice
  - Diagnosis, help systems, text analysis, time series models, ...

Conditional Independence

Definition: X is conditionally independent of Y given Z, if the probability distribution governing X is independent of the value of Y, given the value of Z

$$(\forall i, j, k) P(X = x_i | Y = y_j, Z = z_k) = P(X = x_i | Z = z_k)$$

Which we often write $P(X|Y,Z) = P(X|Z)$

E.g., $P(Thunder|Rain, Lightning) = P(Thunder|Lightning)$

Marginal Independence

Definition: X is marginally independent of Y if

$$(\forall i, j) P(X = x_i, Y = y_j) = P(X = x_i) P(Y = y_j)$$


Equivalently, if

$$(\forall i, j) P(X = x_i | Y = y_j) = P(X = x_i)$$

Equivalently, if

$$(\forall i, j) P(Y = y_i | X = x_j) = P(Y = y_i)$$

Bayes Nets define Joint Probability Distribution in terms of this graph, plus parameters





$$P(X_1, X_2, X_3, X_4, X_5, X_6, X_7, X_8)$$
$$= P(X_1)\, P(X_2)\, P(X_3|\, X_1)\, P(X_4|\, X_2)\, P(X_5|\, X_2)$$
$$P(X_6|\, X_3, X_4)\, P(X_7|\, X_6)\, P(X_8|\, X_5, X_6)$$

Benefits of Bayes Nets:

- Represent the full joint distribution in fewer parameters, using prior knowledge about dependencies
- Algorithms for inference and learning

Bayesian Networks Definition





A Bayes network represents the joint probability distribution over a collection of random variables


A Bayes network is a directed acyclic graph and a set of conditional probability distributions (CPD's)

- Each node denotes a random variable
- Edges denote dependencies
- For each node X_i its CPD defines $P(X_i \mid Pa(X_i))$
- The joint distribution over all variables is defined to be

$$P(X_1 \ldots X_n) = \prod_i P(X_i | Pa(X_i))$$

Pa(X) = immediate parents of X in the graph

Bayesian Network



Nodes = random variables

A conditional probability distribution (CPD) is associated with each node N, defining P(N | Parents(N))

Parents	P(W|Pa)	P(¬W|Pa)
L, R	0	1.0
L, ¬R	0	1.0
¬L, R	0.2	0.8
¬L, ¬R	0.9	0.1

WindSurf

The joint distribution over all variables:

$$P(X_1 \ldots X_n) = \prod_i P(X_i|Pa(X_i))$$


Some helpful terminology

Parents = Pa(X) = immediate parents

Antecedents = parents, parents of parents, ...

Children = immediate children


Descendents = children, children of children, ...



Parents	P(W|Pa)	P(¬W|Pa)
L, R	0	1.0
L, ¬R	0	1.0
¬L, R	0.2	0.8
¬L, ¬R	0.9	0.1

W

Bayesian Network



What can we say about conditional independencies in a Bayes Net?

One thing is this:


Each node is conditionally independent of its non-descendents, given only its immediate parents.

Parents	P(W|Pa)	P(¬W|Pa)
L, R	0	1.0
L, ¬R	0	1.0
¬L, R	0.2	0.8
¬L, ¬R	0.9	0.1

WindSurf

Bayesian Networks

- CPD for each node X_i describes $P(X_i \mid Pa(X_i))$



Parents	P(W|Pa)	P(¬W|Pa)
L, R	0	1.0
L, ¬R	0	1.0
¬L, R	0.2	0.8
¬L, ¬R	0.9	0.1

W

Chain rule of probability says that in general:

$$P(S,L,R,T,W) = P(S)P(L|S)P(R|S,L)P(T|S,L,R)P(W|S,L,R,T)$$

8 params

But in a Bayes net: $P(X_1 \ldots X_n) = \prod_i P(X_i|Pa(X_i))$

$P(S,L,R,T,W) = P(S)\,P(L|S)\,P(R|S)\,P(T|L)\,P(W|L\,R)$

2

Bayesian networks

a BN consists of a Directed Acyclic Graph (DAG) and a set of conditional probability distributions

in the DAG

- each node denotes random a variable
- each edge from X to Y represents that X *directly influences* Y
- formally: each variable X is independent of its non-descendants given its parents

each node X has a *conditional probability distribution* (CPD) representing $P(X \mid Parents(X))$

Bayesian networks

sing the chain rule, a joint probability distribution can be xpressed as

$$P(X_1, \ldots, X_n) = \mathrm{P}(X_1)\prod_{i=2}^{n} P(X_i \mid X_1, \ldots, X_{i-1}))$$

BN provides a compact representation of a joint obability distribution

$$P(X_1, \ldots, X_n) = \prod_{i=1}^{n} P(X_i \mid Parents(X_i))$$

Bayesian network example

- Consider the following 5 binary random variables:

  B = a burglary occurs at your house

  E = an earthquake occurs at your house


  A = the alarm goes off

  J = John calls to report the alarm


  M = Mary calls to report the alarm

- Suppose we want to answer queries like what is

Bayesian network example



Bayesian networks




$$P(B,E,A,J,M) = P(B) \times P(E) \times P(A \mid B,E) \times P(J \mid A) \times P(M \mid A)$$

- a standard representation of the joint distribution for the Alarm example has $2^5 = 32$ parameters
- the BN representation of this distribution has 20 parameters

Bayesian networks

- consider a case with 10 binary random variables

- How many parameters does a BN with the following graph structure have?



= 42

- How many parameters does the standard table representation of the joint distribution have? = 1024

Advantages of the Bayesian network representation

- Captures independence and conditional independence where they exist
- Encodes the relevant portion of the full joint among variables where dependencies exist
- Uses a graphical representation which lends insight int the complexity of inference

Bayesian Belief Networks

- Let we *l* random variables
- The joint probability is given by,

$$p(x_1, x_2, \ldots, x_\ell) = p(x_\ell \mid x_{\ell-1}, \ldots, x_1) \cdot p(x_{\ell-1} \mid x_{\ell-2}, \ldots, x_1) \cdot \ldots$$
$$\ldots \cdot p(x_2 \mid x_1) \cdot p(x_1)$$

Bayesian Belief Networks

The formula

$$p(x_1, x_2, \ldots, x_\ell) = p(x_\ell \mid x_{\ell-1}, \ldots, x_1) \cdot p(x_{\ell-1} \mid x_{\ell-2}, \ldots, x_1) \cdot \ldots$$
$$\ldots \cdot p(x_2 \mid x_1) \cdot p(x_1)$$

can be written as

$$p(x_1, x_2, \ldots, x_\ell) = p(x_1) \cdot \prod_{i=2}^{\ell} p(x_i \mid A_i)$$

where

$$A_i \subseteq \{x_{i-1}, x_{i-2}, \ldots, x_1\}$$

– For example, if ℓ=6, then we could assume:

$$p(x_6 \mid x_5,\ldots,x_1) = p(x_6 \mid x_5, x_4)$$

Then:

$$A_6 = \{x_5, x_4\} \subseteq \{x_5,\ldots,x_1\}$$

– Simialrly, if we assume

$$p(x_5|x_4,\ldots,x_1) = p(x_5|x_4)$$

$$p(x_4|x_3,x_2,x_1) = p(x_4|x_2,x_1)$$

$$p(x_3|x_2,x_1) = p(x_3|x_2)$$

$$p(x_2|x_1) = p(x_2)$$

Then:

$$A_5 = \{x_4\},\ A_4 = \{x_2,x_1\},\ A_3 = \{x_2\}, A_2 = \emptyset$$

– Simialrly, if we assume

$$p(x_5|x_4,\ldots,x_1) = p(x_5|x_4)$$

$$p(x_4|x_3,x_2,x_1) = p(x_4|x_2,x_1)$$

$$p(x_3|x_2,x_1) = p(x_3|x_2)$$

$$p(x_2|x_1) = p(x_2)$$


Then:

$$A_5 = \{x_4\},\ A_4 = \{x_2,x_1\},\ A_3 = \{x_2\}, A_2 = \emptyset$$


– The above is a generalization of the Naïve – Bayes. For the Naïve – Bayes the assumption is:

$$A_i = \emptyset, \text{ for } i=1, 2, \ldots, \ell$$

– A graphical way to portray conditional dependencies




– A graphical way to portray conditional dependencies



- According to this figure we have that:
  - x_6 is conditionally dependent on x_4, x_5.
  - x_5 on x_4
  - x_4 on x_1, x_2
  - x_3 on x_2
  - x_1, x_2 are conditionally independent on other variables.

– A graphical way to portray conditional dependencies



- According to this figure we have that:
  - x_6 is conditionally dependent on x_4, x_5.
  - x_5 on x_4
  - x_4 on x_1, x_2
  - x_3 on x_2
  - x_1, x_2 are conditionally independent on other variables.


- For this case:

$$p(x_1, x_2, \ldots, x_6) = p(x_6 \mid x_5, x_4) \cdot p(x_5 \mid x_4) \cdot p(x_3 \mid x_2) \cdot p(x_2) \cdot p(x_1)$$

- Bayesian Networks
  - a directed acyclic graph (DAG)
  - the nodes correspond to random variables
  - arc represents parent-child (*dependence*) relationship

- A Bayesian Network is specified by:
  - The prior probabilities of its root nodes.
  - The conditional probabilities of the non-root nodes, given their parents, for ALL possible combinations.

– A Bayesian Network from a medical application



- BBN models conditional dependencies concerning smokers (S), tendencies to develop cancer (C) and heart disease (H), together with variables corresponding to heart (H1, H2) and cancer (C1, C2) medical tests.


- any joint probability can be obtained by multiplying the prior (root nodes) and the conditional (non-root nodes) probabilities.

- Training: given a topology, probabilities are estimated from training data. There are also methods that learn the topology.

- Probability Inference: Given an pattern (evidence), the goal is to compute the conditional probabilities for some of the other variables (class)


- Example: Consider the Bayesian network of the figure:

$$P(x1) = 0.60 \quad P(y1|x1) = 0.40 \quad P(z1|y1) = 0.25 \quad P(w1|z1) = 0.45$$
$$P(y1|x0) = 0.30 \quad P(z1|y0) = 0.60 \quad P(w1|z0) = 0.30$$



- Random variables: *x*, *y*, *w*, *z*
- *x*0 means *x* = 0
- *x*1 means *x* = 1

- We can calculate the other probabilities




Example: $p(y_1)$:


$$P(y1) = \sum_x P(y1, x) = P(y1, x1) + P(y1, x0)$$

$$P(y1) = P(y1|x1)P(x1) + P(y1|x0)P(x0) = (0.4)(0.6) + (0.3)(0.4) = 0.36$$

- We can calculate the other probabilities




- Given this info, we can answer any probabilistic query:



a) If x is measured to be $x=1$ ($x1$), compute $P(z1|x1)$ *and* $P(w0|x1)$.

b) If w is measured to be $w=1$ ($w1$) compute $P(z1|w1)$].

a) If x is measured to be $x=1$ ($x1$), compute P(z1|x1) and $P(w0|x1)$.




$$P(z1|x1) = P(z1|y1, x1)P(y1|x1) + P(z1|y0, x1)P(y0|x1)$$

$$= P(z1|y1)P(y1|x1) + P(z1|y0)P(y0|x1)$$

$$= (0.25)(0.4) + (0.6)(0.6) = 0.46$$

a) If x is measured to be $x=1$ ($x1$), compute P(z1|x1) and $P(w0|x1)$.




$$P(w0|x1) = P(w0|z1, x1)P(z1|x1) + P(w0|z0, x1)P(z0|x1)$$

$$= P(w0|z1)P(z1|x1) + P(w0|z0)P(z0|x1)$$


$$= (0.55)(0.46) + (0.7)(0.54) = 0.63$$

b) If w is measured to be w=1 ($w1$) compute P(z1|w1|)].




$$P(z1|w1) = \frac{P(w1|z1)P(z1)}{P(w1)} = \frac{(0.45)(0.47)}{0.37} = 0.57$$


What's about more complex networks?



What's about more complex networks?




We will study this graph




We can show:

- $P(D|E)=P(D)$
- $P(Hb|HD, E, D)= P(Hb|D)$
- $P(CP|Hb, HD, E, D)= P(CP|Hb, HD)$
- $P(BP|CP, Hb, HD, E, D)= P(BP|HD)$
- However, $P(HD|E,D)$ cannot be implified




Exercise:

- *P (CP|HD, BP, E, D)*= ?




Exercise:

- *P (CP|HD, BP, E, D)*= No simplification



Calculate P(HD=yes) ?

Calculate P(HD=yes) ?




$$P(HD = Yes) = \sum_{\alpha}\sum_{\beta} P(HD = yes \mid E = \alpha, D = \beta)P(E = \alpha, D = \beta)$$

where,

α = Set of Values of Exercise(E) = {Yes, No}

β = Set of Values of Diet(D) = {Healthy, Not Healthy}


Calculate P(HD=yes) ?



$$P(HD=Yes)=\sum_{\alpha}\sum_{\beta}P(HD=yes\,|\,E=\alpha,D=\beta)P(E=\alpha,D=\beta)$$


$$=\sum_{\alpha}\sum_{\beta}P(HD=yes\,|\,E=\alpha,D=\beta)P(E=\alpha)P(D=\beta)$$

$$=0.25\times0.7\times0.25+0.45\times0.7\times0.75+0.55\times0.3\times0.25$$
$$+0.75\times0.3\times0.75$$
$$=0.49$$




Calculate P(HD=yes|BP=High)?

Calculate P(HD=yes|BP=High)



$P(HD = yes \mid BP = High)$ can be written as $\frac{P(BP = High \mid HD = yes)P(HD = yes)}{P(BP = High)}$

Calculate P(HD=yes|BP=High)




$$P(BP = High) = \sum_{\gamma} P(BP = high \mid HD = \gamma) P(HD = \gamma)$$

where,

γ = Set of Values of Heart Disease (HD) = {Yes, No}


Calculate P(HD=yes|BP=High)



$$P(BP = High) = \sum_{\gamma} P(BP = high \mid HD = \gamma) P(HD = \gamma)$$


$$= 0.85 \times 0.49 + 0.2 \times 0.51 = 0.5185$$

Calculate P(HD=yes|BP=High)




$$P(HD = yes \mid BP = High) = \frac{P(BP = High \mid HD = yes)P(HD = yes)}{P(BP = High)}$$

$$= \frac{0.85 \times 0.49}{0.5185} = 0.8033$$



Calculate P(HD=yes|BP=high, D=Healthy, E=yes)?

Calculate P(HD=yes|BP=high, D=Healthy, E=yes) ?



$$P(HD = yes \mid BP = high, D = Healthy, E = Yes)$$

$$= \frac{P(BP = high \mid HD = yes, D = Healthy, E = Yes)}{P(BP = high \mid D = Healthy, E = Yes)} \times P(HD = yes \mid D = Healthy, E = Yes)$$

How is this formula true?

$$P(HD = yes \mid BP = high, D = Healthy, E = Yes)$$
$$= \frac{P(BP = high \mid HD = yes, D = Healthy, E = Yes)}{P(BP = high \mid D = Healthy, E = Yes)} \times P(HD = yes \mid D = Healthy, E = Yes)$$

Let $$P(X \mid Y) = \frac{P(Y \mid X)}{P(Y)} \times P(X)$$

Now add *Z* and *W* as condition $$P(X \mid Y, Z, W) = \frac{P(Y \mid X, Z, W)}{P(Y \mid Z, W)} \times P(X \mid Z, W)$$

Similarly,

$$P(HD = yes \mid BP = high) = \frac{P(BP = high \mid HD = yes)}{P(BP = high)} \times P(HD = yes)$$

Now add conditions *D= Healthy* and *E= Yes* to above formula


Similarly,

$$P(BP = high \mid D = Healthy, E = Yes)$$
$$= \sum_{\gamma} P(BP = high \mid HD = \gamma, D = Healthy, E = Yes) \times P(HD = \gamma \mid D = Healthy, E = Yes)$$

Proof:


$$P(BP = high) = \sum_{\gamma} P(BP = high \mid HD = \gamma) \times P(HD = \gamma)$$

Adding conditions *D= Healthy* and *E= Yes*



we get,

$$P(BP = high \mid D = Healthy, E = Yes)$$
$$= \sum_{\gamma} P(BP = high \mid HD = \gamma, D = Healthy, E = Yes) \times P(HD = \gamma \mid D = Healthy, E = Yes)$$
$$= \sum_{\gamma} P(BP = high \mid HD = \gamma) \times P(HD = \gamma \mid D = Healthy, E = Yes)$$



$$P(HD = yes \mid BP = high, D = Healthy, E = Yes)$$

$$= \frac{P(BP = high \mid HD = yes, D = Healthy, E = Yes)}{P(BP = high \mid D = Healthy, E = Yes)} \times P(HD = yes \mid D = Healthy, E = Yes)$$

$$= \frac{P(BP = high \mid HD = yes)}{P(BP = high \mid D = Healthy, E = Yes)} \times P(HD = yes \mid D = Healthy, E = Yes)$$

$$= \frac{P(BP = high \mid HD = yes)}{\sum_{\gamma} P(BP = high \mid HD = \gamma) P(HD = \gamma \mid D = Healthy, E = Yes)} \times P(HD = yes \mid D = Healthy, E = Yes)$$

$$= \frac{0.85 \times 0.25}{0.85 \times 0.25 + 0.2 \times 0.75} = 0.5862$$