

GATE-EC 2016 Q.33

Nelakuditi Rahul Naga - AI20BTECH11029

August 28, 2021

Question

GATE-EC 2016 Q.33

The Discrete Fourier Transform (DFT) of the 4 point sequence $x[n] = \{3, 2, 3, 4\}$ is given by $X[k] = \{12, 2j, 0, -2j\}$. If $X_1[k]$ is the DFT of the 12 point sequence $x_1[n] = \{3, 0, 0, 2, 0, 0, 3, 0, 0, 4, 0, 0\}$, find the value of $\left| \frac{X_1[8]}{X_1[11]} \right|$.

Solution

4-point DFT matrix

The 4-point DFT matrix is given by:

$$W_4 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & \omega^1 & \omega^2 & \omega^3 \\ 1 & \omega^2 & \omega^4 & \omega^6 \\ 1 & \omega^3 & \omega^6 & \omega^9 \end{pmatrix} \quad (1)$$

$$= \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -j & -1 & j \\ 1 & -1 & 1 & -1 \\ 1 & j & -1 & -j \end{pmatrix} \quad (2)$$

$$\text{where } \omega = e^{\frac{-2\pi j}{4}} = -j.$$

Now from the given information we can write:

$$X = W_4 x \quad (3)$$

where

$$x = \begin{pmatrix} 3 \\ 2 \\ 3 \\ 4 \end{pmatrix} \quad (4)$$

and

$$X = \begin{pmatrix} 12 \\ 2j \\ 0 \\ -2j \end{pmatrix} \quad (5)$$

Now we need to find X_1 satisfying the relation :

$$X_1 = W_{12} x_1 \quad (6)$$

where

$$x_1 = \begin{pmatrix} 3 \\ 0 \\ 0 \\ 2 \\ 0 \\ 0 \\ 3 \\ 0 \\ 0 \\ 4 \\ 0 \\ 0 \end{pmatrix} \quad (7)$$

and W_{12} is the 12-point DFT matrix.

12-point DFT matrix

$$W_{12} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & \Omega & \Omega^2 & -j & -j\Omega & -j\Omega^2 & -1 & -\Omega & -\Omega^2 & j & j\Omega & j\Omega^2 \\ 1 & \Omega^2 & -j\Omega & -1 & -\Omega^2 & j\Omega & 1 & \Omega^2 & -j\Omega & -1 & -\Omega^2 & j\Omega \\ 1 & -j & -1 & j & 1 & -j & -1 & j & 1 & -j & -1 & j \\ 1 & -j\Omega & -\Omega^2 & 1 & -j\Omega & -\Omega^2 & 1 & -j\Omega & -\Omega^2 & 1 & -j\Omega & -\Omega^2 \\ 1 & -j\Omega^2 & j\Omega & -j & -\Omega^2 & \Omega & -1 & j\Omega^2 & -j\Omega & j & \Omega^2 & -\Omega \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \\ 1 & -\Omega & \Omega^2 & j & -j\Omega & j\Omega^2 & -1 & \Omega & -\Omega^2 & -j & j\Omega & -j\Omega^2 \\ 1 & -\Omega^2 & -j\Omega & 1 & -\Omega^2 & -j\Omega & 1 & -\Omega^2 & -j\Omega & 1 & -\Omega^2 & -j\Omega \\ 1 & j & -1 & -j & 1 & j & -1 & -j & 1 & j & -1 & -j \\ 1 & j\Omega & -\Omega^2 & -1 & -j\Omega & \Omega^2 & 1 & j\Omega & -\Omega^2 & -1 & -j\Omega & \Omega^2 \\ 1 & j\Omega^2 & j\Omega & j & -\Omega^2 & -\Omega & -1 & -j\Omega^2 & -j\Omega & -j & \Omega^2 & \Omega \end{pmatrix}$$

where

$$\Omega = e^{\frac{-2\pi j}{12}} = \frac{\sqrt{3}-j}{2} \quad (8)$$

Only the red coloured columns in W_{12} give non-zero output when multiplied with x_1 . We can express the matrix W_{12} as a block matrix in the following way:

$$W_{12} = (\mathbf{c}_1 \quad c_2 \quad c_3 \quad \mathbf{c}_4 \quad c_5 \quad c_6 \quad \mathbf{c}_7 \quad c_8 \quad c_9 \quad \mathbf{c}_{10} \quad c_{11} \quad c_{12}) \quad (9)$$

where c_i is the i^{th} column matrix of W_{12} . Now we have :

$$X_1 = (\mathbf{c}_1 \quad c_2 \quad c_3 \quad \mathbf{c}_4 \quad c_5 \quad c_6 \quad \mathbf{c}_7 \quad c_8 \quad c_9 \quad \mathbf{c}_{10} \quad c_{11} \quad c_{12}) \begin{pmatrix} 3 \\ 0 \\ 0 \\ 2 \\ 0 \\ 0 \\ 0 \\ 3 \\ 0 \\ 0 \\ 4 \\ 0 \\ 0 \end{pmatrix} \quad (10)$$

The above equation on simplification gives :

$$X_1 = 3c_1 + 2c_4 + 3c_7 + 4c_{10} \quad (11)$$

We can also express W_4 as block matrix as follows:

$$W_4 = (w_1 \quad w_2 \quad w_3 \quad w_4) \quad (12)$$

where w_i is the i^{th} column matrix of W_4 . Now from (3) we can write:

$$\begin{pmatrix} 12 \\ 2j \\ 0 \\ -2j \end{pmatrix} = (w_1 \quad w_2 \quad w_3 \quad w_4) \begin{pmatrix} 3 \\ 2 \\ 3 \\ 4 \end{pmatrix} \quad (13)$$

$$\Rightarrow 3w_1 + 2w_2 + 3w_3 + 4w_4 = \begin{pmatrix} 12 \\ 2j \\ 0 \\ -2j \end{pmatrix} \quad (14)$$

We can importantly note that :

$$c_1 = \begin{pmatrix} w_1 \\ w_1 \\ w_1 \end{pmatrix} \quad (15)$$

$$c_4 = \begin{pmatrix} w_2 \\ w_2 \\ w_2 \end{pmatrix} \quad (16)$$

$$c_7 = \begin{pmatrix} w_3 \\ w_3 \\ w_3 \end{pmatrix} \quad (17)$$

$$c_{10} = \begin{pmatrix} w_4 \\ w_4 \\ w_4 \end{pmatrix} \quad (18)$$

$$\therefore X_1 = 3c_1 + 2c_4 + 3c_7 + 4c_{10} = \begin{pmatrix} 3w_1 + 2w_2 + 3w_3 + 4w_4 \\ 3w_1 + 2w_2 + 3w_3 + 4w_4 \\ 3w_1 + 2w_2 + 3w_3 + 4w_4 \end{pmatrix} \quad (19)$$

From (14) and (19) we have :

$$X_1 = \begin{pmatrix} 12 \\ 2j \\ 0 \\ -2j \\ 12 \\ 2j \\ 0 \\ -2j \\ 12 \\ 2j \\ 0 \\ -2j \end{pmatrix} \quad (20)$$

Therefore we have:

$$\left| \frac{X_1[8]}{X_1[11]} \right| = \left| \frac{12}{-2j} \right| = |6j| = 6 \quad (21)$$




Figure: Magnitude of $X[k]$ vs k




Figure: Magnitude of $X_1[k]$ vs k