10.05.2.3

EE23BTECH11053-R.Rahul*

QUESTION:

1. In the following APs, find the missing terms in the boxes:

$$(iii)5, _, _, 9\frac{1}{2}$$

(ii) _, 13, _, 3
(iii)5, _, _,9
$$\frac{1}{2}$$

(iv)- 4, _, _, _, _, 6
(v)_,38, _, _, _,'-22'

Solution:

1)

Parameter	Description
n	No. of terms in the A.P
<i>x</i> (0)	first term in the A.P
d	common difference in the A.P
x(n) = x(0) + nd	$(n+1)^{th}$ term in A.P

TABLE I

VARIABLES

The Z-transform of x(n) = 18 - 5n is given by:

$$X(z) = \frac{18 - 23z^{-1}}{(1 - z^{-1})^2} \qquad |z| > 1$$
 (12)

(13)

(19)

(20)

(21)

(22)

(23)

(24)

3)

$$9\frac{1}{2} = 5 + 3d\tag{14}$$

$$3d = \frac{9}{2} \tag{15}$$

$$\therefore d = \frac{3}{2} \tag{16}$$

$$x(1) = 6 \frac{1}{2} \tag{17}$$

$$x(2) = 8 \tag{18}$$

Z-transform of $x(n) = 5 + \frac{3}{2}n$ is given by:

 $X(z) = \frac{5 - \frac{7}{2}z^{-1}}{(1 - z^{-1})^2} \qquad |z| > 1$

6 = -4 + 5d

10 = 5d

 $\therefore d = 2$

$$26 = 2 + 2d \tag{1}$$

$$24 = 2d \tag{2}$$

$$\therefore d = 12 \tag{3}$$

$$x(1) = 14 \tag{4}$$

The Z-transform of x(n) = 2 + 12n is given by:

$$X(z) = \frac{2 + 10z^{-1}}{(1 - z^{-1})^2} \qquad |z| > 1$$
 (5)

(6)

4)

$$x(1) = -2$$
$$x(2) = 0$$

(25)

$$x(3) = 2 \tag{26}$$

$$x(4) = 4 \tag{27}$$

2)

$$3 - 13 = 2d (7)$$

$$-10 = 2d \tag{8}$$

$$\therefore d = -5 \tag{9}$$

$$x(1) = 18 (10)$$

$$x(2) = 8 \tag{11}$$

Z-transform of x(n) = -4 + 2n is given by:

$$X(z) = \frac{-4 + 6z^{-1}}{(1 - z^{-1})^2} \qquad |z| > 1$$
 (28)

(29)

5)

$$-22 - 38 = 4d \tag{30}$$

$$-60 = 4d$$
 (31)

$$\therefore d = -15 \tag{32}$$

$$x(0) = 53$$
 (33)

$$x(2) = 23 \tag{34}$$

$$x(3) = 8 \tag{35}$$

$$x(4) = -7 (36)$$

(37)

Z-transform of x(n) = 53 - 15n is given by:

$$X(z) = \frac{53 - 68z^{-1}}{(1 - z^{-1})^2} \qquad |z| > 1 \qquad (38)$$

(39)




Fig. 2.




Fig. 1.




Fig. 3.




Fig. 4.




Fig. 5.