

Clinical Reasoning & Explainability Report

Patient: See Wuckert | Cycle #8

1. Patient Context

Cancer Type: BREAST
Age: 56 | Gender: F
Chemotherapy Drugs: Tamoxifen, Paclitaxel, Doxorubicin

2. Clinical Data Inputs & Their Impact

2a. Laboratory Values

No lab values available.

2b. Wearable Biometric Data

Metric	Value	Normal	Status	Impact
HRV (RMSSD)	34.1 ms	>30 ms	NORMAL	Healthy autonomic function
Sleep Efficiency	91%	>85%	NORMAL	Adequate sleep quality
Mean Heart Rate	72 bpm	60-100	INFO	Contextual cardiovascular data


2c. Age Modifier

Age 56: No age modifier applied


3. Drug Side-Effect Profiles

Base severity weights for each drug (0-1 scale, higher = more likely/severe)


Tamoxifen -- Known Side Effect Affinities



Paclitaxel -- Known Side Effect Affinities



Doxorubicin -- Known Side Effect Affinities





4. Temporal Severity Model

Side-effect severity follows a temporal curve across the 7-day chemo cycle. Day 1 (infusion) starts mild, severity peaks at Days 3-5 (nadir period when blood counts are lowest), then eases on Days 6-7 (recovery). This pattern is well-established in oncology literature.

Day	Multiplier	Phase
Day 1	0.4x	Infusion
Day 2	0.7x	Onset
Day 3	1.0x	Nadir (Peak)
Day 4	1.0x	Nadir (Peak)
Day 5	0.9x	Late Nadir
Day 6	0.6x	Recovery
Day 7	0.4x	Recovery

5. Per-Side-Effect Severity Reasoning

Final Grade = Drug Base x Clinical Modifier x Temporal Curve x Cycle Factor + Random Noise

Fever	None	None	Mild	Mode	Mild	None	None	Tamoxifen base affinity: 10% Paclitaxel base affinity: 25% Doxorubicin base affinity: 35%
Fatigue	Mild	Mode	Seve	Seve	Seve	Mode	Mild	Tamoxifen base affinity: 45% Paclitaxel base affinity: 70% Doxorubicin base affinity: 80%
Nausea & Vomiting	Mild	Seve	Seve	Seve	Seve	Seve	Mild	Tamoxifen base affinity: 30% Paclitaxel base affinity: 50% Doxorubicin base affinity: 90%
Sore Mouth	Mild	Mild	Seve	Mode	Mode	Mode	Mild	Tamoxifen base affinity: 10% Paclitaxel base affinity: 30% Doxorubicin base affinity: 55%
Diarrhea	None	Mild	Mild	Mild	Mild	None	Mild	Tamoxifen base affinity: 15% Paclitaxel base affinity: 35% Doxorubicin base affinity: 30%
Constipation	None	None	None	Mild	Mild	None	None	Tamoxifen base affinity: 15% Paclitaxel base affinity: 25% Doxorubicin base affinity: 25%
Loss of Appetite	Mild	Mode	Mode	Seve	Mode	Mild	Mild	Tamoxifen base affinity: 20% Paclitaxel base affinity: 45% Doxorubicin base affinity: 60%
Swallowing Difficulty	None	Mild	None	None	None	None	None	Tamoxifen base affinity: 5% Paclitaxel base affinity: 10% Doxorubicin base affinity: 15%
Anxiety / Depression	None	Mild	Mode	Mode	Mild	Mild	Mild	Tamoxifen base affinity: 35% Paclitaxel base affinity: 35% Doxorubicin base affinity: 45%
Edema (Hands/Feet)	None	None	Mild	Mild	None	None	None	Tamoxifen base affinity: 25% Paclitaxel base affinity: 30% Doxorubicin base affinity: 20%
Itching / Rash	None	None	None	Mild	Mild	None	None	Tamoxifen base affinity: 15% Paclitaxel base affinity: 20% Doxorubicin base affinity: 20%
Shortness of Breath	None	None	Mild	None	Mild	Mild	None	Tamoxifen base affinity: 10% Paclitaxel base affinity: 20% Doxorubicin base affinity: 25%
Muscle / Joint Pain	Mild	Mode	Seve	Seve	Seve	Mode	Mild	Tamoxifen base affinity: 40% Paclitaxel base affinity: 75% Doxorubicin base affinity: 30%
Numbness / Tingling	Mild	Mode	Seve	Seve	Seve	Mild	Mild	Tamoxifen base affinity: 10% Paclitaxel base affinity: 80% Doxorubicin base affinity: 15%

6. Methodology & Data Sources

- SEVERITY CALCULATION: $\text{Final severity} = \text{Drug_Base_Weight} \times \text{Clinical_Modifier} \times \text{Temporal_Curve} \times \text{Cycle_Factor} + \text{Gaussian_Noise}(0, 0.08)$
- DRUG PROFILES: Based on published chemotherapy side-effect incidence rates from oncology literature.
- CLINICAL MODIFIERS: Lab values (WBC, Hemoglobin, GFR, Pain) adjust severity multipliers based on clinical thresholds.
- WEARABLE DATA: HRV (RMSSD) and sleep efficiency provide real-time physiological stress indicators.
- TEMPORAL MODEL: 7-day cycle follows established nadir pattern (peak toxicity Days 3-5 post-infusion).
- CYCLE FACTOR: Later cycles accumulate toxicity at +2% per cycle (capped at 40% increase).
- GRADING: Continuous 0-1 score mapped to 4 grades: None (<0.20), Mild (0.20-0.44), Moderate (0.45-0.69), Severe (>=0.70).
- DATA SOURCES: Synthea synthetic patient data (demographics, conditions, medications, observations, encounters) + generated wearable biometrics.