



Imaging Healthcare Specialists - Hillcrest
150 W. Washington St
San Diego, CA 92103
Phone #: (858)658-6500

Your Partner in Health - For Every Imaging Need

Online results, images and scheduling available at:
imaginghealthcare.com/results

Patient:	MARTINEZ, LELIA M	DOB:	11/09/1948	Age:	75
Exam Date:	04/27/2024	ID:	30000062756	Sex:	Female
Accession #:	IHS3646810	CC:			
At the	DOWNING, KRISTOPHER, MD				
Request of:	12264 EL CAMINO REAL SUITE 101				
	SAN DIEGO, CA 92130				

PROCEDURE:
MR SHOULDER RT W/O CONTRAST

COMPARISON:
None.

INDICATIONS:
Right shoulder pain .

TECHNIQUE:
A comprehensive examination was performed without intravenous gadolinium contrast utilizing a variety of imaging planes and imaging parameters to optimize visualization of suspected pathology.

FINDINGS:
ROTATOR CUFF REGION
CUFF TENDONS: There is a large full-thickness tear that involves the entire supraspinatus tendon and the anterior fibers of the infraspinatus tendon with retraction of the torn tendon fibers to the superior glenoid. There is a high-grade (75-90%) tear of the subscapularis tendon as well.
CUFF MUSCLES: There is moderate atrophy of the supraspinatus muscle belly. See below. The other muscles are normal in bulk.
CUFF GOUTALLIER STAGE: Stage II: Less fat than muscle within the muscle.
DELTOID: Normal. No significant atrophy or tear.
LONG BICEPS TENDON: Intact with moderate to severe intra-articular tendinosis.

LABRUM/BICEPS ANCHOR
SUPERIOR: Degenerative signal in the superior labrum and the anchor but no clearly visible tear.
ANTERIOR/INFERIOR: Normal. No visible tear or attrition.
POSTERIOR: Normal. No posterior labrum abnormality.

CAPSULE
ANTERIOR/INFERIOR: Normal. No visible capsular laxity or thickening.
POSTERIOR: Normal. No visible capsular laxity or thickening.

AC JOINT REGION
AC JOINT: Mild-to-moderate osteoarthritis of the acromioclavicular joint.
AC LIGAMENTS: Normal acromioclavicular ligament.

Patient:	MARTINEZ, LELIA M	DOB:	11/09/1948	Age:	75
Exam Date:	04/27/2024	ID:	30000062756	Sex:	Female
Accession #:	IHS3646810				
At the	DOWNING, KRISTOPHER, MD	CC:			
Request of:	12264 EL CAMINO REAL SUITE 101 SAN DIEGO, CA 92130				

CC LIGAMENTS: Normal coracoclavicular ligaments.

ACROMION: Type 2 anatomic configuration with a large os acromiale and inferior acromial spurring. There is a large effusion in the synchondrosis between the acromion and the os.

SUBACROMIAL BURSA: Normal. No significant effusion.

HYALINE CARTILAGE: Normal. No visible cartilage narrowing or focal defect.

OTHER BONES: Normal proximal humerus, glenoid, and coracoid.

OTHER OBSERVATIONS: Moderate-sized glenohumeral joint effusion with a 9 x 4 mm loose body in the posterior aspect of the axillary recess.

CONCLUSION:

1. Large full-thickness tear involves the entire supraspinatus tendon and the anterior fibers of the infraspinatus tendon with tendon retraction to the superior glenoid and moderate fatty atrophy.
2. High-grade (75-90%) tear of the subscapularis tendon.
3. Moderate to severe intra-articular long biceps tendinosis.
4. Moderate-sized glenohumeral joint effusion with a 9 x 4 mm loose body in the posterior aspect of the axillary recess.
5. Mild to moderate acromioclavicular joint osteoarthritis.
6. Large os acromiale with a large effusion in the synchondrosis between the acromion and the os acromiale.

Dictated and Electronically Authenticated by: Cooper, James, M.D. on 04/27/2024 03:53 PM Proofread by: Cooper, James, M.D. on 04/27/2024 03:53 PM

IMAGING HEALTHCARE SPECIALISTS
150 W. Washington St
San Diego, CA 92103 (858)658-6500