PARKINOVA

Comprehensive Parkinson's Research Assessment

A. Patient & Assessment Information

Patient Reference	PT-101
Age	65
Gender	Male
Assessment Date	2026-10-27 10:00:00
Selected Modules	Brain MRI Scans
Report Reference ID	TEST-101

B. Brain MRI Neuroimaging Analysis

Analysis Status	Completed
Model Prediction	Healthy/Control
Model Architecture	Parkinson CNN (244×244×3)
Input Preprocessing	244×244 Lanczos Resizing, Channel Standardization
Analysis Timestamp	2026-10-27T10:00:00

Model Findings & Interpretation:

Neuroimaging pattern analysis completed.

Model Limitations:

The MRI classifier identifies spatial and intensity patterns correlated with the training cohort. An arbitrary MRI scan alone cannot independently confirm or rule out Parkinson's disease. Model confidence is not clinical probability of disease.

C. Spiral Drawing Analysis

Spiral Analysis Status	Not Submitted / In Standby
Reason	Patient has not yet completed the spiral drawing module
Impact	Combined evaluation relies solely on MRI scan and clinical questionnaire

D. Combined Multi-Modal Assessment Summary

Unimodal MRI Assessment: Assessment is currently based exclusively on brain MRI neuroimaging. Spiral analysis was not submitted. Multi-modal synthesis will update when spiral drawing evaluation is provided.

E. Parkinson's Health, Nutrition & Movement Education

Evidence-based supportive lifestyle strategies tailored for individuals with Parkinson's and clinical cohorts:

1. Indian Nutrition & Antioxidants	Focus on neuroprotective, antioxidant-rich foods: cooked spinach (palak), methi, drumstick sambar, ripe papaya, soft ragi kanji/dosas, yellow moong dal khichdi, and soaked almonds. Avoid hard, dry, or splintery textures.
2. Levodopa & Protein Coordination	Dietary large neutral amino acids (LNAAs) compete with levodopa for transport across gut and blood-brain barrier. Take Levodopa 30-60 minutes before or 1-2 hours after protein meals. Do NOT eliminate essential dietary protein.
3. Hydration & Gut Motility	Autonomic constipation affects up to 80% of individuals with Parkinson's. Maintain 6-8 glasses of fluid daily (lukewarm jeera/cumin water, tender coconut water) and adequate soluble fiber (oats, stewed prunes).
4. Safe Physical Activity & Fall Prevention	Daily walking with rhythmic auditory/counting cues (1-2-1-2 cadence), daily calf/chest stretching, supervised balance exercises, and proactive home fall-proofing (removing loose throw rugs, installing night lights).

Regulatory Notice & Academic Disclaimer

This application is an academic AI research prototype. Its output is generated by a machine-learning model and is not a medical diagnosis. The model's performance and reliability may vary depending on image quality, dataset characteristics, and the images used during training. Consult a qualified neurologist for medical evaluation.