

## Aluminium – Calcium – Silicon

*Rainer Schmid-Fetzer*

### Literature Data

The liquidus surface of the Al-corner has been studied and fundamental agreement was obtained between most of the earlier experimental work [1926Doa, 1927Gro, 1928Shi] and reviewing papers [1934Fus, 1943Mon, 1952Han, 1960Spe, 1969Wat]. Samples were made from Al–Ca and Al–Si master alloys under a  $\text{CaCl}_2$  or  $\text{KCl–NaCl–BaCl}_2$  protective layer. Their composition was checked by chemical analysis, and thermal and metallographic analyses were performed [1926Doa]. A dominating  $\text{CaSi}_2$  liquidus surface extending close to the Al-corner and an eutectic type Al– $\text{CaSi}_2$  pseudobinary system was deduced from these data [1926Doa] and essentially confirmed by similar studies [1928Shi, 1962Kol] and [1966Tom]. The coexistence of (Al) and a compound, assumed to be  $\text{CaSi}_2$ , was also seen in micrographs of Al-rich samples that were cast, forged or hot rolled, annealed for 30 min at  $460^\circ\text{C}$  and then used for Brinell and tensile tests [1927Gro]. Similar interpretations are given in [1934Fus, 1943Mon] and also by [1952Han] who made additional experiments to quantify the location of ternary eutectics and accepts the binary phase  $\text{CaAl}_4$  instead of  $\text{CaAl}_3$  as given in previous literature. The interpretation of an (Al)– $\text{CaSi}_2$  equilibrium is, however, inconsistent with reports on some ternary compounds that form after annealing for a long time. [1967Gla] studied 39 samples in the Al–Si– $\text{CaSi}_2$ – $\text{CaAl}_2$  subsystem at  $400^\circ\text{C}$  by X-ray methods and metallography and detected the compound  $\text{CaAl}_2\text{Si}_2$  ( $\beta$ ) in equilibrium with (Al), (Si),  $\text{CaSi}_2$ ,  $\text{CaAl}_4$ ,  $\text{CaAl}_2$  and another ternary compound,  $\text{CaAl}_{1-x}\text{Si}_{1+x}$  ( $\gamma$ ) [1965Bod]. [1967Gla] also prepared single crystals of  $\beta$  and performed a detailed X-ray structure analysis. The  $\beta$  phase was also observed by electron microprobe in Al-20 mass% Si samples containing 0.13 to 2.6 mass% Ca, which were equilibrated at  $800^\circ\text{C}$ , cast and found to consist of (Al)+(Si)+ $\beta$  [1976Tag]. The precipitation of  $\beta$  and  $\text{Ca}_2\text{Si}$  was found in metallurgical grade silicon ( $< 0.5$  mass% Al,  $< 0.3$  mass% Ca) by optical metallography and electron microscopy (SEM, TEM, EMPA) [1990Ang].

The  $\text{CaAl}_{1-x}\text{Si}_{1+x}$  ( $\gamma$ ) phase was also prepared at  $x = 0$  by [2002Ima] and in the range  $-0.4 < x < 0.2$  by [2002Lor], studied by X-ray diffraction and also for their superconducting behavior with a transition temperature of 7.8 K for  $\text{CaAlSi}$  [2002Ima].

The congruent melting point of  $\beta$  was reported at  $975^\circ\text{C}$  by DTA [1994Ang] and the pseudobinary eutectic  $\text{L}=\beta+(\text{Si})$  at  $927^\circ\text{C}$ , and the “pseudobinary” eutectic  $\text{L}=\beta+\text{CaSi}_2$  at  $925^\circ\text{C}$ . [1994Ang] also performed a brief Calphad-type thermodynamic analysis of the ternary system and produced calculated isopleths. This included the modelling of the Ca-activity in ternary liquids on the  $\text{Al}_{0.5}\text{Si}_{0.5}$ –Ca section at  $1350^\circ\text{C}$ . This Ca-activity was experimentally determined by [1975Sch] with 11 alloys at  $1350^\circ\text{C}$  and 3 alloys at  $1210^\circ\text{C}$  using the boiling point method. The activity coefficients of Al and Ca in molten Si at  $1450$ – $1550^\circ\text{C}$  were determined by the Knudsen effusion method and also by a chemical equilibrium technique from the distribution between liquid silicon and lead at  $1450^\circ\text{C}$  [1999Mik]. Two other compounds,  $\text{Ca}_3\text{Al}_6\text{Si}_2$  ( $\delta$ ) and  $\text{Ca}_2\text{Al}_4\text{Si}_3$  ( $\epsilon$ ) were prepared by pressing  $\text{CaAl}_2$ –Si mixtures, heating to  $700$ – $1000^\circ\text{C}$ , quenching and studying by X-ray analysis [1955Chr, 1956Chr]. The reaction was complete only above  $900^\circ\text{C}$  and the mutual equilibria  $\text{CaAl}_2+\delta$ ,  $\delta+\epsilon$  and  $\epsilon+(\text{Si})$  were observed at  $1000^\circ\text{C}$ , but not the  $\beta$  phase located between  $\epsilon$  and (Si). The  $\delta$  phase completely decomposes into  $\epsilon$  and some Al and Ca at  $1200^\circ\text{C}$ , while the  $\epsilon$  phase only starts to decompose at this temperature, probably due to Ca-loss to the gas phase [1956Chr]. The most Ca-rich compound,  $\text{Ca}_3\text{Al}_2\text{Si}_2$  ( $\alpha$ ) was fused from the elements in corundum crucibles under argon, continuously agitated for 4h at  $1150$ – $1200^\circ\text{C}$  and slowly cooled to room temperature within 12h. Single crystals could be extracted from the sample and a detailed X-ray structure analysis was performed [1977Wid]. The composition change during preparation was found to be negligible by chemical analysis of a similarly prepared  $\text{Ba}_3\text{Al}_2\text{Si}_2$  compound [1977Wid].

The present evaluation continues and updates the one published in [1990Sch] with respect to both new ternary and binary data.

## Binary Systems

The Al–Si system is accepted from [Mas2]. The Al–Ca system is taken from [2002Ted]. The Ca–Si system is accepted from [2000Man], however, the shape of the liquidus lines may need some revision as indicated by thermodynamic calculations [2003Gro].

## Solid Phases

Data on all solid phases reported in this system are given in Table 1.

Electronic structure calculations of the stability of  $\text{Al}_2\text{Si}_2^{6-}$  chains in the compound  $\text{Ca}_3\text{Al}_2\text{Si}_2$  suggest that all Si atoms reside in the twofold sites and all Al atoms in the threefold ones [1988Li].

Bonding in the  $\text{CaAl}_2\text{Si}_2$  ( $\beta$ ) structure type was studied theoretically [1988Zhe]. The same structure type was experimentally found to form also in 13 different, though related, ternary systems [1980Klu].

[1975Eml] mentioned the possible existence of a phase  $\text{Ca}_{0.8}\text{Al}_{1.2}\text{Si}$ , which is not included in Table 1.

## Pseudobinary Systems

The Al– $\text{CaSi}_2$  section has been quoted as a eutectic pseudobinary system in the earlier literature [1926Doa] to [1952Han]. The primary crystallization of  $\text{CaSi}_2$  from Al-rich liquids is probably metastable in view of the formation of  $\beta$  on that section. The  $\beta$  phase may have been misinterpreted as  $\text{CaSi}_2$  in the metallographic examination, since the conclusions in the basic early work on Al–Ca–Si [1926Doa] rely on the assumption that no ternary compounds exist. The calculated isopleths Al– $\text{CaSi}_2$  and Si– $\text{CaAl}_2$  [1994Ang] contain the  $\beta$  phase, however, they are based on old versions of the Ca–Si and Al–Ca binaries and cannot be accepted. In addition, the phases  $\alpha$ ,  $\gamma$ ,  $\delta$  and  $\epsilon$  had not been considered by [1994Ang]. Thermodynamic calculations performed in the present assessment were based on the recent binaries and they show that both sections are not pseudobinary systems. The Al-poor side of the Al– $\text{CaSi}_2$  section exhibits phase fields of liquid with  $\text{CaSi}$  and  $\text{Ca}_{14}\text{Si}_{19}$ . However the partial section Al– $\text{CaAl}_2\text{Si}_2$  ( $\beta$ ) is a pseudobinary eutectic as shown in Fig. 1, according to the present calculation. Similarly, the Si-poor side of the Si– $\text{CaAl}_2$  section exhibits phase fields of liquid with  $\text{CaSi}$ . However the partial section Si– $\text{CaAl}_2\text{Si}_2$  ( $\beta$ ) is also a pseudobinary eutectic as shown in Fig. 2, according to the present calculation. The calculated eutectic temperature 933°C is in close agreement with the experimental value 927°C [1994Ang].

## Invariant Equilibria

The three invariant equilibria of the Al-corner, given in Table 2, are from the present thermodynamic calculation. Earlier work [1952Han] assumed the participation of  $\text{CaSi}_2$  instead of  $\beta$  in the equilibria max, E and D (with reported temperatures of 637, 615.8, and 576.5°C). This cannot be accepted as described in the previous section. Also a eutectic reaction  $\text{L} \rightleftharpoons (\text{Al}) + (\text{Si}) + \beta$  was given with a liquid composition of 0.7 at.% Ca [1966Tom]. This is virtually impossible based on the thermodynamic calculation, which shows that this invariant (D) is essentially degenerate to the binary Al–Si eutectic. [1966Tom] also reported from microradiograph examinations that the Ca-content in (Si) is much smaller than in (Al).

## Liquidus Surface

The liquidus surface of the Al-corner given in Fig. 3 is from the present thermodynamic calculation. It deviates from that given by [1952Han] as discussed in the previous section.

## Isothermal Sections

An isothermal section at about 400°C is given in Fig. 4 [1967Gla, 1955Chr] and [1956Chr]. The equilibria around the  $\epsilon$  and  $\delta$  phases are estimated by dashed lines. Both phases form above 900°C with appreciable reaction rates [1955Chr] and are presumably stable down to 400°C. The reported  $\epsilon$  – (Si) equilibrium at 1000°C [1955Chr] cannot be accepted in view of the congruent melting point of  $\beta$  at 975°C.

The equilibria above 33 at.% Ca have not been studied experimentally. The dashed tie line  $\text{Ca}_5\text{Si}_3$ – $\text{CaAl}_2$  (and the three more Ca-rich ones) given dashed in Fig. 4 are based on the present thermodynamic

calculation, disregarding the  $\alpha$  and  $\gamma$  phases and the mutual solubilities along the  $\text{CaAl}_2$ – $\text{CaSi}_2$  section. No tie lines can be given around these phases.

The precipitation of  $\beta$  and  $\text{Ca}_2\text{Si}$  in metallurgical grade silicon ( $< 0.5$  mass% Al,  $< 0.3$  mass% Ca) [1990Ang] supports the existence of the  $(\text{Si}) + \beta + \text{Ca}_2\text{Si}$  equilibrium with negligible solubility in (Si).

### Thermodynamics

The measured Ca-activities [1975Sch] in the ternary liquid phase along the equal molar fractions of Al and Si at  $1350^\circ\text{C}$  are well represented by the thermodynamic calculation of [1994Ang] shown in Fig. 5

### Notes on Materials Properties and Applications

Si and Ca are important additions to Al-alloys. The  $\text{CaAl}_{1-x}\text{Si}_{1+x}(\gamma)$  phase shows superconducting behavior [2002Ima, 2002Lor]. The de-oxidation of steel using complex Ca–Al–Si de-oxidizers was discussed by [1983Gho].

### Miscellaneous

The dissolution of  $\text{CaSi}_2$  from Al–Ca–Si alloys in HCl was discussed by [1953Tou].

### References

- [1926Doa] Doan, G., “On the Al–Ca–Si System” (in German), *Z. Metallkd.*, **18**, 350–355 (1926) (Equi. Diagram, Experimental, #, 15)
- [1927Gro] Grogan, J.D., “The Influence of Calcium on Aluminium Containing Silicon”, *J. Inst. Met.*, **37**, 77–91 (1927) (Experimental, 6)
- [1928Shi] Shinoda, G., “Improvement of Aluminium Alloy with X-Ray Study” (in Japanese), *Nippon Kogyo Kwai Shi*, **44**, 544–557 (1928) (Equi. Diagram, Experimental, 12)
- [1934Fus] Fuss, V., *Metallography of Al and its Alloys* (in German), translated from Anderson, R.J., Sherwood Press Inc., Cleveland, Berlin, 163–166 and 207–219 (1936) (Equi. Diagram, Review, #, 300)
- [1943Mon] Mondolfo, L.F., *Metallography of Aluminum Alloys*, Wiley & Sons, Inc., New York, 61–63 (1943) (Equi. Diagram, Review, #, 3)
- [1952Han] Haneman, H., Schrader, A., “Ternary Aluminum-Alloys” in “*Atlas Metallographicus (III/2)*” (in German), Verlag Stahleisen, Düsseldorf, 62–65 (1952) (Equi. Diagram, Crys. Structure, Review, Experimental, #, \*, 5)
- [1953Tou] Tournaire, M., Renouard, M., “Contribution to a Study of Al–Mg–Si Alloys” (in French), *Revue de Metallurgie/Memoires*, **50**, 328–332 (1953) (Experimental, 0)
- [1955Chr] Chrétien, A., Freundlich, W., Deschanvres, A., “On Two New Ternary Ca–Al–Si Compounds” (in French), *Compt. Rend.*, **241**, 1781–1783 (1955) (Equi. Diagram, Experimental, #, 2)
- [1956Chr] Chrétien, A., Freundlich, W., Deschanvres, A., “Properties of the New Ternary Compounds,  $\text{Ca}_3\text{Al}_6\text{Si}_2$  and  $\text{Ca}_2\text{Al}_4\text{Si}_3$ ” (in French), *Compt. Rend.*, **242**, 784–785 (1956) (Crys. Structure, Experimental, 2)
- [1960Spe] Spengler, H., “The Importance of Research on Eutectics and its Application to Ternary Eutectic Aluminium Alloys” (in German), *Metallwiss. Tech.*, **14**(3), 201–206 (1960) (Experimental, 11)
- [1962Kol] Kolachev, B.A., “Quasi-Binary Sections of Ternary Systems, Studied by Withdrawing a Solid Phase from a Melt” (in Russian), *Tr. Mosk. Aviats. Tekhnol. Inst.*, **5**, 124–132 (1962) (Equi. Diagram, Experimental, #, 4)
- [1965Bod] Bodak, O.I., Gladyshevsky, E.I., Zarechnyuk, O.S., Cherkashin, E.E., “Compounds with Structures of the  $\text{AlB}_2$  Type in the Ternary Systems” (in Russian), *Visn. Lviv. Univ., Ser. Khim.*, **8**, 75–79 (1965) (Crys. Structure, Experimental, 8)

- [1966Tom] Tomochiro, I., Yamamoto, R., Fujisawa, K., “Mechanism of Structural Changes of Lo-Ex” (in Japanese), *Keikinzoku*, **16**, 30-35 (1966) (Equi. Diagram, Experimental, #, 2)
- [1967Gla] Gladyshevsky, E.I., Kripyakevich, P.I., Bodak, O.I., “Crystal Structure of  $\text{CaAl}_2\text{Si}_2$  and Analogous Compounds” (in Russian), *Ukrain. Fiz. Zhur.*, **12**, 447-453 (1967) (Equi. Diagram, Experimental, #, \*, 16)
- [1969Wat] Watanabe, H., Sato, E., “Phase Diagrams of Aluminum-Base Systems” (in Japanese), *Keikinzoku*, **19**(11), 499-535 (1969) (Equi. Diagram, 232)
- [1975Eml] Emlin, B.I., Gasik, M.I., Kileso, S.N., Elt’sova, Z.V., Vukelich, S.B., “Phase Composition of Al-Si Alloys” (in Russian), *Izvest. v. u. z. Tsvetn. Metall.*, **4**, 156-158 (1975) (Equi. Diagram, Experimental, 7)
- [1975Sch] Schuermann, E., Fuebers, P., Litterscheidt, H., “Vapor Pressure of Ca Above Ca-Si, Ca-Al And Ca-Al-Si Alloys” (in German), *Arch. Eisenhuettenwes.*, **46**(8), 473-476 (1975) (Experimental, Thermodyn., 5)
- [1976Tag] Tagami, M., Serita, Y., Komatsu, K., “The Influence of Ca Additions on the Grain Size of Primary Si Crystals in a Hypereutectic Al-20% Si Alloy”, *J. Jpn. Inst. Light Met.*, **26**, 385-390 (1976) (Equi. Diagram, Experimental, \*, 20)
- [1977Wid] Widera, A., Schaefer, H., “Preparation and Crystal Structure of  $\text{A}_3\text{Al}_2\text{Si}_2$  (A = Ca, Sr, Ba)” (in German), *Z. Naturforsch. B*, **32B**, 1349-1351 (1977) (Crys. Structure, Experimental, 3)
- [1980Klu] Kluefers, P., Neumann, H., Mewis, A., Schuster, H.-U., “ $\text{AB}_2\text{X}_2$  Compounds with the  $\text{CaAl}_2\text{Si}_2$  Structure, VIII (1)” (in German), *Z. Naturforsch. B*, **35**(10), 1317-1318 (1980) (Crys. Structure, Experimental, 4)
- [1983Gho] Ghosh, A., Naik, V., “Deoxidation of Steel with Ca-Si-Al and Mg-Si-Al Complex Deoxidizers: a Thermodynamic Analysis”, *Tool Alloy Steels*, **17**(7), 239-244 (1983) (Experimental, 16)
- [1988Li] Li, J., Hoffmann, R., “ $\text{Ca}_3\text{Al}_2\text{Si}_2$ : An Inorganic Structure Analogous to but not Isoelectronic with Polyacene”, *J. Phys. Chem.*, 1988, **92**(4), 887-893 (1988) (Crys. Structure, Theory, 18)
- [1988Zhe] Zheng, C., Hoffmann, R., “Complementary Local and Extended Views of Bonding in the  $\text{ThCr}_2\text{Si}_2$  and  $\text{CaAl}_2\text{Si}_2$  Structures”, *J. Solid State Chem.*, **72**(1), 58-71 (1988) (Crys. Structure, Theory, 13)
- [1990Ang] Anglezio, J.C., Servant, C., Dubrous, F., “Characterization of Metallurgical Grade Silicon” *J. Mater. Res.*, **5**, 1894-1899 (1990) (Equi. Diagram, Crys. Structure, Experimental, 11)
- [1990Sch] Schmid-Fetzer, R., “Al-Ca-Si (Aluminium - Calcium - Silicon),” MSIT Ternary Evaluation Program, in *MSIT Workplace*, Effenberg, G. (Ed.), MSI, Materials Science International Services GmbH, Stuttgart; Document ID: 10.15085.1.20, (1990) (Crys. Structure, Equi. Diagram, Assessment, 15)
- [1993Mil] Miller, G.J., Li, F., Franzen, H.F., “The Structural Phase Transition in Calcium-Aluminum Compound ( $\text{CaAl}_4$ ): A Concerted Application of Landau Theory and Energy Band Theory”, *J. Am. Chem. Soc.*, **115**(9), 3739-3745 (1993) (Crys. Structure, 26)
- [1994Ang] Anglezio, J.C., Servant, C., Ansara, I., “Contribution to the Experimental and Thermodynamic Assessment of the Al-Ca-Fe-Si System - I. Al-Ca-Fe, Al-Ca-Si, Al-Fe-Si and Ca-Fe-Si Systems”, *Calphad*, **18**(3), 273-309 (1994) (Equi. Diagram, Calculation, 71)
- [1998Hua] Huang, B., Corbett, D., “Two New Binary Calcium-Aluminium Compounds:  $\text{Ca}_{13}\text{Al}_{14}$  with a Novel Two-Dimensional Aluminium Network, and  $\text{Ca}_8\text{Al}_3$ , an  $\text{Fe}_3\text{Al}$ -Type Analogue”, *Inorg. Chem.*, **37**(22), 5827-5833 (1998) (Crys. Structure, Experimental, 30)
- [1999Mik] Miki, T., Morita, K., Sano, N., “Thermodynamic Properties of Si-Al, -Ca, -Mg Binary and Si-Ca-Al, -Ti, -Fe Ternary Alloys”, *Mater. Trans., JIM*, **40**(10), 1108-1116 (1999) (Thermodyn., Experimental, 22)
- [2000Man] Manfrinetti, P., Fornasini, M.L., Palenzona, A., “The Phase Diagram of the Ca-Si System”, *Intermetallics*, **8**, 223-228 (2000) (Equi. Diagram, Crys. Structure, Experimental, #, 31)
- [2002Ima] Imai, M., Nishida, K., Kimura, T. Abe, H., “Superconductivity of  $\text{Ca}(\text{Al}_{0.5}\text{Si}_{0.5})_2$ , a Ternary Silicide with the  $\text{AlB}_2$ -Type Structure”, *Appl. Phys. Lett.*, **80**(6), 1019-1021 (2002) (Crys. Structure, Experimental, Superconduct., 26)

- [2002Lor] Lorenz, B., Lenzi, J., Cmaidalka, J., Meng, R.L., Sun, Y.Y., Xue, Y.Y., Chu, C.W., “Superconductivity in the C32 Intermetallic Compounds  $AAI_{(2-x)}Si_x$ , with A = Ca and Sr; and  $0.6 < x < 1.2$ ”, *Physica C*, **383**(3), 191-196 (2002) (Crys. Structure, Electr. Prop., Experimental, Phys. Prop., 10)
- [2002Ted] Tedenac, J.-C., Kevorkov, D., Velikanova, T., “Al–Ca (Aluminum - Calcium)”, MSIT Binary Evaluation Program, in *MSIT Workplace*, Effenberg, G. (Ed.), MSI, Materials Science International Services GmbH, Stuttgart, Document ID: 20.12711.1.20 (2002) (Equi. Diagram, Assessment, Crys. Structure, 13)
- [2003Gro] Gröbner, J., Chumak, I., Schmid-Fetzer, R., “Experimental Study of Ternary Ca–Mg–Si Phase Equilibria and Thermodynamic Assessment of Ca–Si and Ca–Mg–Si Systems”, *Intermetallics*, in print (2003)

**Table 1:** Crystallographic Data of Solid Phases


Phase/ Temperature Range [°C]	Pearson Symbol/ Space Group/ Prototype	Lattice Parameters [pm]	Comments/References
( $\beta$ Al)	<i>hP2</i> <i>P6<sub>3</sub>/mmc</i> Mg	$a = 269.3$ $c = 439.8$	at 25°C, 20.5 GPa [Mas2]
( $\alpha$ Al) < 660.452	<i>cF4</i> <i>Fm<math>\bar{3}m</math></i> Cu	$a = 404.96$	at 25°C [Mas2]
( $\gamma$ Ca)	?	?	at 25°C, 1.5 GPa [Mas2]
( $\beta$ Ca) 842-443	<i>cI2</i> <i>Im<math>\bar{3}m</math></i> W	$a = 448.0$	[Mas2]
( $\alpha$ Ca) < 443	<i>cF4</i> <i>Fm<math>\bar{3}m</math></i> Cu	$a = 558.84$	at 25°C [Mas2]
(Si) < 1414	<i>cF8</i> <i>Fd<math>\bar{3}m</math></i> C-diamond	$a = 543.06$	at 25 °C [Mas2]
CaAl <sub>4</sub> (h) 700 - 170	<i>tI10</i> <i>I4/mmm</i> Al <sub>4</sub> Ba	$a = 435.3$ $b = 1107$	[V-C2]
CaAl <sub>4</sub> (l) < 170	<i>m*10</i> ? ?	$a = 615.26 \pm 0.15$ $b = 617.30 \pm 0.13$ $c = 632.90 \pm 0.14$ $\beta = 118.026 \pm 0.016$	[1993Mil]
CaAl <sub>2</sub> < 1086	<i>cF24</i> <i>Fd<math>\bar{3}m</math></i> Cu <sub>2</sub> Mg	$a = 804.0$	[V-C2] Powder X-ray diffraction
Ca <sub>13</sub> Al <sub>14</sub> < 633	<i>mC54</i> <i>C2/m</i> Ca <sub>13</sub> Al <sub>14</sub>	$a = 1555.1 \pm 0.4$ $b = 987.3 \pm 0.2$ $c = 972.6 \pm 0.2$ $\beta = 108.09 \pm 0.02^\circ$	[1998Hua] Single-crystal X-ray diffraction

Phase/ Temperature Range [°C]	Pearson Symbol/ Space Group/ Prototype	Lattice Parameters [pm]	Comments/References
Ca <sub>8</sub> Al <sub>3</sub> < 578	<i>aP</i> 22 <i>P</i> $\bar{1}$ Ca <sub>8</sub> In <sub>3</sub>	$a = 948.4 \pm 0.3$ $b = 959.2 \pm 0.3$ $c = 967.1 \pm 0.3$ $\alpha = 99.02 \pm 0.03^\circ$ $\beta = 101.13 \pm 0.03^\circ$ $\gamma = 119.55 \pm 0.03^\circ$	[1998Hua] Single-crystal X-ray diffraction
CaSi <sub>2</sub> < 1030	<i>hR</i> 18 <i>R</i> $\bar{3}m$ CaSi <sub>2</sub>	$a = 386.3$ $c = 3071.0$	[2000Man]
Ca <sub>14</sub> Si <sub>19</sub> 1085- $\approx$ 900	<i>hR</i> 198 <i>R</i> $\bar{3}c$ Ca <sub>14</sub> Si <sub>19</sub>	$a = 867.2$ $c = 6844.5$	[2000Man]
Ca <sub>3</sub> Si <sub>4</sub> < $\approx$ 910	<i>hP</i> 42 <i>P</i> 6 <sub>3</sub> / <i>m</i> Ca <sub>3</sub> Si <sub>4</sub>	$a = 854.1$ $c = 1490.6$	[2000Man]
CaSi < 1320	<i>oC</i> 8 <i>Cmcm</i> CrB	$a = 459$ $b = 1079.5$ $c = 391$	[2000Man]
Ca <sub>5</sub> Si <sub>3</sub> < 1240	<i>tI</i> 32 <i>I</i> 4/ <i>mcm</i> Cr <sub>5</sub> B <sub>3</sub>	$a = 764.1$ $c = 1487.6$	[2000Man]
Ca <sub>2</sub> Si < 1270	<i>oP</i> 12 <i>Pnma</i> anti-PbCl <sub>2</sub>	$a = 766.7$ $b = 479.9$ $c = 900.2$	[2000Man]
* $\alpha$ , Ca <sub>3</sub> Al <sub>2</sub> Si <sub>2</sub>	<i>oI</i> 14 <i>Immm</i> Ca <sub>3</sub> Al <sub>2</sub> Ge <sub>2</sub>	$a = 400 \pm 1$ $b = 1824 \pm 2$ $c = 457.6 \pm 1.0$	ordered variant of Ta <sub>3</sub> B <sub>4</sub> [1977Wid]
* $\beta$ , CaAl <sub>2</sub> Si <sub>2</sub> < 975	<i>hP</i> 5 <i>P</i> $\bar{3}m1La2O2S$	$a = 413 \pm 1$ $c = 714.5 \pm 1.5$	La <sub>2</sub> O <sub>3</sub> -type superstructure [1967Gla]. Congruent melting [1994Ang]
* $\gamma$ , CaAl <sub>1-x</sub> Si <sub>1+x</sub>	<i>hP</i> 3 AlB <sub>2</sub>	$a = 419.05$ $c = 439.92$	at $x = 0$ [2002Ima]  composition range $x = -0.35$ to $+0.55$ [1967Gla] or $x = -0.4$ to $+0.2$ [2002Lor]
* $\delta$ , Ca <sub>3</sub> Al <sub>6</sub> Si <sub>2</sub> < $\approx$ 1150	<i>hP</i>	$c/a = 1.64$	[1956Chr]
* $\epsilon$ , Ca <sub>2</sub> Al <sub>4</sub> Si <sub>3</sub>	<i>cP</i> 18	$a = 715$	[1956Chr]


**Table 2:** Invariant Equilibria in the Al-rich Corner

Reaction	$T\text{ [}^{\circ}\text{C]}$	Type	Phase	Composition (at.%)		
				Al	Ca	Si
$L + \text{CaAl}_2 \rightleftharpoons \beta + \text{CaAl}_4$	688	U	L	85.4	11.7	2.9
			$\text{CaAl}_2$	66.7	33.3	0
			$\beta$	40	20	40
			$\text{CaAl}_4$	80	20	0
$L \rightleftharpoons (\text{Al}) + \beta$	649	$e_{\text{max}}$	L	97.96	0.65	1.39
$L \rightleftharpoons (\text{Al}) + \beta + \text{CaAl}_4$	610	E	L	94	5.4	0.6
			(Al)	99.99	0	0.01
			$\beta$	40	20	40
			$\text{CaAl}_4$	80	20	0
$L \rightleftharpoons (\text{Al}) + (\text{Si}), \beta$	577	D	L	87.9	0	12.1
			(Al)	98.5	0	1.5
			(Si)	0	0	100
			$\beta$	40	20	40


**Fig. 1: Al–Ca–Si.**  
The pseudobinary section  $\text{CaAl}_2\text{Si}_2$  - Al



**Fig. 2: Al-Ca-Si.**  
The pseudobinary  
section  $\text{CaAl}_2\text{Si}_2$  - Si




**Fig. 3: Al-Ca-Si.**  
Liquidus surface of  
the Al-rich corner  
(>80 at.% Al)





**Fig. 4: Al–Ca–Si.**  
Isothermal section at 400°C; some tie lines above 33 at.% Ca are estimated. Equilibria with the phases  $\alpha$ , CaSi, and  $\text{Ca}_3\text{Si}_4$  are not given



**Fig. 5: Al–Ca–Si.**  
Calculated activities of Ca in the liquid phase at 1350°C and  $x_{\text{Al}} = x_{\text{Si}}$

