

Homework Assignment N°1

Thibault Douzon
Rajavarman Mathivanan

September 4th, 2018

Contents

1	Exercise 1: Geometry	3
1.1	Part a	3
1.2	Part b	4
1.3	Part c	4
1.4	Part d	5
1.5	Part e	5
2	Exercise 2: Calculus	5
2.1	Part a	5
2.2	Part b	6
2.3	Part c	6

1 Exercise 1: Geometry

1.1 Part a

We have:

$$w_0 + 3w_1 + w_2 = 0$$

$$w_0 - w_1 + 2w_2 = 0$$

Normalize by w_2 it gives

$$W_0 + 3W_1 = -1$$

$$W_0 - W_1 = -2$$

Where $W_0 = \frac{w_0}{w_2}$ and $W_1 = \frac{w_1}{w_2}$
Cramer's rule gives us the results:

$$W_0 = \frac{(-1 \times -1) - (3 \times -2)}{(1 \times -1) - (1 \times 3)} = -\frac{7}{4}$$

$$W_1 = \frac{(1 \times -2) - (-1 \times 1)}{(1 \times -1) - (1 \times 3)} = \frac{1}{4}$$

Thus, we can chose $w_2 = 1$ and then

$$w_0 = -\frac{7}{4}$$

$$w_1 = \frac{1}{4}$$


$$w_2 = 1$$

We can check our first system is solved with this solution. Hence the equation of the line is the following:

$$-\frac{7}{4} + \frac{x_1}{4} + x_2 = 0$$


1.2 Part b

Our figure is the following:



1.3 Part c

The point (4,3) is in the positive side of the line, the most simple «proof» is this plot:



It shows our point «above» the blue line, hence it is in the positive side of the line. For this point, because we saw it was on the positive side, $w_0 + w_1x_1 + w_2x_2$ should give a positive result.

$$-\frac{7}{4} + \frac{4}{4} + 3 = \frac{9}{4} > 0$$

1.4 Part d

First we must notice that (w_1, w_2) is orthogonal to our line. To figure out the direction vector of the line, we can rewrite its equation like this

$$-w_1x - w_0 = w_2y$$

Hence its direction vector is $(w_2, -w_1)$ and now we can prove that (w_1, w_2) is orthogonal to the line by showing that

$$(w_1, w_2) \cdot (w_2, -w_1) = 0$$

Now we know the perpendicular direction to the line, we can compute the distance between the point and the line.

$$d_{line}(P) = |[(x_1, x_2) - (3, 1)] \cdot (w_1, w_2)| / \|(w_1, w_2)\|$$

$$d_{line}(P) = |x_1w_1 + x_2w_2 - 3w_1 - 1w_2| / \|(w_1, w_2)\|$$

Or because $w_0 + 3w_1 + 1w_2 = 0$ ($(3, 1)$ is on the line) we can add $w_0 + 3w_1 + 1w_2$ without changing the value. Hence:

$$d_{line}(P) = \frac{|w_0 + x_1w_1 + x_2w_2|}{\|(w_1, w_2)\|}$$

1.5 Part e

We use what we learned from part d:

$$d_{line}((0, 0)) = \frac{|w_0|}{\|(w_1, w_2)\|}$$

Hence we have

$$d_{line}((0, 0)) = \frac{|w_0|}{\sqrt{w_1^2 + w_2^2}} = \frac{\frac{7}{4}}{\sqrt{\frac{1}{4} + 1}} \approx 1.7$$

2 Exercice 2: Calculus

$$\sigma(y) = \frac{1}{1 + e^{-y}}$$

2.1 Part a

Calculus says $\left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2}$, then:

$$\sigma'(y) = \frac{e^{-y}}{(1 + e^{-y})^2}$$

$$\sigma'(y) = \frac{(1 + e^{-y}) - 1}{(1 + e^{-y})^2}$$

$$\sigma'(y) = \sigma(y) \times \frac{(1 + e^{-y}) - 1}{1 + e^{-y}}$$

$$\sigma'(y) = \sigma(y) \times (1 - \sigma(y))$$

2.2 Part b

We know that $\lim_{y \rightarrow \infty} e^{-y} = 0$ thus because the denominator does not tend to zero:

$$\lim_{y \rightarrow \infty} \sigma(y) = \frac{1}{1 + \lim_{y \rightarrow \infty} e^{-y}} = 1$$

and now we know the limit exists and its value:

$$\lim_{y \rightarrow \infty} \sigma'(y) = \lim_{y \rightarrow \infty} \sigma(y) \times (1 - \sigma(y)) = 1 \times (1 - 1) = 0$$

2.3 Part c