Brachistochrone Problem

Rajeev Atla

Physics Club

November 20, 2020

This Section

This Section

- More advanced

This Section

- More advanced
- Goals

This Section

- More advanced
- Goals
    - Get everyone to pass F=ma exam

This Section

- More advanced
- Goals
  - Get everyone to pass F=ma exam
  - USAPhO Qualifiers!!

This Section

- More advanced
- Goals
  - Get everyone to pass F=ma exam
  - USAPhO Qualifiers!!
- Prerequisites (recommended)

This Section

- More advanced
- Goals
  - Get everyone to pass F=ma exam
  - USAPhO Qualifiers!!
- Prerequisites (recommended)
  - Taken/currently taking a physics class

This Section

- More advanced
- Goals
  - Get everyone to pass F=ma exam
  - USAPhO Qualifiers!!
- Prerequisites (recommended)
  - Taken/currently taking a physics class
  - Or...

This Section

- More advanced
- Goals
  - Get everyone to pass F=ma exam
  - USAPhO Qualifiers!!
- Prerequisites (recommended)
  - Taken/currently taking a physics class
  - Or...
  - Willingness to learn

PSA: Problems

- On classroom

PSA: Problems

- On classroom
- Due date: next meeting

PSA: Problems

- On classroom
- Due date: next meeting
- We hope to continue this pattern for the rest of this year

Brachistochrone
What Do I mean?

- Etymology

- Etymology
  - Brachistos ($\beta\rho\alpha\chi\iota\sigma\tau\sigma$) means "shortest"

Brachistochrone

What Do I mean?

- Etymology
  - Brachistos ($\beta\rho\alpha\chi\iota\sigma\tau\sigma$) means "shortest"
  - Chronos ($\chi\rho o\nu o\sigma$) means "time"

Brachistochrone

What Do I mean?

- Etymology
  - Brachistos ($\beta\rho\alpha\chi\iota\sigma\tau\sigma$) means "shortest"
  - Chronos ($\chi\rho o\nu o\sigma$) means "time"
- A brachistochrone curve is the path such that a ball traveling along this path takes the least amount of time

Brachistochrone

What Do I mean?

- Etymology
  - Brachistos ($\beta\rho\alpha\chi\iota\sigma\tau\sigma$) means "shortest"
  - Chronos ($\chi\rho o\nu o\sigma$) means "time"
- A brachistochrone curve is the path such that a ball traveling along this path takes the least amount of time
- This is our problem

- Etymology
  - Brachistos ($\beta\rho\alpha\chi\iota\sigma\tau\sigma$) means "shortest"
  - Chronos ($\chi\rho o\nu o\sigma$) means "time"
- A brachistochrone curve is the path such that a ball traveling along this path takes the least amount of time
- This is our problem
- Formal problem statement

Brachistochrone

What Do I mean?

- Etymology
  - Brachistos ($\beta\rho\alpha\chi\iota\sigma\tau\sigma$) means "shortest"
  - Chronos ($\chi\rho o\nu o\sigma$) means "time"
- A brachistochrone curve is the path such that a ball traveling along this path takes the least amount of time
- This is our problem
- Formal problem statement
  - Constraints: given two points $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$

Brachistochrone

What Do I mean?

- Etymology
  - Brachistos ($\beta\rho\alpha\chi\iota\sigma\tau\sigma$) means "shortest"
  - Chronos ($\chi\rho o\nu o\sigma$) means "time"
- A brachistochrone curve is the path such that a ball traveling along this path takes the least amount of time
- This is our problem
- Formal problem statement
  - Constraints: given two points $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$
  - Find function $y = f(x)$ such that the time it takes for a ball to travel under the influence of gravity from P_1 to P_2

Getting Started

- Let s be a position vector

Getting Started

- Let s be a position vector
- Let v be the associated velocity vector

Getting Started

- Let s be a position vector
- Let v be the associated velocity vector
- From last lecture, recall that

$$v = \frac{ds}{dt} \Rightarrow dt = \frac{ds}{v} \Rightarrow t_{12} = \int_{P_1}^{P_2} \frac{ds}{v}$$

Energy Conservation

Energy Conservation

- Kinetic energy $K = \frac{1}{2}mv^2$

Energy Conservation

- Kinetic energy $K = \frac{1}{2}mv^2$
- Gravitational potential energy $U = mgy$

Energy Conservation

- Kinetic energy $K = \frac{1}{2}mv^2$
- Gravitational potential energy $U = mgy$
- Conservation of energy means that these two are equal

Energy Conservation

- Kinetic energy $K = \frac{1}{2}mv^2$
- Gravitational potential energy $U = mgy$
- Conservation of energy means that these two are equal

$$\frac{1}{2}mv^2 = mgy \Rightarrow v = \sqrt{2gy}$$

Energy Conservation

- Kinetic energy $K = \frac{1}{2}mv^2$
- Gravitational potential energy $U = mgy$
- Conservation of energy means that these two are equal

$$\frac{1}{2}mv^2 = mgy \Rightarrow v = \sqrt{2gy}$$

- We can substitute this into the last equation

Pythagorean Theorem

$$
\begin{aligned}
ds^2 &= dx^2 + dy^2 \\
ds^2 &= dx^2 \left(1 + \left(\frac{dy^2}{dx^2}\right)\right) \\
ds^2 &= dx^2 \left(1 + \left(\frac{dy}{dx}\right)^2\right) \\
ds^2 &= dx^2 \left(1 + y'^2\right) \\
ds &= dx\sqrt{1 + y'^2}
\end{aligned}
$$

Putting It All Together

- Original equation:

$$t_{12} = \int_{P_1}^{P_2} \frac{ds}{v}$$

Putting It All Together

- Original equation:

$$t_{12} = \int_{P_1}^{P_2} \frac{ds}{v}$$

- Conservation of energy:

$$v = \sqrt{2gy}$$

Putting It All Together

- Original equation:

$$t_{12} = \int_{P_1}^{P_2} \frac{ds}{v}$$

- Conservation of energy:

$$v = \sqrt{2gy}$$

- Pythagorean theorem:

$$ds = dx\sqrt{1 + y'^2}$$

Lagrangians

$$t_{12} = \int_{P_1}^{P_2} \sqrt{\frac{1 + y'^2}{2gy}} dx$$

Lagrangians

$$t_{12} = \int_{P_1}^{P_2} \sqrt{\frac{1 + y'^2}{2gy}} dx$$

- We want to minimize this by...

Lagrangians

$$t_{12} = \int_{P_1}^{P_2} \sqrt{\frac{1 + y'^2}{2gy}} dx$$

- We want to minimize this by...
- picking a function $y = f(x)$ to minimize integral

Lagrangians

$$t_{12} = \int_{P_1}^{P_2} \sqrt{\frac{1 + y'^2}{2gy}} dx$$

- We want to minimize this by...
- picking a function $y = f(x)$ to minimize integral
- How do we do it???

Lagrangians

$$t_{12} = \int_{P_1}^{P_2} \sqrt{\frac{1 + y'^2}{2gy}} dx$$

- We want to minimize this by...
- picking a function $y = f(x)$ to minimize integral
- How do we do it???
- Lagrangians

More About Lagrangians

- Let the Lagrangian be

$$\mathcal{L} = \sqrt{\frac{1 + y'^2}{2gy}}$$

More About Lagrangians

- Let the Lagrangian be

$$\mathcal{L} = \sqrt{\frac{1 + y'^2}{2gy}}$$

- Remeber that $y = f(x)$

$$\mathcal{L}(x) = \sqrt{\frac{1 + f'(x)^2}{2gf(x)}}$$

- $f'(x) = \frac{df(x)}{dx}$ (Lagrangian notation)

Least Action Principle

- We need to choose $f(x)$ minimize the time

$$t_{12} = \int_{P_1}^{P_2} \mathcal{L}(x)dx$$

Least Action Principle

- We need to choose $f(x)$ minimize the time

$$t_{12} = \int_{P_1}^{P_2} \mathcal{L}(x)dx$$

- Any ideas?

Least Action Principle

- We need to choose $f(x)$ minimize the time

$$t_{12} = \int_{P_1}^{P_2} \mathcal{L}(x) dx$$

- Any ideas?
- Euler-Lagrange equation

$$\frac{d}{dx}\left(\frac{\partial \mathcal{L}}{\partial f'(x)}\right) = \frac{\partial \mathcal{L}}{\partial f(x)}$$

Partial Derivatives

- Symbol is ∂

Partial Derivatives

- Symbol is ∂
- Hold all other variables constant while taking a derivative

Partial Derivatives

- Symbol is ∂
- Hold all other variables constant while taking a derivative
- Let $f(x, y) = 2x + 3y$, what are $\frac{\partial f(x,y)}{\partial x}$ and $\frac{\partial f(x,y)}{\partial y}$?

Partial Derivatives

- Symbol is ∂
- Hold all other variables constant while taking a derivative
- Let $f(x, y) = 2x + 3y$, what are $\frac{\partial f(x,y)}{\partial x}$ and $\frac{\partial f(x,y)}{\partial y}$?

$$\frac{\partial f(x, y)}{\partial x} = 2$$
$$\frac{\partial f(x, y)}{\partial y} = 3$$

Partial Derivatives

- Symbol is ∂
- Hold all other variables constant while taking a derivative
- Let $f(x, y) = 2x + 3y$, what are $\frac{\partial f(x,y)}{\partial x}$ and $\frac{\partial f(x,y)}{\partial y}$?

$$\frac{\partial f(x,y)}{\partial x} = 2$$
$$\frac{\partial f(x,y)}{\partial y} = 3$$

Beltrami's Identity

$$\frac{d}{dx}\left(\frac{\partial \mathcal{L}}{\partial f'(x)}\right) = \frac{\partial \mathcal{L}}{\partial f(x)}$$

$$\mathcal{L}(x) = \sqrt{\frac{1 + f'(x)^2}{2gf(x)}}$$

Beltrami's Identity

$$\frac{d}{dx}\left(\frac{\partial \mathcal{L}}{\partial f'(x)}\right) = \frac{\partial \mathcal{L}}{\partial f(x)}$$

$$\mathcal{L}(x) = \sqrt{\frac{1 + f'(x)^2}{2gf(x)}}$$

- Anyone want to do this???

Beltrami's Identity

$$\frac{d}{dx}\left(\frac{\partial \mathcal{L}}{\partial f'(x)}\right) = \frac{\partial \mathcal{L}}{\partial f(x)}$$

$$\mathcal{L}(x) = \sqrt{\frac{1 + f'(x)^2}{2gf(x)}}$$

- Anyone want to do this???
- Time for a trick: Beltrami's Identity
  - Notice that $\mathcal{L}(x)$ doesn't *explicitly* depend on x

$$\mathcal{L} - f'(x)\frac{\partial \mathcal{L}}{\partial f'(x)} = C$$

Lagrangians

Using Beltrami's Identity

$$C = \mathcal{L} - f'(x)\frac{\partial \mathcal{L}}{\partial f'(x)}$$

$$\frac{\partial \mathcal{L}}{\partial f'(x)} = \frac{f'(x)}{\sqrt{(2gf(x))\left(1 + f'(x)^2\right)}}$$

$$C = \frac{1}{\sqrt{2gf(x)\left(1 + f'(x)^2\right)}}$$

$$\frac{1}{2gC^2} = f(x)\left(1 + f'(x)^2\right)$$

Lagrangians

Solution

$$\frac{1}{2gC^2} = f(x)\left(1 + f'(x)^2\right)$$

Lagrangians

Solution

$$\frac{1}{2gC^2} = f(x)\left(1 + f'(x)^2\right)$$

- So what's the solution???

Lagrangians

Solution

$$\frac{1}{2gC^2} = f(x)\left(1 + f'(x)^2\right)$$

- So what's the solution???
- It can be shown that this is a cycloid curve

Lagrangians

Solution

$$\frac{1}{2gC^2} = f(x)\left(1 + f'(x)^2\right)$$

- So what's the solution???
- It can be shown that this is a cycloid curve
- Has to be parametrized

$$x(\theta) = \frac{1}{4gC^2}(\theta - \sin\theta)$$

$$y(\theta) = \frac{1}{4gC^2}(1 - \cos\theta)$$

Lagrangians

Solution

$$\frac{1}{2gC^2} = f(x)\left(1 + f'(x)^2\right)$$

- So what's the solution???
- It can be shown that this is a cycloid curve
- Has to be parametrized

$$x(\theta) = \frac{1}{4gC^2}(\theta - \sin\theta)$$
$$y(\theta) = \frac{1}{4gC^2}(1 - \cos\theta)$$

- We can use $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$ to find C

Lagrangian

Visualizing the Solution

$$x(\theta) = \frac{1}{4gC^2}(\theta - \sin\theta)$$
$$y(\theta) = \frac{1}{4gC^2}(1 - \cos\theta)$$

Lagrangian

Visualizing the Solution

$$x(\theta) = \frac{1}{4gC^2}(\theta - \sin\theta)$$
$$y(\theta) = \frac{1}{4gC^2}(1 - \cos\theta)$$



Thank You!