


We just need to
use one of
those Logo
Versions around
60 px by 60 px



- change to Select Medications

- shrink down size
- add more white spare
around elements for
breathing space

center button
- change to Build Report?
- when button is pressed can it change back
to Building Report?

A tre we able to add

a progress bar or

Spinning graphic

to show that the
report is being

Compiled?

For up dialogue

that verifies a

medication has


successfully been

added = Medication

Added =

Table is too wide for mobile & causes honzontal - Just 2 columns to simplify

rane Action
Generic/Brand Add Didd a visually
Strong button



A We need to add

an icon that allows user

to go back to APP

homepage to

start another

report

be changel

A way to return a way to return hack to the main website

■■ Bell Wi-Fi 🗢 6:40 PM

⊕ 4 86% □

PRIVACY 🌣


practices may follow different treatment protocols that vary from published drug monographs. In addition, your child's response will be unique to them and their diagnosis.

Disclaimer of Project Responsibility and Liability

Yut Dem strives to provide general evidence based information and original content for families, mental health, social work, and teaching professionals working with vulnerable families. It does not replace medical advice from a licensed doctor or health professional. The program does not take responsibility for how this information is provided, interpreted and/or enacted upon by other professionals and consumers.

Change to I Seed up Medication Reporter =

YUT DEM :: Medication List All rights reserved.



X We need to add

an icon that allows user

to go back to APP

homepage to

start another

inst

> need one correct logo

A way to return a way to return hack to the main website









• Purpose:

Class: Traditional Mood Stabilizer

Carbamazepine belongs to a group of medications called traditional mood-stabilizers which are effective in treating mood disorders such as bipolar disorder. Carbamazepine is also used for pain control for the treatment of certain types of neurologic pain, as an anticonvulsant to treat seizure disorders (epilepsy) and to treat alcohol withdrawal syndrome.Â

More specifically, Carbamazepine is used for the treatment of the following symptoms:

- o Mania, frenzied, abnormally excited or irritated mood, overly distracted
- o Mixed episodes of mania and depression that happen at the same time
- o Depression, feeling sad, hopeless, guilty, tearful
- Aggression, impulsive or risky behaviourÂ
- Severe PTSDÂ
- Drug and alcohol withdrawal

Carbamazepine works by altering electrical activity in the brain and by decreasing this activity, your child's mood should return to a normal baseline, and stabilize as long as proper dosing is maintained. In turn, your child or teen should feel in better control of their emotions and have an improvement in their overall functioning.Â

Carbamazepine works by altering electrical activity in the brain and by decreasing this activity, your child's mood should return to a normal baseline, and stabilize as long as proper dosing is maintained. In turn, your child or teen should feel in better control of their emotions and have an improvement in their overall functioning.Â

Generally, traditional mood stabilizers are effective agents to stabilize mood and manage overt aggressive and disruptive behaviours. However, these medicines are being used less on young children due to harsh long term side effects on the body compounded with the need for ongoing blood work to monitor plasma levels. Instead, atypical antipsychotics and SSRI antidepressants have become the preferred 1st line agents to help with mood stabilization.Â

Carbamazepine comes in various preparations, so you must discuss with the prescribing healthcare provider if your child doesnâct tolerate swallowing capsules, tablets or prefers a liquid or suspension. Extended-release products are also available and usually given twice daily. Dosing may be increased over several weeks, but your child should see some medication effects within 1-2 weeks.

Dosing:

Carbamazepine comes in various preparations, so you must discuss with the prescribing healthcare provider if your child doesn't tolerate swallowing capsules, tablets or prefers a liquid or suspension. Extended-release products are also available and usually given twice daily. Dosing may be increased over several weeks, but your child should see some medication effects within 1-2 weeks.

Usual dosing according to **Lexicomp** drug database:

Children <6 vr

Please advise
- on HTML email
output, Strange
Symbols are
inserted
- What is the fix?

Connerts:

- All browsers displaying App now
- Rendering time for report is much faster t consistently displays
- PDF + Email HTML rendering is working
- working on both Windows + OSX + 105 platforms

A Please verien app from end User perspective to Optimize ravigation, ensure simplicity, USEasility, and easy understanding of information hiarchez