

```

import numpy as np

def problem_1a (A, B):
    return A + B

def problem_1b (A, B, C):
    return (A * B) + np.transpose(C)

def problem_1c (x, y):
    return x.T.dot(y)

def problem_1d (A, j):
    return np.sum(A[0::2, j])

def problem_1e (A, c, d):
    return np.mean(A[np.nonzero((A >= c) & (A <=d ))])

def problem_1f (x, k, m, s):
    return np.random.randn(x + m, s * np.eye(x.shape[0])), k).T

def problem_1g (A):
    return A[:, np.random.permutation(A.shape[0])]

def problem_1h (x):
    return (x - np.mean(x)) / np.std(x)

def problem_1i (x, k):
    return np.repeat(np.atleast_2d(x), k, axis=0)

def problem_2a ():
    X = np.arange(9).reshape(3,3)
    print(X)
    print("\n")
    row_min = X.min(axis=1)
    print(row_min)
    print("\n")
    print(X-row_min)
    print("\n")

print("Part 2a:\n")
problem_2a()
print("This code is not subtracting each row's minimum element from the respective row because the line 'print(X - row_min)' isn't the correct way of implementing this solution.")
print("The code before this line adequately creates the array X and obtains the minimum value in each row, but the array row_min is not used correctly.")
print("The result of 'X - row_min' is:\n")
print(X-row_min)
print("\nAnd this is because the statement is subtracting the row vector [0, 3, 6] from each row in X.")

```

Part 2a:

```
[[0 1 2]
 [3 4 5]
 [6 7 8]]
```

```
[0 3 6]
```

```
[[ 0 -2 -4]
 [ 3  1 -1]
 [ 6  4  2]]
```

This code is not subtracting each row's minimum element from the respective row because the line 'print(X - row_min)' isn't the correct way of implementing this solution.

The code before this line adequately creates the array X and obtains the minimum value in each row, but the array row_min is not used correctly.

The result of 'X - row_min' is:

```
[[0 1 2]
 [0 1 2]
 [0 1 2]]
```

And this is because the statement is subtracting the row vector [0, 3, 6] from each row in X.

```
def problem_2b():
    X = np.arange(9).reshape(3,3)
    row_min = X.min(axis=1)

    # Part b - change to correctly perform the computation with a 3d array
    Y = np.arange(27).reshape(3,3,3)
    print(Y)
    print("\n")
    # Expand for broadcasting
    # Broadcasting allows this new np array to be used in computations with arrays with different dimensions
    # The row_min array will now subtract 0 from first plane, 3 from second plane, and 6 from third plane of Y
    row_min = row_min[:, np.newaxis, np.newaxis]
    print(Y-row_min)

print("Part 2b:\n")
problem_2b()
print("\nBroadcasting allows this new np array to be used in computations with arrays with different dimensions.")
```

```
print("The row_min array will now subtract 0 from first plane, 3 from second plane, and 6 from third plane of Y.")
```

Part 2b:

```
[[[ 0  1  2]
   [ 3  4  5]
   [ 6  7  8]]
```

```
[[ 9 10 11]
 [12 13 14]
 [15 16 17]]
```

```
[[18 19 20]
 [21 22 23]
 [24 25 26]]]
```

```
[[[ 0  1  2]
   [ 3  4  5]
   [ 6  7  8]]
```

```
[[ 6  7  8]
 [ 9 10 11]
 [12 13 14]]
```

```
[[12 13 14]
 [15 16 17]
 [18 19 20]]]
```

Broadcasting allows this new np array to be used in computations with arrays with different dimensions.

The row_min array will now subtract 0 from first plane, 3 from second plane, and 6 from third plane of Y.

```
def linear_regression (X_tr, y_tr):
    X_t = X_tr.T
    Xdoty = np.matmul(X_t, y_tr)
    XdotXT = np.matmul(X_t, X_tr)
    return np.linalg.solve(XdotXT, Xdoty)

def train_age_regressor ():
    # Load data
    X_tr = np.reshape(np.load("age_regression_Xtr.npy"), (-1, 48*48))
    y_tr = np.load("age_regression_ytr.npy")
    X_te = np.reshape(np.load("age_regression_Xte.npy"), (-1, 48*48))
    y_te = np.load("age_regression_yte.npy")

    w = linear_regression(X_tr, y_tr)
    yhat_tr = np.matmul(X_tr, w)
    yhat_te = np.matmul(X_te, w)
```

```

# Calculate MSE for training data
trainDifference = np.square(yhat_tr - y_tr)
trainMSE = np.mean(trainDifference)

# Calculate MSE for testing data
testDifference = np.square(yhat_te - y_te)
testMSE = np.mean(testDifference)

return trainMSE, testMSE

print("Part 3:\n")
print(train_age_regressor())
Part 3:
(80.83988427156137, 749.3051827446681)

import matplotlib.pyplot as plt
from scipy.stats import poisson

def poisson_distribution():
    data = np.load('PoissonX.npy')
    plt.hist(data, density=True, edgecolor='blue')
    plt.show

    rateParameters = [2.5, 3.1, 3.7, 4.3]
    figure, axis = plt.subplots(2,2)

    Y = poisson.pmf(data,rateParameters[0])
    axis[0, 0].plot(data,Y)
    axis[0, 0].set_title("Rate Parameter of 2.5")

    Y = poisson.pmf(data,rateParameters[1])
    axis[0, 1].plot(data,Y)
    axis[0, 1].set_title("Rate Parameter of 3.1")

    Y = poisson.pmf(data,rateParameters[2])
    axis[1, 0].plot(data,Y)
    axis[1, 0].set_title("Rate Parameter of 3.7")


    Y = poisson.pmf(data,rateParameters[3])
    axis[1, 1].plot(data,Y)
    axis[1, 1].set_title("Rate Parameter of 4.3")


    plt.subplots_adjust(bottom=0.1, right=1.1, top=1.1)
    plt.show

print("Part 4a:\n")
poisson_distribution()

```

Part 4a:





```
print("Part 4b:\n")
print("I.")
print("It appears that the smaller magnitudes of x tend to correspond
to the larger values of y.")
print("This can be seen in the first histogram where larger values of
y match up with smaller values of x.")
print("The Poisson distribution graphs also show this with the highest
y values corresponding to values of x between 2.5 and 5.0.\n")
print("II.")
print("It appears that the extreme larger and extreme lower magnitudes
of x exhibit more uncertainty when it comes to predicting the value of
y.")
print("This can be seen in the four histograms of the Poisson
distribution where the y values are the lowest for the highest and
lowest values of x.")
```

Part 4b:

I.

It appears that the smaller magnitudes of x tend to correspond to the

larger values of y .

This can be seen in the first histogram where larger values of y match up with smaller values of x .

The Poisson distribution graphs also show this with the highest y values corresponding to values of x between 2.5 and 5.0.

II.

It appears that the extreme larger and extreme lower magnitudes of x exhibit more uncertainty when it comes to predicting the value of y .

This can be seen in the four histograms of the Poisson distribution where the y values are the lowest for the highest and lowest values of x .