

①

①

Set theory:

(i) Set: A set is a collection of objects under study, sets are usually denoted by capital letters A, B, C, X, Y, Z, etc.

For EX: Students of electrical engineering in a college, rivers of India, states of India etc.

(ii) Experiment: An experiment is a physical process that is observed and whose result is noted.

Ex: ... all throwing a dice, etc.

For Ex: turning a switch on or off, throwing a dice, etc.

Experiments are of two types

- (a) deterministic
- (b) non-deterministic experiment

(a) Deterministic experiment:

If we are sure about the outcome of the experiment before conducting the experiment, then it is known as deterministic experiment. For Ex: constructing a wall, throwing a stone upwards, etc.

(b) Non-Deterministic Experiment:

An experiment is called a non-deterministic, random probabilistic or stochastic if we are not sure that which of the possible outcomes will occur when an experiment is conducted. Here after an experiment means random experiment.

for EX: tossing a coin, throwing a dice, etc.

(iii) Trial? A single performance of an experiment is called a trial.

For Ex: tossing a coin.

Scanned with CamScanner

(iv) Sample space or sample point:

A collection of all possible outcomes of a random experiment is called a sample space and usually denoted by S . The elements of a sample space are called sample points.

For Ex: If a dice is thrown, then any one of 1, 2, 3, 4, 5 or 6, will appear on the face.

$$S = \{1, 2, 3, 4, 5, 6\}$$

↓
sample points of S .



Smita Sonker is presenting



Swati Meena
and 53 more



2:35 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

Find

2 / 18 162%

(v) Discrete sample space:

A sample space is called to be discrete, if it has finitely many or a countably infinite number of elements.

For EX: tossing a coin the sample space

$$S = \{ H, T \}$$

throwing a dice

$$S = \{ 1, 2, 3, 4, 5, 6 \}$$

etc.

Meeting details ^



Turn on captions



Smita Sonker
is presenting





Smita Sonker is presenting



aditya jain
and 53 more



2:35 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

2 / 18 162% Find

(vi) continuous Sample Space
A sample space is said to be continuous; if the elements of the sample space constitute a continuum
for Ex: All the points on a line segment.

(vii) Event: Any subset A of a sample space S and each is called an event.
In tossing a coin $S = \{H, T\}$
Here \emptyset and S & $\{H, T\}$ are events and each of

Meeting details ^



Turn on captions

Up arrow icon
Smita Sonker
is presenting





Smita Sonker is presenting



neelam
and 37 more



2:36 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

Print Preview Back Forward Find

Here \emptyset , $\{H\}$, $\{T\}$, $\{H, T\}$ are subsets of S and each of them is an event.

Scanned with CamScanner

Ex. 3 simple or Elementary event

If an event has only one sample point and cannot be further divided into smaller events then it is known as simple or elementary event.

Prahlad Kumar has left the meeting

For Ex : 1, 2, 3, 4, 5, 6

Meeting details ^



Turn on captions

Up arrow icon
Smita Sonker
is presenting

⋮



Smita Sonker is presenting



saloni vij
and 34 more



2:37 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help



3 / 18

162%

Find

⑥ compound event:

If an event has more than one sample point and can be obtained by combining the several elementary events is called a compound event.

for Ex:

when a coin is tossed twice then sample space

$$S = \{HH, HT, TH, TT\}$$

and each of S is a compound event.

hariom dagar has left the meeting

Meeting details ^



Turn on captions



Smita Sonker
is presenting





Smita Sonker is presenting



suman saurav
and 33 more



2:37 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help



3 / 18

162%

Find

(viii) favourable Event:

(ix) Equally Likely events : All six faces are equally likely to occur.

(x) complement of an Event \bar{A} or A' = $S - A$

(xi) Mutually Exclusive Events
 $A \cap B = \emptyset$

(xii) Exhaustive events:

The total number of possible outcomes in any of a random experiment is called. exhaustive

Tushar has left the meeting

Meeting details ^



Turn on captions



Smita Sonker
is presenting





Smita Sonker is presenting



Bomma Sahithya
and 33 more



2:37 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

Print Email Find

events.

(xiii) odd in favour of an event and odd against an event

Let there be m outcomes favourable to a certain event and n outcomes are not in favour to the event in a sample space S , then odd in favour of the event
odd not in favour or against of the event $= \frac{m}{m+n}$.

Scanned with CamScanner

(xiv) Permutation

$${}^n P_r = \frac{m!}{(m-r)!}$$

Combination

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting





Smita Sonker is presenting



Sahil Shende
and 32 more



2:38 PM



You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

Printer icon | Print icon | Up arrow | Down arrow | Page number (4) / 18 | Magnifying glass | Find | 162% |

(xv) combination

$${}^m_{\text{C}_r} \text{ or } \binom{m}{r} = \frac{\cancel{m}}{\cancel{r} \ m-r}$$

(xvi) union of Events :
Let A and B be two events, then A union B ($A \cup B$)
is the event that consists all the points either in A or in
B or in both A and B.

(xvii) Intersection of Events :
A intersection B ($A \cap B$)

Meeting details ^



Turn on captions



Smita Sonker
is presenting





Smita Sonker is presenting



Vipul Bhakar
and 28 more



2:38 PM



(xvii) Intersection of Events :

let A and B be two events then A intersect B ($A \cap B$)
is the set of all those points that are contained in
both A and B.

classical.

Defⁿ of Probability:

$$P(A) = \frac{m}{n} = \frac{\text{no. of favourable outcomes}}{\text{Total no. of outcomes}}$$

Hitesh Chaudhary has left the meeting

$m = \text{no. of favourable outcomes do A.}$

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting

⋮



Smita Sonker is presenting



aryan kabra
and 25 more



2:38 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

Print Preview Back Forward Page 4 / 18 162% Find

$m = \text{no. of favourable outcomes to } A.$

$\therefore n - m = \text{non-favourable no. of outcomes to } A$
and set of non-favourable no. of outcomes is
denoted by A' $\Rightarrow P(A') = \frac{n-m}{n} = 1 - \frac{m}{n} = 1 - P(A)$

$$\Rightarrow P(A) + P(A') = 1$$

If $P(A) = \text{prob. of success of an event } A = p$ and

$P(A') = \dots, \text{failure}, \dots, A = q.$

$$p = P(A) = \frac{m}{n}, 0 \leq p \leq 1$$

mayank kumar has left the meeting



Dhanavath jayanthnaik
joined

Meeting details ^



Turn on captions

Smita Sonker
is presenting





Smita Sonker is presenting



mohit meena
and 25 more



2:39 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

5 / 18 162% Find

$\Rightarrow p + q = 1$ $q = P(A') = \frac{n-m}{n} = 1 - \frac{m}{n} = 1-p$, $0 \leq q \leq 1$

Scanned with CamScanner

(5)

Empirical or statistical defⁿ of probability.

Let m = frequency of occurrence of event A.

n = number of independent trials of a random experiment which are repeated under the same conditions.

urvashi mittal has left the meeting

lim $\frac{m}{n}$,

Meeting details ^



Turn on captions

Up arrow icon
Smita Sonker
is presenting

⋮



Smita Sonker is presenting



prathmesh dakh...
and 23 more




2:39 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help



5 / 18

162%

Find

$$\text{Then } P(A) = \lim_{n \rightarrow \infty} \frac{m}{n},$$

provided limit is unique and finite.

where $\frac{m}{n}$ = relative frequency of the event A in
 n trials.

If $n \rightarrow \infty$ then $\frac{m}{n}$ is very close to actual probability.
This def" of prob. is also known as relative frequency
def" of prob.

suman saurav has left the meeting

To records show that 800 of 1000 tested ceramic

Meeting details ^



Turn on captions



Smita Sonker
is presenting





Smita Sonker is presenting



Amisha
and 23 more



2:39 PM



set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

Find

Ex: If records show that 800 of 1000 tested ceramic insulators were able to withstand a certain thermal shock. Find the prob. that any one untested insulator will be able to withstand the thermal shock.

Sol: Here m = 800, n = 1000. Let A be the event that any untested insulator will be able to withstand the thermal shock.

$$P(A) = \frac{800}{1000} = 0.80$$

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting

⋮

6 (6) WhatsApp x | Meet - jpb-gfxi-jas x | Meet - nvo-hwuy-usf x | New Tab x | +

← → C 🔒 meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

nemali girisai and 16 more

19 5 2:40 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

5 / 18 162% Find

EX-9 : An electric engineer is studying the

Scanned with CamScanner

(iii) Axiomatic Def^m of Probability
If s is the sample space and A be any event of

Meeting details ^

Meeting controls: microphone, video, red circle button

CC Turn on captions

Smita Sonker is presenting

Type here to search

Windows Start button

14:40 11-09-2020

Smita Sonker is presenting

r ramesh vishnoi and 15 more 2:41 PM You

(iii) Axiomatic Defⁿ of Probability

If s is the sample space and A be any event of random experiment, then

- (a) $0 \leq P(A) \leq 1$, for each event $A \in S$
- (b) $P(S) = 1$
- (c) If A_1 and A_2 are two mutually exclusive events in S , then $P(A_1 \cup A_2) = P(A_1) + P(A_2)$

(a), (b) & (c) are known as axioms of Probability.

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting

...



Type here to search



14:41
11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting Dhanavath jayanth... and 13 more 16 people 5 messages 2:41 PM You

Ex-10: Mr. X who is a broker feels that the prob that a given stock will go up in value during the day's trading is 0.6 and the prob. that it will go down in value is 0.1. What is the prob. that it will go up or down?

Sol: $P(A) = 0.6$, $P(B) = 0.1$

$\therefore P(A \cup B) = P(A) + P(B) = 0.6 + 0.1 = 0.7$

Bomma Sahithya has left the meeting

generalization of third Axioms of prob.

Meeting details ^



CC

Turn on captions



Smita Sonker
is presenting



Type here to search



14:41
11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

M Mannat Jain and 12 more

15 5 2:41 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

6 / 18 162% Find

$\therefore P(A \cup B) = P(A) + P(B) = 0.6 + 0.1 = 0.7$

Generalization of third Axiom of prob.

If A_1, A_2, \dots, A_n are mutually exclusive events in a sample space S , then

$P(A_1 \cup A_2 \cup A_3 \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$

$P\left(\bigcup_{i=1}^n A_i\right) = \sum_{i=1}^n P(A_i)$

Vipul Bhakar has left the meeting

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting



Type here to search



14:41 11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

R Raman Jain and 12 more

15 5 2:41 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

7 / 18 162% Find

Conditional Law of Probability or Theorem of Total Probability.

Theorem 6.4.1:

Let A and B be the mutually exclusive events, then prove that $P(A \cup B) = P(A) + P(B)$

Proof: Let n be the total number of equally likely cases out of which m_1 be the number that event A occurs and m_2 be the number that event B occurs

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting

...



Type here to search



14:41
11-09-2020
ENG
10

6 (6) WhatsApp x Meet - jpb-gfxi-jas x Meet - nvo-hwuy-usf x New Tab x +

← → C 🔒 meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting a arnav goel and 10 more 13 5 2:41 PM You ...

set theory.pdf - Adobe Reader File Edit View Document Tools Window Help 7 / 18 162% Find

A occurs and m_2 be the number that event B occurs then the number of favourable ones of occurring A or B = $m_1 + m_2$

$\therefore P(A \text{ or } B) = P(A \cup B) = \frac{m_1 + m_2}{n} = \frac{m_1}{n} + \frac{m_2}{n}$.

but $P(A) = \frac{m_1}{n}$, $P(B) = \frac{m_2}{n}$.

$\therefore P(A \cup B) = P(A) + P(B)$

Theorem 6.4.2 :-

Dhanavath jayanthnaik has left the meeting

Let A and B be two events then show that

Meeting details ^



CC Turn on captions

Up Smita Sonker is presenting



Type here to search



14:41 11-09-2020




Smita Sonker is presenting



prathmesh dakh...
and 10 more




2:42 PM



Theorem 6.4.2 :-

Let A and B be two events then show that

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$



Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting

⋮



Type here to search



14:42
11-09-2020
ENG
10

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

Harshit Mehra and 9 more

12 5 2:42 PM You

Boof: using Venn diagram

$A \cup B = A \cup (\bar{A} \cap B)$

where A and $\bar{A} \cap B$ are mutually exclusive events.

$\therefore P(A \cup B) = P(A) + P(\bar{A} \cap B)$

$= P(A) + P(\bar{A} \cap B) + P(A \cap B) - P(A \cap B)$

$= P(A) + P(B) - P(A \cap B)$

$\because (\bar{A} \cap B)$ and $(A \cap B)$ are mutually exclusive events

Meeting details ^

CC Turn on captions

Smita Sonker is presenting

Type here to search

14:42 11-09-2020

Smita Sonker is presenting

Krishnam Goyal
and 9 more

2:42 PM You

$$= P(A) + P(B) - P(A \cap B)$$

$$\because (\bar{A} \cap B) \text{ and } (A \cap B) \text{ are mutually exclusive events}$$

$$\text{and } B = (A \cap B) \cup (\bar{A} \cap B)$$

$$\therefore P(B) = P(A \cap B) + P(\bar{A} \cap B)$$

$$\Rightarrow P(A \cup B) = P(A) + P(B) - P(A \cap B)$$
 Hence Proved

Remark: If A and B are mutually exclusive events,
then $P(A \cap B) = 0$

Meeting details ^



cc

Turn on captions

Smita Sonker
is presenting



Type here to search



14:42 11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting Sahil Shende and 9 more 2:42 PM

set theory.pdf - Adobe Reader

$P(A \cap B) = 0$

and $P(A \cup B) = P(A) + P(B)$

Theorem 6.4.3:

If A, B and C are any three events, then

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$$

Proof: $P(A \cup B \cup C) = P[A \cup (B \cup C)] = P[A] + P[B \cup C] - P(A \cap B \cap C)$

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting



Type here to search



14:42
11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

Shubham Bansal and 9 more

12 5 2:42 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

8 / 18 162% Find

Boof: $P(A \cup B \cup C) = P[A \cup (B \cup C)] = P[A] + P(B \cup C) - P(A \cap B \cap C)$

using the - 6.4.2

$= P(A) + P(B) + P(C) - P(B \cap C) - [P(A \cap B) \cup P(A \cap C)]$

$= P(A) + P(B) + P(C) - P(B \cap C) - [P(A \cap B) + P(A \cap C) - P(A \cap B \cap C)]$

$= P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$

Hence Proved

Scanned with CamScanner

m 6.4.1..

Meeting details ^

Meeting controls: microphone, video, share screen

CC Turn on captions

Smita Sonker is presenting

Type here to search

Windows Start button

14:42 11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

arnav goel and 9 more

12 5 2:42 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

9 / 18 162% Find

Scanned with CamScanner

m 6.1.4 :

more than

Boof : $P(\emptyset) = 0$

$s \cup \emptyset = s$ (s and \emptyset are mutually exclusive events)

$\therefore P(s \cup \emptyset) = P(s)$

$\Rightarrow P(s) + P(\emptyset) = P(s)$

$\Rightarrow P(\emptyset) = 0$

Th...

Meeting details ^

Turn on captions

Smita Sonker is presenting



Type here to search



14:42 11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting r ramesh vishnoi and 9 more 12 people 5 messages 2:42 PM

Theorem 6-4-5 Show that $P(A') = 1 - P(A)$

Proof: $A \cup A' = S$

(A and A' are mutually exclusive events).

$\therefore P(A \cup A') = P(A) + P(A') = P(S) = 1$ ($\because P(S) = 1$)

$\Rightarrow P(A') = 1 - P(A)$

Hence Proved

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting

...



Type here to search



14:42
11-09-2020

6 (6) WhatsApp x Meet - jpb-gfxi-jas x Meet - nvo-hwuy-usf x New Tab x +

← → C https://meet.google.com/nvo-hwuy-usf 2:43 PM You

Smita Sonker is presenting

Harshit Mehra and 9 more

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

9 / 18 162% Find

Ex-13 A single dice is thrown once. Find the prob. of getting a 3 or 5.
 $S = \{1, 2, 3, 4, 5, 6\}$

Sol: $P(1) = P(2) = \dots = P(6) = \frac{1}{6}$

$P(3 \text{ or } 5) = P(3) + P(5) = \frac{1}{6} + \frac{1}{6} = \frac{1}{3}$ (all are mutually exclusive events)

Ex-15 In throwing a pair of dice, find the prob. of getting 3, 5 or 11.

Sol: $P(A) = \frac{2}{36}, P(B) = \frac{4}{36}, P(C) = \frac{2}{36}$

Meeting details ^



CC
Turn on captions

Smita Sonker
is presenting



Type here to search



14:43 11-09-2020

Smita Sonker is presenting

M Mannat Jain and 9 more

2:43 PM You

Ex-13 A single dice is thrown once. Find the prob. of getting a 3 or 5.
 $S = \{1, 2, 3, 4, 5, 6\}$

Sol: $P(1) = P(2) = \dots = P(6) = \frac{1}{6}$

$P(3 \text{ or } 5) = P(3) + P(5) = \frac{1}{6} + \frac{1}{6} = \frac{1}{3}$ (all are mutually exclusive events)

Ex-15 In throwing a pair of dice, find the prob. of getting 3, 5 or 11.

Sol: $P(A) = \frac{2}{36}, P(B) = \frac{4}{36}, P(C) = \frac{2}{36}$

Meeting details ^



CC
Turn on captions

Smita Sonker
is presenting



Type here to search



14:43
11-09-2020

Smita Sonker is presenting S Sahil Shende and 8 more 2:43 PM You

~~Ques - 18~~ In throwing a pair of dice, find the prob
of getting 3, 5 or 11

Solⁿ: $P(A) = \frac{2}{36}$, $P(B) = \frac{4}{36}$, $P(C) = \frac{2}{36}$
 $P(A \cup B \cup C) = P(A) + P(B) + P(C)$ ($\because A, B$ and C are mutually
 $= \frac{2}{36} + \frac{4}{36} + \frac{2}{36} = \frac{2}{9}$ exclusive events)

Scanned with CamScanner

Raman Jain has left the meeting

conditional Probability:

Meeting details ^



CC Turn on captions

Up Smita Sonker is presenting



Type here to search



14:43 11-09-2020

6 (6) WhatsApp x Meet - jpb-gfxi-jas x Meet - nvo-hwuy-usf x New Tab x +

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting Smita Sonker and 8 more 2:43 PM You

set theory.pdf - Adobe Reader File Edit View Document Tools Window Help 10 / 18 162% Find

conditional Probability:
In our daily life, it is not always possible that two or more events always happen independently but their occurrence may depend on one another, i.e., the probability of one event depends on the happening of another event.

Defⁿ:
Let A and B be two events, then conditional probability of A given B is

$$= P(A/B) = \frac{P(A \cap B)}{P(B)}, P(B) > 0.$$

similarly Probability of B given A = $P(B/A) = \frac{P(A \cap B)}{P(A)}$, $P(A) > 0$

Meeting details ^



CC
Turn on captions

Up
Smita Sonker
is presenting

...



Type here to search



14:43
11-09-2020
ENG
10

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

p prathmesh dakh... and 8 more

11 5 2:43 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

10 / 18 162% Find

6.5.1 Theorem on Compound Probability of multiplication,
Law of Probability

If A and B are any two events in sample spaces,
then

$$P(A \cap B) = P(A \cdot B) = P(A) \cdot P(B/A), \text{ if } P(A) > 0$$
$$= P(B) \cdot P(A/B), \text{ if } P(B) > 0$$

Similarly let A, B and C be three events, then

$$P(A \cap B \cap C) = P(A) \cdot P(B/A) \cdot P(C/B)$$

Meeting details ^



cc

Turn on captions

Smita Sonker
is presenting



Type here to search



14:43 11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

ramesh vishnoi and 8 more

5 2:43 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

10 / 18 162% Find

6.5.1 Theorem on Compound Probability of Multiplication
Law of Probability

If A and B are any two events in sample spaces,
then

$$P(A \cap B) = P(AB) = P(A) \cdot P(B/A), \text{ if } P(A) > 0$$
$$= P(B) \cdot P(A/B), \text{ if } P(B) > 0$$

Similarly let A, B and C be three events, then

$$P(A \cap B \cap C) = P(A) \cdot P(B/A) \cdot P(C/A \cap B),$$

Meeting details ^



cc

Turn on captions

Smita Sonker
is presenting

...



Type here to search



14:43 11-09-2020

Smita Sonker is presenting

Krishnam Goyal
and 8 more

2:43 PM You

set theory.pdf - Adobe Reader

Independent Events:

Two events are said to be independent if happening or non-happening of the events do not depend on each-other.

If A and B are two events, then A and B are said to be independent if

$P(A|B) = P(A)$ and $P(B|A) = P(B)$

or $P(AB) = P(A \cap B) = P(A) \cdot P(B)$

This is called special multiplication rule for independent events.

Meeting details ^

Turn on captions

Smita Sonker is presenting

Type here to search



Type here to search



14:43
11-09-2020
ENG
10

Smita Sonker is presenting

Sahil Shende and 8 more

2:44 PM You

Remark: Therefore, we can say that two events A and B are independent, if and only if

(i) $P(A|B) = P(A)$, (ii) $P(B|A) = P(B)$ and (iii) $P(A \cap B) = P(A)P(B)$

and if $P(A \cap B) \neq P(A) \cdot P(B)$, then A and B are not independent events.

(iii) can be extended: for more than two events.

Defⁿ: Let $E_1, E_2, E_3, \dots, E_n$ be n events, then E_1, E_2, \dots, E_n

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting



Type here to search



14:44
11-09-2020
ENG
10

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

Harshit Mehra and 8 more

11 5 2:44 PM You

set theory.pdf - Adobe Reader

Defⁿ: Let $E_1, E_2, E_3, \dots, E_n$ be n events, then E_1, E_2, \dots, E_n are independent if and only if

$$P(E_1 \cap E_2 \cap \dots \cap E_n) = P(E_1) \cdot P(E_2) \cdot P(E_3) \dots \cdot P(E_n)$$

Meeting details ^

Turn on captions

Smita Sonker is presenting



Type here to search



14:44 11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting Sahil Shende and 7 more 2:44 PM

set theory.pdf - Adobe Reader

Theorem :

C If A and B are independent events, then show that

- (a) \bar{A} and \bar{B} are independent events.
- (b) \bar{A} and B are independent events.
- (c) A and \bar{B} are independent events.

Proof:

(a) $\overline{A \cap B} = \bar{A} \cup \bar{B}$

$\Rightarrow P(\overline{A \cap B}) = P(\bar{A} \cup \bar{B}) = 1 - P(A \cup B)$ [using result
 $P(\bar{A}) = 1 - P(A)$]

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting

...



Type here to search



14:44
11-09-2020

Smita Sonker is presenting

Smita Sonker and 7 more

2:44 PM You

$P(\bar{A}) = 1 - P(A)$

$$\begin{aligned} \Rightarrow P(\bar{A} \cap \bar{B}) &= 1 - [P(A) + P(B) - P(A \cap B)] \\ &= 1 - [P(A) + P(B) - P(A) \cdot P(B)] \\ &= [1 - P(A)] - P(B)[1 - P(A)] \quad (\because A \text{ & } B \text{ are independent events}) \\ &= [1 - P(A)][1 - P(B)] \\ &= P(\bar{A}) \cdot P(\bar{B}) \end{aligned}$$

Hence result -

(b) $B = (A \cap B) \cup (\bar{A} \cap B)$

Meeting details ^



CC

Turn on captions

Smita Sonker
is presenting



Type here to search



14:44
11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

rameshvishnoi... and 7 more

10 5 2:45 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

12 / 18 162% Find

(b) $B = (A \cap B) \cup (\bar{A} \cap B)$

$A \cap B$ and $\bar{A} \cap B$ are mutually exclusive

$\therefore P(B) = P(A \cap B) + P(\bar{A} \cap B)$

$P(B) = P(A) \cdot P(B) + P(\bar{A} \cap B)$

$\Rightarrow P(\bar{A} \cap B) = P(B) [1 - P(A)] = P(B) P(\bar{A})$

$\therefore \bar{A}$ and B are independent events.

Meeting details ^



CC
Turn on captions

Up
Smita Sonker
is presenting



Type here to search



14:45
11-09-2020

Smita Sonker is presenting

Sahil Shende and 5 more

2:45 PM You

set theory.pdf - Adobe Reader

13 / 18 162% Find

Mark

13

Term 'Independent' is defined in terms of 'probability' of events whereas 'mutually exclusive' is defined in terms of 'events'. Moreover, mutually exclusive events never have an outcome You can't unmute someone else's presentation but independent events do have common outcomes, provided each event is non-empty. Clearly 'independent' and 'mutually exclusive' do not have the same meaning.

You can't unmute someone else's presentation

Meeting details ^



CC
Turn on captions


Up arrow
Smita Sonker
is presenting



Type here to search



14:45
11-09-2020
ENG
10



Meeting details ^



CC
Turn on captions

▲
Smita Sonker
is presenting



Type here to search



14:45
11-09-2020
ENG
10

Smita Sonker is presenting

Mannat Jain and 5 more

2:45 PM You

Ex-17: A dice is rolled. If the outcome is an even number what is the probability that it is prime?

Sol: In rolling a dice, the sample space

$$S = \{1, 2, 3, 4, 5, 6\}.$$

Let A be the event that the number is even

$$= \{2, 4, 6\}$$

and B be the event that number is prime

$$= \{2, 3, 5\}.$$

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting



Type here to search



14:45
11-09-2020

Smita Sonker is presenting

Smita Sonker and 5 more

2:45 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

13 / 18 162% Find

Then $A \cap B = \{2\} \Rightarrow P(A \cap B) = \frac{1}{6}$

$A' = \{2, 3, 5\}$

$P(A') = \frac{3}{6} = \frac{1}{2}$

$\therefore P(\text{getting a prime number} / \text{getting an even number})$

$= \frac{P(A \cap B)}{P(A')} = \frac{\frac{1}{6}}{\frac{1}{2}} = \frac{1}{3}$

Meeting details ^



cc

Turn on captions

Smita Sonker
is presenting

...



Type here to search



14:45 11-09-2020



Smita Sonker is presenting



ramesh vishnoi
and 5 more



2:46 PM



Ex-18 A family has two children. What is the probability that both are girls, given that at least one of them is a girl?

Solⁿ The sample space $S = \{(b, b), (b, g), (g, b), (g, g)\}$

where $b = \text{boy}$, $g = \text{girl}$.

Let A and B be the events that both are girls and at least one is girl respectively. Then.

$$\therefore P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{1}{4}}{\frac{3}{4}} = \frac{1}{3}$$

Meeting details ^



CC
Turn on captions

Up
Smita Sonker
is presenting



Type here to search



14:46
11-09-2020

6 (6) WhatsApp X Meet - jpb-gfxi-jas X Meet - nvo-hwuy-usf X New Tab X +

← → C https://meet.google.com/nvo-hwuy-usf 2:46 PM You

Smita Sonker is presenting

Sahil Shende and 5 more

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

14 / 18 162% Find

Ex-18 A family has two children. What is the probability that both are girls, given that at least one of them is a girl?

Solⁿ The sample space $S = \{(b, b), (b, g), (g, b), (g, g)\}$

where $b = \text{boy}$, $g = \text{girl}$.

Let A and B be the events that both are girls and at least one is girl respectively. Then.

$\therefore P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{1/4}{3/4} = \frac{1}{3}$

Meeting details ^



CC

Turn on captions

Smita Sonker
is presenting

...



Type here to search



14:46 11-09-2020

Smita Sonker is presenting

prathmesh dakh... and 5 more

2:46 PM You

set theory.pdf - Adobe Reader

Extension of Multiplication law of Probability to n Events

(15 → 13)

Let A_1, A_2, \dots, A_n be n events, then.

$P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1) \cdot P(A_2 | A_1) \cdot P(A_3 | A_1 \cap A_2) \dots P(A_n | A_1 \cap A_2 \cap \dots \cap A_{n-1})$

where $P(A_i | A_1 \cap A_2 \cap \dots \cap A_{i-1})$

be the conditional probability of the event A_i given that the event A_1, A_2, \dots, A_{i-1} have already happened.

(i) Extension of Multiplication Law of Probability for n-independent Events

Meeting details ^



cc

Turn on captions

Smita Sonker
is presenting

...



Type here to search



14:46
11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

M Mannat Jain and 5 more

5 2:46 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

15 / 18 162% Find

(1) Extension of Multiplication Law of Probability for n -independent Events -

Let A_1, A_2, \dots, A_n be n -independent events then

$$P(A_1 \cap A_2 \cap A_3 \cap \dots \cap A_n) = P(A_1) \cdot P(A_2) \cdot \dots \cdot P(A_n)$$

The above result can be proved using the concept of independence; if A_1, A_2, \dots, A_n are independent events then

$$\begin{aligned} P(A_2 | A_1) &= P(A_2) \cdot P(A_3 | A_1 \cap A_2) = P(A_3), \dots, P(A_n | A_1 \cap A_2 \cap \dots \cap A_{n-1}) \\ &= P(A_n) \end{aligned}$$

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting



Type here to search



14:46
11-09-2020

Smita Sonker is presenting

Smita Sonker and 5 more 2:46 PM You

$= P(A_m)$

$\Rightarrow P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1) \cdot P(A_2) \cdots P(A_n).$

The above result can be proved using the concept of independence, if A_1, A_2, \dots, A_n are independent events then

$P(A_2 | A_1) = P(A_2), P(A_3 | A_1 \cap A_2) = P(A_3), \dots,$

$P(A_m | A_1 \cap A_2 \cap \dots \cap A_{n-1}) = P(A_m)$

$\Rightarrow P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1) \cdot P(A_2) \cdots P(A_n)$

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting



Type here to search



14:46
11-09-2020

6 (6) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

r ramesh vishnoi and 4 more

5 2:47 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

16 / 18 162% Find

1 pairwise Independent Events.

Let A_1, A_2, \dots, A_n defined on the same space .

$P(A_i) > 0 ; i = 1, 2, \dots, n$, then these events are said to be pairwise independent, if every pair of two events is independent, i.e .

$P(A_i \cap A_j) = P(A_i) \cdot P(A_j) , i \neq j = 1, 2, \dots, n$

Harshit Mehra has left the meeting

Meeting details ^

Turn on captions

Smita Sonker is presenting



Type here to search



14:47 11-09-2020 ENG



10

Smita Sonker is presenting

Sahil Shende and 4 more

2:47 PM You

(iii) Mutually Independent Events :

If A_1, A_2, \dots, A_n be n events in a sample space S , then they are said to be mutually independent if

$$P(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}) = P(A_{i_1}) P(A_{i_2}) \dots P(A_{i_k})$$

Hence, the events are mutually independent if they are independent by pairs; by triplets, and by quadruplets and so on.

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting



Type here to search



14:47
11-09-2020
ENG
10

(7) WhatsApp x Meet - jpb-gfxi-jas x Meet - nvo-hwuy-usf x New Tab x +

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting r ramesh vishnoi and 4 more 5 2:47 PM You

set theory.pdf - Adobe Reader File Edit View Document Tools Window Help 17 / 18 162% Find

Theorem
Before we discuss Baye's theorem, we shall define the rule of elimination or theorem of total probability.

Rule of Elimination or Theorem of Total Probability.
If the event A can occur along with event E. suppose that event E occurs in n mutually exclusive ways E_1, E_2, \dots, E_n , then

$$P(A) = \sum_{i=1}^n P(E_i) \cdot P(A|E_i), \text{ provided } P(E_i) > 0, i=1, 2, \dots, n$$

Proof.

Meeting details ^

CC Turn on captions Smita Sonker is presenting

Type here to search

14:47 11-09-2020

(7) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting Smita Sonker and 3 more 2:47 PM You

set theory.pdf - Adobe Reader

Proof: $A = A \cap E_1 \cup A \cap E_2 \cup \dots \cup A \cap E_n$.
Events $(A \cap E_1), (A \cap E_2), \dots, (A \cap E_n)$ are mutually exclusive events.
 $\therefore P(A) = P[(A \cap E_1) \cup (A \cap E_2) \cup \dots \cup (A \cap E_n)]$
 $= P(A \cap E_1) + P(A \cap E_2) + \dots + P(A \cap E_n)$
 $= \sum_{i=1}^n P(A \cap E_i)$
using conditional probability

Meeting details ^



cc

Turn on captions

Smita Sonker
is presenting

...



Type here to search

14:47
11-09-2020

(7) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting Sahil Shende and 3 more 2:47 PM

set theory.pdf - Adobe Reader

$$P(A) = P(A \cap E_1) + P(A \cap E_2) + \dots + P(A \cap E_n)$$
$$= \sum_{i=1}^n P(A \cap E_i)$$

using conditional probability

$$P(A \cap E_i) = P(E_i) \cdot P(A|E_i)$$

$$\therefore P(A) = \sum_{i=1}^n P(E_i) \cdot P(A|E_i)$$

Baye's Theorem

Meeting details ^

Turn on captions

Smita Sonker is presenting



Type here to search



14:47
11-09-2020

(7) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

prathmesh dakh... and 3 more

5 2:48 PM You

set theory.pdf - Adobe Reader

File Edit View Document Tools Window Help

17 / 18 162% Find

Baye's Theorem

Let E_1, E_2, \dots, E_n be n mutually exclusive events of which one of them must occur. Let A be any event, then

$$P(E_i | A) = \frac{P(A \cap E_i)}{P(A)} = \frac{P(E_i)P(A|E_i)}{\sum_{i=1}^n P(E_i) \cdot P(A|E_i)}$$

Scanned with CamScanner

Meeting details ^



CC
Turn on captions

Up arrow
Smita Sonker
is presenting

...



Type here to search



14:48
11-09-2020

(7) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

Sahil Shende and 3 more

2:48 PM You

set theory.pdf - Adobe Reader

EX-34 : $\frac{6}{18}$

A bag contains 3 black and 4 white balls. Two balls are drawn at random one at a time without replacement. What is probability that second ball is white?

Sol: Let E_1 & E_2 be the events that first ball is black and first ball is white respectively. Let A be the event that second ball is white.

then

$$A = (A \cap E_1) \cup (A \cap E_2)$$

Meeting details ^



CC
Turn on captions

Up
Smita Sonker
is presenting



Type here to search



14:48
11-09-2020

(7) WhatsApp Meet - jpb-gfxi-jas Meet - nvo-hwuy-usf New Tab

meet.google.com/nvo-hwuy-usf

Smita Sonker is presenting

p prathmesh dakh... and 3 more

5 2:48 PM You

set theory.pdf - Adobe Reader

A = (A ∩ E₁) ∪ (A ∩ E₂)

$$P(A \cap E_1) = P(E_1) \cdot P(A/E_1)$$
$$= \left(\frac{3}{7}\right) \left(\frac{4}{8}\right) = \frac{2}{7}$$

Similarly P(A ∩ E₂) = P(E₂) · P(A/E₂)

$$= \left(\frac{4}{7}\right) \left(\frac{3}{8}\right) = \frac{2}{7}$$
$$\therefore P(A) = P(A \cap E_1) + P(A \cap E_2) = \frac{4}{7}$$

Meeting details ^



CC
Turn on captions

Up
Smita Sonker
is presenting

...



Type here to search



14:48
11-09-2020