

RAG Pipeline

Retrieval Results Report

Generated: 24 May 2026, 03:52 PM · Total queries: 20

Pipeline Configuration

Configuration	
Retrieval mode	bm25
Reranker	OFF
Adaptive weights	OFF
Fusion strategy	N/A
BM25 / Sem weights	N/A
Filtering	OFF
Cross-modal boost	ON
text_k / rerank_k / image_k	10 / 5 / 5
Embed model	BAAI/bge-large-en-v1.5
Image embed model	ViT-L-14 (laion2b_s32b_b82k)
LLM model(s)	gemma4:e2b
Chunk tokens / overlap	384 / 128

Query 1 of 20

User Query

The trajectory diagram of Voyager 1 and Voyager 2 shows clearly diverging paths after Saturn. Using both the trajectory image and the mission text, explain why the two paths diverge at that point, and what specific orbital geometry decision made Voyager 1's post-Saturn path irrecoverable for further planetary encounters.

Retrieved Text Context

[1] Flight trajectories of Voyager 1 and 2 through the outer solar system (left) and the launch on top of a Titan III-Centaur rocket (right). Images courtesy of NASA.

Taking advantage of this alignment would be two Voyager spacecraft, both beginning their long journeys in 1977. Voyager 2 launched first, on August 20, followed by Voyager 1 on September 5. Both spacecraft would first fly by Jupiter and use that planet's massive gravity to bend their trajectories to then fly by Saturn. Voyager 1 would also be targeted to fly by Saturn's moon Titan, which was known to have a dense atmosphere, a trajectory that would preclude any future planetary flybys. But the option (Source: Voyager Grand Tour PDF.pdf, page 1)

[2] Both Voyagers quietly coasted as they headed toward their next target, the ringed planet Saturn. Once again, Voyager 1 was first, beginning observations in August 1980 before making its closest approach 77,000 miles above Saturn's cloud tops on November 12, and finishing studies in December.

Significantly, Voyager 1 made a close flyby of Saturn's large moon Titan, larger than our own Moon and with a dense atmosphere, coming within 4,000 miles of its surface near its South Pole. The Titan encounter changed Voyager 1's trajectory so it could not make any further planetary encounters – thus beginning its long journey out of our solar system. Voyager 2 started its examination of Saturn in June 1981, making its closest approach at 63,000 miles on August 25, and finishing observations in September.

The instruments aboard the Voyagers returned a wealth of new information about Saturn and the cameras returned thousands of images of the planet, its rings and satellites. Among significant discoveries, the Voyagers found that Saturn's rings were far more complex than previously believed, Titan may have lakes of liquid hydrocarbons, Saturn has auroras near its poles much like Earth does, and that there were three previously unknown satellites.

(Source: Voyager Grand Tour PDF.pdf, page 3)

[3] Voyager's Grand Tour

By John Uri

Manager, History Office

NASA Johnson Space Center


<https://www.jsc.nasa.gov/history/>

In the early days of the Space Age, scientists realized that given the right planetary alignments it might be possible to use the gravity of one planet to change the trajectory of a spacecraft and send it on to another planet without expending any fuel. This slingshot or gravity assist trajectory principle was first tested by Mariner 10, which used the gravity of Venus to slingshot its way to Mercury in 1974.

A very rare planetary alignment would occur in the late 1970's allowing a spacecraft to visit all the outer planets (Jupiter, Saturn, Uranus, Neptune and Pluto) using gravity assists at each planet to send it on to the next. This unique alignment would not occur again for another 175 years! The initial ambitious plan, called the Grand Tour, was to send two pairs of spacecraft, one pair to visit Jupiter, Saturn and Pluto, the other to fly by Jupiter, Uranus and Neptune. However, the original plan was scaled back in the budget conscious early 1970's to just two less capable spacecraft visiting only Jupiter and Saturn, and Titan, Saturn's largest moon.

(Source: Voyager Grand Tour PDF.pdf, page 1)

Retrieved Images



Flight trajectories of Voyager 1 and 2 through the outer solar system (left) and the launch on top of a Titan III-Centaur rocket (right).
Images courtesy of NASA.

Query 2 of 20

User Query

The Earth-Moon photograph taken by Voyager 1 is described as the first single-frame image of both bodies together. Using the image itself and the surrounding mission text, explain what engineering and operational conditions had to be simultaneously satisfied to produce it, and why this milestone was considered a meaningful preview of capabilities rather than merely a publicity photograph.

Retrieved Text Context

[1] Just 13 days after its launch, Voyager 1 scored the first of its many firsts: at a distance of 7.25 million miles, it turned its camera back toward Earth and snapped the first ever photograph of the Earth-Moon system in a single frame, giving a sneak preview of the discoveries that lay ahead.

(Source: Voyager Grand Tour PDF.pdf, page 2)

[2] was kept open, if Voyager 1's Titan flyby was successful, to retarget Voyager 2 to send it on to Uranus and maybe even Neptune – assuming it would survive that long!

Just 13 days after its launch, Voyager 1 scored the first of its many firsts: at a distance of 7.25 million miles, it turned its camera back toward Earth and snapped the first ever photograph of the Earth-Moon system in a single frame, giving a sneak preview of the discoveries that lay ahead.

(Source: Voyager Grand Tour PDF.pdf, page 2)


[3] First photograph of the Earth-Moon system in a single frame (left), taken by Voyager 1 on September 18, 1977. Voyager spacecraft as it would appear in space (right). Images courtesy of NASA.

On their way to Jupiter, both Voyagers successfully passed through the Asteroid Belt without incident – two Pioneer spacecraft had made the crossing safely in 1973 and 1974 as pathfinders, proving that it was safe to do so. Voyager 1 arrived first, passing 217,000 miles above the giant planet's cloud tops on March 5, 1979. Two months before the close encounter, Voyager 1 began taking photographs of Jupiter and its moons and gathering other scientific data about the planet and its environment, a process that continued for about a month after the fly-by. Voyager 2 followed not too far behind, beginning scientific observations in April 1979 before passing 350,000 miles above Jupiter's cloud tops on July 9, 1979, and finishing observations about a month later.

The 11 instruments aboard each Voyager made amazing discoveries, not to mention the thousands of stunning photographs of Jupiter and its many moons. Among the more notable discoveries were that Jupiter has a faint ring system, its Great Red Spot is a large swirling storm system, the moon Io has active volcanoes, the moon Europa has a geologically young surface made of ice possibly floating on an ocean of liquid water, and that there were three previously undiscovered small moons.

(Source: Voyager Grand Tour PDF.pdf, page 2)

Retrieved Images



was kept open, if Voyager 1's Titan flyby was successful, to retarget Voyager 2 to send it on to Uranus and maybe even Neptune – assuming it would survive that long! Just 13 days after its launch, Voyager 1 scored the first of its many firsts: at a distance of 7.25 million miles, it turned its camera back toward Earth and snapped the first ever photograph of the Earth-Moon system in a single frame, giving a sneak preview of the discoveries that lay ahead. First photograph of the Earth-Moon system in a single frame (left), taken by Voyager 1 on September 18, 1977. Voyager spacecraft as it would appear in space (right). Images courtesy of NASA.

Query 3 of 20

User Query

Curiosity's Sky Crane landing image and the text description of the landing sequence together tell a more complete story than either alone. Using both the image of the sky crane lowering Curiosity and the technical description of the EDL sequence, explain why the sky crane approach was necessary rather than airbag-based landing used by earlier rovers, and what physical constraints of the crater site made this the only viable solution.

Retrieved Text Context

[1] National Aeronautics and Space Administration
Mars Science Laboratory/Curiosity


NASA's Mars Science Laboratory mission set down a large, mobile laboratory — the rover Curiosity — at Gale Crater, using precision landing technology that made one of Mars' most intriguing regions a viable destination for the first time. Within the first eight months of a 23-month primary mission, Curiosity met its major objective of finding evidence of a past environment well suited to supporting microbial life. Curiosity touches down on Mars after being lowered by its Sky Crane.

The rover studies the geology and environment of selected areas in the crater and analyzes samples drilled from rocks or scooped from the surface. The Mars Science Laboratory spacecraft launched from Cape Canaveral Air Force Station, Florida, on Nov. 26, 2011. Mars rover Curiosity carries the most advanced payload of scientific gear ever used on Mars' surface, landed successfully on the floor of Gale Crater on Aug. 6, 2012 Universal Time (evening of Aug. 5, Pacific Time), at 4.6 degrees south latitude, 137.4 degrees east longitude and minus 4,501 meters (2.8 miles) elevation. Investigate whether conditions have been favorable for microbial life and for preserving clues in the rocks about possible past life. Engineers designed the spacecraft to steer it- More than 400 scientists from around the world participate in the science operations. with a series of S-curve maneuvers similar to those used by astronauts piloting NASA space shuttles. During the three minutes before touchdown, the spacecraft slowed its descent with a parachute, then used retrorockets mounted (Source: mars-science-laboratory.pdf, page 1)

[2] south latitude, 137.4 degrees east longitude favorable for microbial life and for preserving and minus 4,501 meters (2.8 miles) elevation. clues in the rocks about possible past life. Engineers designed the spacecraft to steer it- More than 400 scientists from around the self during descent through Mars' atmosphere world participate in the science operations. with a series of S-curve maneuvers similar to those used by astronauts piloting NASA space shuttles. During the three minutes before touchdown, the spacecraft slowed its descent with a parachute, then used retrorockets mounted around the rim of its upper stage. In the final seconds, the upper stage acted as a sky crane, lowering the upright rover on a tether to land on The touchdown site, Bradbury Landing, is near the foot of a layered mountain, Aeolis Mons The rover's landing site, Gale Crater, is about the size of Connecticut and Rhode Island combined. ("Mount Sharp"). Selection of Gale Crater fol- (Source: mars-science-laboratory.pdf, page 1)

[3] capability of landing within a target area only about 20 kilometers (12 miles) long. That precision, about a five-fold improvement on earlier Mars landings, made sites eligible this main destination in September 2014. In the low lay- that would otherwise be excluded for encompassing ers of Mount Sharp during the rover's extended mission, nearby unsuitable terrain. The Gale Crater landing site is so researchers anticipate finding further evidence about hab- close to the crater wall and Mount Sharp that it would not itable past environments and about the evolution of the have been considered safe if the mission were not using Martian environment from a wetter past to a drier present. this improved precision. Science findings began months before landing. Measure- Curiosity is about twice as long (about 3 meters or 10 feet) ments that Curiosity made of natural radiation levels during and five times as heavy as NASA's twin Mars Exploration the flight from Earth to Mars will help NASA design for as- Rovers, Spirit and Opportunity, launched in 2003. It inher- tronaut safety on future human missions to Mars. ited many des

Retrieved Images



Mars Science Laboratory/Curiosity Curiosity touches down on Mars after being lowered by its Sky Crane. The Mars Science Laboratory spacecraft launched from Cape Canaveral Air Force Sta

position early in the mission provided evidence that the planet has lost much of its original atmosphere by a process favoring loss from the top of the atmosphere rather than interaction with the surface.

In the initial months of the surface mission, the rover team drove Curiosity eastward toward “Yellowknife Bay” to Curiosity’s first sample drilling, at a rock called “John Klein.”

Mars Science Laboratory/Curiosity

2

NASA Facts

(Source: mars-science-laboratory.pdf, page 2)

[3] capability of landing within a target area only about 20 kilometers (12 miles) long. That precision, about a five-fold improvement on earlier Mars landings, made sites eligible this main destination in September 2014. In the low lay-out that would otherwise be excluded for encompassing the base of Mount Sharp during the rover’s extended mission, nearby unsuitable terrain. The Gale Crater landing site is so researchers anticipate finding further evidence about habitable past environments and about the evolution of the Martian environment from a wetter past to a drier present. this improved precision.

Science findings began months before landing. Measurements that Curiosity made of natural radiation levels during the flight from Earth to Mars will help NASA design for as-Rovers, Spirit and Opportunity, launched in 2003. It inherited many design elements from them, including six-wheel In the first few weeks after landing, images from th

Retrieved Images



Curiosity is about twice as long (about 3 meters or 10 feet) and five times as heavy as NASA's twin Mars Exploration Rovers, Spirit and Opportunity, launched in 2003. It inherited many design elements from them, including six-wheel Curiosity's first sample drilling, at a rock called "John Klein." Mars Science Laboratory/Curiosity 2 NASA Facts

Query 5 of 20

User Query

The Hubble deep field image and the panel of faint early galaxies shown alongside it are described as evidence for galaxy growth over cosmic time. Using the visual appearance of those galaxies and the text about hierarchical galaxy formation, explain how the morphological evidence in the images connects to the theoretical prediction that large present-day galaxies assembled through mergers rather than forming fully intact.

Retrieved Text Context

[1] Tracing the Growth of Galaxies

Like documenting a child's development in a scrapbook, astronomers have used Hubble to capture the appearance of many developing galaxies throughout cosmic time. This is possible because of the mathematical relationship between cosmic distance and time: the deeper Hubble peers into space, the farther back it looks in time. As it happens, the most distant and earliest galaxies spied by Hubble are smaller and more irregularly shaped than today's grand spiral and elliptical galaxies. This is evidence that galaxies grew over time through mergers with other galaxies to become the giant systems we see today.

amounts of stars and gas in galaxies, the types and amounts of identifiable chemical elements present, and star-formation rates.

And the evolution continues. Hubble observations of our neighboring Andromeda galaxy (M31) have allowed astronomers to predict with certainty that a titanic collision between our Milky Way galaxy and Andromeda will inevitably take place beginning 4 billion years from now. The galaxy is now 2.5 million light-years away, but it is inescapably moving toward the Milky Way under the mutual pull of gravity between the two galaxies and the invisible dark matter that surrounds them both. The merger will likely result in the creation of a giant elliptical galaxy billions of years from now.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 4)

[2] Because the universe was smaller in the past, galaxies were more likely to interact with one another gravitationally. Some of Hubble's cosmic "snapshots" show fantastic stellar streamers pulled out and flung across space by colliding galaxies. They apparently settled over time into the more familiarly shaped galaxies seen closer to Earth and hence nearer to the present time. By carefully studying galaxies at different epochs, astronomers can see how galaxies changed and evolved over time. Among the things they investigate are the relative

Left: Hubble's Ultra Deep Field is one of the most distant looks into space ever. The cumulative exposure time needed to capture the image was about a million seconds (11 days).

Right: A sample of the faintest and farthest galaxies in the Hubble Ultra Deep Field shows that they were irregularly shaped and frequently interacting in the distant past.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 4)


[3] Viewing Galactic Details and Mergers

When Edwin Hubble discovered that the universe was a vast frontier of innumerable galaxies beyond our Milky Way, he categorized them according to three basic shapes: spiral, elliptical, and irregular. Later, the sharp vision of the space telescope named in his honor revealed unprecedented details in such galaxies. In addition, Hubble's images uncovered a plethora of oddball, peculiarly shaped galaxies that are more numerous the farther back into time the telescope looks. This is because the expanding universe was smaller long ago, and galaxies were both younger and more likely to interact since they were closer to one another.

Among this zoo of odd galaxies are "tadpole-like" objects and apparently merging systems dubbed "train-wrecks." For all their violence, galactic collisions take place at a glacial rate by human standards—timescales on the order of several hundred million years. Hubble's images, therefore, capture snapshots of galaxies at various stages of interaction. The merging of galaxies produces turbulence and tides that can induce new vigorous bursts of star formation within their interstellar gas clouds. These observed mergers form a preview of the coming collision between our own Milky Way and the neighboring Andromeda galaxy 4 billion years from now.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 18)

Retrieved Images



Left: Hubble's Ultra Deep Field is one of the most distant looks into space ever. The cumulative exposure time needed to capture the image was about a million seconds (11 days). Right: A sample of the faintest and farthest galaxies in the Hubble Ultra Deep Field shows that they were irregularly shaped and frequently interacting in the distant past.

Query 6 of 20

User Query

The dark matter page shows two views of galaxy cluster CI 0024+17 — one in visible light showing blue arcs, and one with a blue dark matter density overlay. Using both images together with the gravitational lensing text, explain the full chain of inference that connects the visible arc morphology in the first image to the quantitative dark matter mass estimate cited in the text.

Retrieved Text Context

[1] Top left: This sketch shows paths of light from a distant galaxy that is being gravitationally lensed. Top right: Gravitational lensing has created

three distorted images of the same background galaxy and five of a background quasar in this image of galaxy cluster SDSS J1004+4112.

Bottom: These two pictures show a massive galaxy cluster, CI 0024+17. The visible-light image (left) shows blue arcs among the yellowish

galaxies. These arcs are magnified, warped images of galaxies located behind the cluster whose light has been distorted and bent. The blue

overlay (right) shows the dark matter density needed to account for the gravitational distortions.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 6)

[2] Shining a Light on Dark Matter

Dark matter is an invisible form of matter that makes up most of the universe's mass and creates its underlying structure. Dark

matter's gravity drives normal matter (gas and dust) to collect and build up into stars and galaxies. Although astronomers cannot

see dark matter, they can detect its influence by observing how the gravity of massive galaxy clusters, which contain dark matter,

bends and distorts the light of more distant galaxies located behind the cluster. This phenomenon is called gravitational lensing.

Hubble's uniquely sharp vision allows astronomers to map the distribution of dark matter in space using gravitational lensing.

Large galaxy clusters contain both dark matter and normal matter. By observing the areas around massive clusters of galaxies

astronomers can identify warped background galaxies and reverse-engineer their distortions to reveal where the densest

concentrations of matter lie. Mathematical models of these results shed light on the location and properties of the lensing

material, both visible and invisible (dark). The universe appears to have about five times more dark matter than regular matter

and seems to be organized around an immense network of dark matter filaments that have grown over time.

Massive visible

structures, like galaxy clusters, are found at the intersections of these filaments.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 6)

[3] To the surprise of astronomers, Hubble observations along

with those of ground-based observatories have also shown

that the universe is not just expanding, but accelerating—a

discovery that won the 2011 Nobel Prize in Physics. Many

scientists believe this acceleration is caused by a "dark

energy" that pervades the universe. Dark energy can be

thought of as a sort of "antigravity" that is pushing galaxies

apart by stretching space at an increasing pace. The nature of this energy is a complete mystery, even though astronomers estimate that it makes up about 70 percent of the mass and energy in the entire universe. Though it cannot be measured directly using current technology, dark energy can be characterized by its effect on matter in the visible universe. By observing how dark energy behaves over time, astronomers hope to gain a better understanding of what it is and how it might affect the future of the cosmos.

Background photo: R. Gendler

(Source: [highlights_of_hubbles_exploration_of_the_universe.pdf](#), page 3)

Retrieved Images



Top left: This sketch shows paths of light from a distant galaxy that is being gravitationally lensed. Top right: Gravitational lensing has created three distorted images of the same background galaxy and five of a background quasar in this image of galaxy cluster SDSS J1004+4112. Bottom: These two pictures show a massive galaxy cluster, CI 0024+17. The visible-light image (left) shows blue arcs among the yellowish galaxies. These arcs are magnified, warped images of galaxies located behind the cluster whose light has been distorted and bent. The blue overlay (right) shows the dark matter density needed to account for the gravitational distortions.

Query 7 of 20

User Query

The image series of Supernova 1987A from 1994 through 2006 shows a ring of material progressively lighting up over twelve years. Using both the time-sequence images and the text description of the shock wave dynamics, explain what physical process the brightening ring documents, why it is observable only over a multi-year baseline, and what this reveals about the pre-explosion mass loss history of the progenitor star.

Retrieved Text Context

[1] Sept. 24, 1994

Feb. 6, 1998

Mar. 23, 2001

Supernova 1987A as seen by
Hubble in 1994.

Jan. 5, 2003

Dec. 15, 2004

Dec. 6, 2006

A shock wave of material from the central, exploded star of SN 1987A is slamming into an existing ring of material around the doomed star. As it does, various spots in the ring are heated and begin to glow. The nature of the strangely shaped pink feature in the center of the ring is not well understood.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 15)

[2] A shock wave of material from the central, exploded star of SN 1987A is slamming into an existing ring of material around the doomed star. As it does, various spots in the ring are heated and begin to glow. The nature of the strangely shaped pink feature in the center of the ring is not well understood.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 15)


[3] Documenting the Death Throes of Stars

Hubble has revealed unprecedented details in the appearance of Sun-like stars that have entered the death throes of their lives. Ground-based images suggested that many of these objects, called planetary nebulas (though they are unrelated to planets), have simple spherical shapes. Hubble has shown, however, that their shapes are much more varied and complex. Some look like pinwheels, others like butterflies, and still others like hourglasses. Such images yield insights into the complex dynamics that accompany a star's release of its outer gaseous layers before it collapses to form a white dwarf. Turning its sights to the tattered remains of a more massive star's explosive death, Hubble observations of Supernova 1987A revealed three mysterious rings of material encircling the doomed star. The telescope also spied bright spots on the middle ring's inner region caused by an expanding wave of matter slamming into it from the explosion. Likewise, Hubble's view of M1, the Crab Nebula, showed details never before seen about this mighty blast and the rapidly spinning pulsar that remains at its core.

Hubble has revealed the astounding variety and amazing complexity of planetary nebulas.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 14)

Retrieved Images



Sept. 24, 1994 Feb. 6, 1998 Mar. 23, 2001 Jan. 5, 2003 Dec. 15, 2004 Dec. 6, 2006 A shock wave of material from the central, exploded star of SN 1987A is slamming into an existing ring of material around the doomed star. As it does, various spots in the ring are heated and begin to glow. The nature of the strangely shaped pink feature in the center of the ring is not well understood.

Query 8 of 20

User Query

The Saturn montage image shows the planet alongside several of its moons at different scales. Using the image to identify visible moons and the text description of Voyager 1's Titan encounter, explain the specific trade-off that made Titan the highest-priority Saturn target, why that priority decision permanently closed Voyager 1's path to the outer planets, and what Voyager 2's different Saturn geometry preserved for the mission as a whole.

Retrieved Text Context

[1] Both Voyagers quietly coasted as they headed toward their next target, the ringed planet Saturn. Once again, Voyager 1 was first, beginning observations in August 1980 before making its closest approach 77,000 miles above Saturn's cloud tops on November 12, and finishing studies in December.

Significantly, Voyager 1 made a close flyby of Saturn's large moon Titan, larger than our own Moon and with a dense atmosphere, coming within 4,000 miles of its surface near its South Pole. The Titan encounter changed Voyager 1's trajectory so it could not make any further planetary encounters – thus beginning its long journey out of our solar system. Voyager 2 started its examination of Saturn in June 1981, making its closest approach at 63,000 miles on August 25, and finishing observations in September.

The instruments aboard the Voyagers returned a wealth of new information about Saturn and the cameras returned thousands of images of the planet, its rings and satellites. Among significant discoveries, the Voyagers found that Saturn's rings were far more complex than previously believed, Titan may have lakes of liquid hydrocarbons, Saturn has auroras near its poles much like Earth does, and that there were three previously unknown satellites.

(Source: Voyager Grand Tour PDF.pdf, page 3)

[2] Montage of Jupiter and its four largest moons, (left) and of Saturn and several of its moons (right). Images are not to scale. Courtesy of NASA.

Two more stops: Uranus and Neptune

Voyager 2's flyby of Saturn set it up to automatically fly by Uranus after a coast of five and a half years. It began to study the Uranian system in November 1985 and flew within 50,600 miles of its cloud tops on January 24, 1986, before concluding its observations in February. In conducting the first ever robotic reconnaissance of Uranus, Voyager 2 observed the planet's atmosphere, satellites and thin ring system, discovering 11 new moons and a previously unknown magnetic field and returning about 8,000 pictures. And then, it was on to Neptune, arriving there after another coast of three and a half years. Beginning its study in June 1989, Voyager 2 flew within just 3,076 miles to Neptune's cloud tops on August 25, concluding its observations in October. That close pass to Neptune allowed detailed observations of its largest moon Triton. During this first ever close up study of Neptune, Voyager 2 returned about 10,000 photographs of the planet, its satellites and dark rings, discovering 6 new moons, a Great Dark Spot on

(Source: Voyager Grand Tour PDF.pdf, page 3)

[3] Flight trajectories of Voyager 1 and 2 through the outer solar system (left) and the launch on top of a Titan III-Centaur rocket (right). Images courtesy of NASA.

Taking advantage of this alignment would be two Voyager spacecraft, both beginning their long journeys in 1977. Voyager 2 launched first, on August 20, followed by Voyager 1 on September 5. Both spacecraft would first fly by Jupiter and use that planet's massive gravity to bend their trajectories to then fly by Saturn. Voyager 1 would also be targeted to fly by Saturn's moon Titan, which was known to have a dense atmosphere, a trajectory that would preclude any future planetary flybys. But the option

(Source: Voyager Grand Tour PDF.pdf, page 1)

Retrieved Images



*Montage of Jupiter and its four largest moons, (left) and of Saturn and several of its moons (right). Images are not to scale.
Courtesy of NASA.*

Query 9 of 20

User Query

The Uranus montage and the Neptune-Triton montage each show a planet alongside several moons. Using these images in combination with the text on Voyager 2's encounters, explain why Voyager 2 discovered 11 moons at Uranus and 6 at Neptune despite having far less observing time at each world than it had spent at Jupiter, and what this implies about the relationship between mission geometry and small-body discovery yield.

Retrieved Text Context

[1] Montage of Jupiter and its four largest moons, (left) and of Saturn and several of its moons (right). Images are not to scale. Courtesy of NASA.

Two more stops: Uranus and Neptune

Voyager 2's flyby of Saturn set it up to automatically fly by Uranus after a coast of five and a half years. It began to study the Uranian system in November 1985 and flew within 50,600 miles of its cloud tops on January 24, 1986, before concluding its observations in February. In conducting the first ever robotic reconnaissance of Uranus, Voyager 2 observed the planet's atmosphere, satellites and thin ring system, discovering 11 new moons and a previously unknown magnetic field and returning about 8,000 pictures. And then, it was on to Neptune, arriving there after another coast of three and a half years. Beginning its study in June 1989, Voyager 2 flew within just 3,076 miles to Neptune's cloud tops on August 25, concluding its observations in October. That close pass to Neptune allowed detailed observations of its largest moon Triton. During this first ever close up study of Neptune, Voyager 2 returned about 10,000 photographs of the planet, its satellites and dark rings, discovering 6 new moons, a Great Dark Spot on (Source: Voyager Grand Tour PDF.pdf, page 3)

[2] Montage of Uranus and several of its larger moons (left) and of Neptune and its largest moon Triton (right). Courtesy of NASA.

On February 14, 1990, more than 12 years after it began its journey and shortly before its cameras were permanently turned off to conserve power, Voyager 1 spun around and pointed them back into the solar system. In a mosaic of 60 images, it was able to capture a "family portrait" of six of the solar system's planets, including a pale blue dot called Earth. It was fitting that these were the last pictures from either Voyager.

(Source: Voyager Grand Tour PDF.pdf, page 4)

[3] First photograph of the Earth-Moon system in a single frame (left), taken by Voyager 1 on September 18, 1977. Voyager spacecraft as it would appear in space (right). Images courtesy of NASA.

On their way to Jupiter, both Voyagers successfully passed through the Asteroid Belt without incident – two Pioneer spacecraft had made the crossing safely in 1973 and 1974 as pathfinders, proving that it was safe to do so. Voyager 1 arrived first, passing 217,000 miles above the giant planet's cloud tops on March 5, 1979. Two months before the close encounter, Voyager 1 began taking photographs of Jupiter and its moons and gathering other scientific data about the planet and its environment, a process that continued for about a month after the fly-by. Voyager 2 followed not too far behind, beginning scientific observations in April 1979 before passing 350,000 miles above Jupiter's cloud tops on July 9, 1979, and finishing observations about a month later.

The 11 instruments aboard each Voyager made amazing discoveries, not to mention the thousands of stunning photographs of Jupiter and its many moons. Among the more notable discoveries were that Jupiter has a faint ring system, its Great Red Spot is a large swirling storm system, the moon Io has active volcanoes, the moon Europa has a geologically young surface made of ice possibly floating on an ocean of liquid water, and that there were three previously undiscovered small moons.

(Source: Voyager Grand Tour PDF.pdf, page 2)

Retrieved Images



*Montage of Jupiter and its four largest moons, (left) and of Saturn and several of its moons (right). Images are not to scale.
Courtesy of NASA. Two more stops: Uranus and Neptune*

Query 10 of 20

User Query

Curiosity's self-portrait assembled from MAHLI images is shown in the document. Using the image to assess what the composite reveals about rover configuration and condition, and the technical text describing MAHLI's capabilities and arm placement, explain why this imaging method provides more diagnostic value for mission engineers than a single wide-angle shot from a fixed camera would.

Retrieved Text Context

[1] The mission uses radio relays via Mars orbiters as the principal means of communication between Curiosity and the Deep Space Network's antennas on Earth. In the first two years after Curiosity's landing, the orbiters downlinked 48 gigabytes of data from Curiosity.

In April 2004, NASA solicited proposals for specific instruments and investigations to be carried by Mars Science Laboratory. The agency selected eight of the proposals later that year and also reached agreements with Russia and Spain to carry instruments those nations provided. A suite of instruments named Sample Analysis at Mars analyzes samples of material collected and delivered by A self-portrait of Curiosity, built up from pictures taken by its Mars Hand Lens Imager.

the rover's arm, plus atmospheric samples. It includes a gas chromatograph, a mass spectrometer and a tunable drive, a rocker-bogie suspension system, and cameras laser spectrometer with combined capabilities to identify a mounted on a mast to help the mission's team on Earth wide range of carbon-containing compounds and determine exploration targets and driving routes. Unlike earlier mine the ratios of different isotopes of key elements. Iso-rovers, Curiosity carries equipment to gather and process tope ratios are clues to understanding the history of Mars' samples of rocks and soil, distributing them to onboard atmosphere and water. The principal investigator is Paul test chambers inside analytical instruments.

Mahaffy of NASA's Goddard Space Flight Center, Green-NASA's Jet Propulsion Laboratory (JPL), Pasadena, Calif., builder of the Mars Science Laboratory, engineered Curi-An X-ray diffraction and fluorescence instrument called osity to roll over obstacles up to 65 centimeters (25 inch-CheMin also examines samples gathered by the robotic es) high and to travel up to about 200 meters (660 feet) (Source: mars-science-laboratory.pdf, page 3)


[2] a stereo Navigation Camera on its mast and low-slung, of the area illuminated by the beam. The laser and tele-stereo Hazard-Avoidance cameras. The wide view of the scope sit on the rover's mast. Chemcam also serves as Navigation Camera is also used to aid targeting of other a passive spectrometer to measure composition of the

instruments and to survey the sky for clouds and dust. surface and atmosphere. Roger Wiens of Los Alamos Equipment called the Sample Acquisition/Sample Preparation and Handling System includes tools to remove dust from rock surfaces, scoop up soil, drill into rocks to collect powdered samples from rocks' interiors, sort samples by particle size with sieves, and deliver samples to laboratory. This information is necessary for planning human exploration of Mars and is relevant to assessing the planet's ability to harbor life. The principal investigator is Donald B. Hassler of Southwest Research Institute, Boulder, Colo. In the two minutes before landing, the Mars Descent Imager captured color, high-definition video of the landing sphere, to aid in design of future missions. region to provide geological context for the investigations Program/Project Management on the ground and to aid precise determination of the landing site. The Mars Science Laboratory is managed for NASA's Science Mission Directorate, Washington, D.C., by JPL, a division of the California Institute of Technology in Pasadena. Michael Malin is principal investigator.

(Source: mars-science-laboratory.pdf, page 4)

[3] samples of rocks and soil, distributing them to onboard test chambers inside analytical instruments

Retrieved Images



A self-portrait of Curiosity, built up from pictures taken by its Mars Hand Lens Imager. drive, a rocker-bogie suspension system, and cameras mounted on a mast to help the mission's team on Earth select exploration targets and driving routes. Unlike earlier

Query 11 of 20

User Query

The Cepheid star brightness sequence images from the Andromeda galaxy and the distant supernova comparison images from 1995 and 2002 are presented as two complementary distance-measurement methods. Using both image sets and the text about refining the Hubble constant, explain why astronomers need both techniques rather than relying on only one, and how the combination of refined distances and velocity measurements changed the estimated age of the universe.

Retrieved Text Context

[1] Above: Astronomers use cyclical changes in the brightness of Cepheid stars to determine astronomical distances. The arrow points to a Cepheid star in the Andromeda galaxy observed by Hubble (inset boxes).

Right: Certain supernovas have a characteristic maximum brightness that can be used to calculate their distances from Earth. Refining celestial distances enables astronomers to better calculate the expansion rate of the universe. The arrow points to a distant supernova discovered in an area of the sky first imaged in 1995 called the Hubble Deep Field. Astronomers found the supernova when they targeted the same area of sky again in 2002 and saw a change.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 3)

[2] Discovering a Runaway Universe

Our cosmos is getting bigger. Nearly a century ago Edwin Hubble measured the expansion rate of the universe. This value, called the Hubble constant, is an essential ingredient needed to determine the age, size, and fate of the cosmos. Before Hubble was launched, the value of the Hubble constant was imprecise, and calculations of the universe's age ranged from 10 billion to 20 billion years. Now, astronomers using Hubble have refined their estimates of the universe's present expansion rate and are working to make it more accurate. They do this by getting better galaxy distance measurements from Hubble and coupling these values with the best galaxy-velocity measurements obtained from other telescopes. Scientists measure distances by comparing the brightness of a known object in our galaxy (like a star or an exploded star) to that of a similar object in a distant galaxy. With Hubble's refined distance values, calculations currently put the age of the universe as 13.8 billion years.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 3)


[3] Because the universe was smaller in the past, galaxies were more likely to interact with one another gravitationally. Some of Hubble's cosmic "snapshots" show fantastic stellar streamers pulled out and flung across space by colliding galaxies. They apparently settled over time into the more familiarly shaped galaxies seen closer to Earth and hence nearer to the present

time. By carefully studying galaxies at different epochs, astronomers can see how galaxies changed and evolved over time. Among the things they investigate are the relative
Left: Hubble's Ultra Deep Field is one of the most distant looks into space ever. The cumulative exposure time needed to capture the image was about a million seconds (11 days).

Right: A sample of the faintest and farthest galaxies in the Hubble Ultra Deep Field shows that they were irregularly shaped and frequently interacting in the distant past.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 4)

Retrieved Images



Background photo: R. Gendler Above: Astronomers use cyclical changes in the brightness of Cepheid stars to determine astronomical distances. The arrow points to a Cepheid star in the Andromeda galaxy observed by Hubble (inset boxes).

Query 12 of 20

User Query

The M84 galaxy images show a camera view with a dark dust band and a spectrograph plot with dramatic color shifts from blue to red. Using both images and the text on black hole mass measurements, explain exactly what physical phenomenon the color shift encodes, why the spectrograph slit position relative to the core is critical to the measurement, and how this technique established the mass relationship between black holes and their host galaxies.

Retrieved Text Context

[1] Shown here are two Hubble views of the galaxy M84. Left: This camera view shows the bright core at the center of the galaxy surrounded by a (vertical) dark band of gas and dust. Right: This plot was generated by passing light near the core of the galaxy through a spectrograph. The thin, vertical rectangle in the center of the left panel shows the size and shape of the spectrograph's sampling slit. Spectra taken left and right of the slit's position, which is centered on the core of the galaxy, show a dramatic shift in color to blue or red. Blueshifted light indicates that the emitting light source is moving toward Earth, while redshifted light indicates objects that are moving away. At 880,000 miles per hour, stars and glowing gases nearest to the core of M84 are moving the fastest. They are circling a black hole at the center of the galaxy, and so appear to be moving rapidly toward Earth on the left and receding on the right.
(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 7)


[2] Realizing Monster Black Holes Are Everywhere
Hubble provided decisive evidence that the hubs of most galaxies contain enormous black holes, which have the mass of millions or even billions of stars. Not only are black holes resident in almost every galaxy, but somehow their sizes correspond. A Hubble census of galaxies showed that a black hole's mass is dependent on the mass of its host galaxy's central bulge of stars: the larger the galaxy, the larger the black hole. This close relationship may be evidence that black holes grew along with their galaxies, devouring a fraction of the galaxy's mass. Hubble also provided astronomers the first-ever views of material encircling black holes in large, flat disks. Combining images with data from Hubble's spectrographs, researchers have peered into the center of many galaxies and established the existence of large black holes. In a census performed by Hubble in the late 1990s, galaxies NGC 3379 and NGC 3377 were found to have black holes that "weighed in" at 50 million and over 100 million solar masses, respectively, and NGC 4486B was revealed to have a double nucleus at its core.
(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 7)

[3] Combining images with data from Hubble's spectrographs, researchers have peered into the center of many galaxies and established the existence of large black holes. In a census performed by Hubble in the late 1990s, galaxies NGC 3379 and NGC 3377 were found to have black holes that "weighed in" at 50 million and over 100 million solar masses, respectively, and NGC 4486B was revealed to have a double nucleus at

its core.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 7)

Retrieved Images



Realizing Monster Black Holes Are Everywhere

Query 13 of 20

User Query

The Jupiter impact sequence showing cloud darkening from Comet Shoemaker-Levy 9, the close-up plume image from the first large fragment impact, and the Saturn aurora image are all on the same or adjacent pages. Using these three images and the text covering both the impact events and aurora observations, explain what each image type contributes uniquely to understanding planetary atmospheric dynamics, and why continuous ultraviolet monitoring capability is the common prerequisite for all three scientific results.

Retrieved Text Context

[1] This image series, starting in the lower right corner, shows the darkening of Jupiter's clouds as a result of the impact of Comet Shoemaker-Levy 9.

Hubble captured a colossal plume rising above the limb of Jupiter from the impact of the first large fragment of Comet Shoemaker-Levy 9.

(Times along the left side are in hour:minute format.)

Jupiter's moons also have yielded important clues in the search for life beyond Earth. Hubble provided the best evidence yet for an underground saltwater ocean on Ganymede, the largest moon in the solar system, by detecting related activity in Ganymede's own auroras. This subterranean ocean is thought to have more water than all the water on Earth's surface.

Hubble also recorded evidence of transient changes in the atmosphere above the surface of Jupiter's moon Europa.

Astronomers suspect that these disturbances are caused by gas plumes expelled from a subsurface ocean. Identifying liquid water is crucial in the search for habitable worlds beyond Earth and in the quest to find life as we know it.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 8)

[2] Studying the Outer Planets and Moons

Hubble has witnessed impacts on Jupiter that were produced by minor bodies in the solar system. The latest collision observed by Hubble occurred in 2009 when a suspected asteroid plunged into Jupiter's atmosphere, leaving a temporary dark feature the size of the Pacific Ocean. In 1994 Hubble watched 21 fragments of Comet Shoemaker-Levy 9 bombard the giant planet sequentially—the first time astronomers witnessed such an event. Each impact left a temporary black, sooty scar within Jupiter's clouds.

Jupiter is well known for its Great Red Spot, a giant storm roughly the size of Earth that has been continuously visible since at least the late 1800s. The mammoth storm has been shrinking in size for at least 80 years. Astronomers now use Hubble regularly to measure the Red Spot's size and investigate why it is slowly disappearing.

Hubble also made the first images of bright auroras at the northern and southern poles of Saturn and Jupiter.

Auroras are brilliant curtains of light that appear in the upper atmosphere of a planet with a magnetic field. They develop when electrically charged particles trapped in the magnetic


field spiral inward at high speeds toward the north and south magnetic poles. When these particles hit the upper atmosphere, they excite atoms and molecules there causing them to glow in a similar process to that of a fluorescent light. This image series, starting in the lower right corner, shows the darkening of Jupiter's clouds as a result of the impact of Comet Shoemaker-Levy 9.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 8)

[3] Top left: Jupiter's trademark Great Red Spot—a swirling, 300-mile-per-hour storm—has shrunk to its smallest size ever. Astronomers have followed this gradual downsizing since the 1930s and are now keenly monitoring its shrinkage and investigating its cause with Hubble. Top right: Hubble's ability to capture ultraviolet light revealed a region of glowing auroras over the pole of Saturn. While sharing some features of Earth's auroras and others of Jupiter's, Saturn's auroras are different from both and are likely unique in the solar system. Below left: The blue areas mapped on Jupiter's moon Europa mark spots where Hubble found spectroscopic evidence of oxygen and hydrogen—the two elements that form the water molecule. Scientists believe this is most likely the result of water vapor plumes erupting from beneath the surface, with recent visual evidence of these plumes also seen by Hubble (below right). The pictures of Europa are a combination of shots gathered by NASA's Voyager and Galileo missions.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 9)

Retrieved Images



Studying the Outer Planets and Moons This image series, starting in the lower right corner, shows the darkening of Jupiter's clouds as a result of the impact of Comet Shoemaker-Levy 9. Hubble captured a colossal plume rising above the limb of Jupiter from the impact of the first large fragment of Comet Shoemaker-Levy 9. (Times along the left side are in hour:minute format.)

Query 14 of 20

User Query

The TW Hydrae disk image shows a gap structure around the star, and the 1997 and 2012 Beta Pictoris disk images show an edge-on disk changing over fifteen years. Using both sets of images alongside the planet-formation text, explain what specific morphological features in each image constitute evidence for an embedded or nearby planet, why temporal baseline matters for the Beta Pictoris case but not for TW Hydrae, and how these two systems together support a dynamic rather than static picture of disk evolution.

Retrieved Text Context

[1] Finding Planetary Construction Zones

Astronomers used Hubble to confirm that planets form in dust disks around stars. The telescope first resolved protoplanetary disks around nearly 200 stars in the bright Orion Nebula.

Looking at nearby stars elsewhere in the sky, Hubble completed the largest and most sensitive visible-light imaging survey of dusty debris disks, which were probably created by collisions between leftover objects from planet formation.

Two particular stars illustrate these findings: TW Hydrae and Beta Pictoris. Using a mask to block the star's bright light, Hubble scientists spotted a mysterious gap in a vast protoplanetary disk of gas and dust swirling around the star TW Hydrae. The gap is most likely caused by a growing, unseen planet that is gravitationally sweeping up material and carving out a lane in the disk like a snowplow. It is 1.9 billion miles wide and not yet completely cleared of material. Researchers have also noted changes in the planetary disk surrounding Beta Pictoris. By masking out the star's light, scientists have studied changes in the orbiting material caused by a massive planet embedded within its dust disk. Astronomers spotted the planet using the European Southern Observatory.

Here are a few of the many protoplanetary disks that were discovered by Hubble in the large, majestic Orion Nebula. The disks appear as silhouettes, illuminated from behind by the glowing gases of the nebula.

Below left and right: Hubble has provided detailed pictures showing changes in the large, edge-on, gas-and-dust disk encircling the 20-million-year-old star Beta Pictoris.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 17)

[2] Below left and right: Hubble has provided detailed pictures showing changes in the large, edge-on, gas-and-dust disk encircling the 20-million-year-old star Beta Pictoris.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 17)

[3] Here are a few of the many protoplanetary disks that were discovered by Hubble in the

large, majestic Orion Nebula.

The disks appear as silhouettes,
illuminated from behind by the
glowing gases of the nebula.

Below left and right: Hubble has provided detailed pictures showing
changes in the large, edge-on, gas-and-dust disk encircling the
20-million-year-old star Beta Pictoris.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 17)

Retrieved Images



*Below left and right: Hubble has provided detailed pictures showing changes in the large, edge-on, gas-and-dust disk encircling the
20-million-year-old star Beta Pictoris.*

Query 15 of 20

User Query

The Golden Record carried by both Voyager spacecraft is described in the text, and the image shows both the record face and its engraved instruction cover. Using the image of the record and the text description of its contents and purpose, explain what specific categories of information were encoded on the record, what physical provision was made to ensure an extraterrestrial finder could play it, and why the record was considered a scientifically motivated decision rather than merely a symbolic gesture given the trajectory constraints of the Voyager Interstellar Mission.

Retrieved Text Context

[1] Each Voyager carries a Golden Record, providing potential finders with information about its creator, the location of and information about the home planet. Courtesy of NASA.

The Golden Record is available commercially, for streaming, mp3 or Audio CD. A detailed description of the contents are summarized in the 1978 book *Murmurs of Earth: The Voyager Interstellar Record*, by Carl Sagan et al., also available commercially.

(Source: Voyager Grand Tour PDF.pdf, page 5)

[2] With their planetary encounters over, the spacecraft began the Voyager Interstellar Mission (VIM) in 1989, in which they are going where no man-made spacecraft has gone before, exploring uncharted areas of the outer solar system and eventually crossing the boundary into true interstellar space.

Because this part of space is so poorly understood, it took some time to analyze the data that Voyager sent back and determine when that transition took place – it is now believed that Voyager 1 crossed the boundary in August 2012, and Voyager 2 should follow suit in the next few years. Forty years after they were launched, both spacecraft are still operating more than 10 billion miles from their home planet, although most science instruments have been turned off to conserve power. Based on current projections, it is estimated that they can continue to function until about 2025.

In their current paths, neither Voyager will come within less than a light-year from any star in the next tens of thousands of years, long after they will have ceased to communicate with Earth. But, just in case any civilization is able to capture them to determine their origin, each Voyager is equipped with a golden record (think 1970's technology!) with coded information about the spacecraft and its makers.

Contained on the record are recordings of sounds from Earth, including music, greetings in multiple languages and even whale songs, as well as photographs of various Earthly activities. Just hope those aliens have turntables! Even a stylus was thoughtfully provided with the record, along with instructions on how to play it.

(Source: Voyager Grand Tour PDF.pdf, page 5)

[3] Hubble's discoveries and memorable photos have also reinvigorated the public's interest in astronomy. Along with pictures of the telescope and the astronauts who launched and serviced it during six space shuttle missions, certain memorable science images have become cultural icons. They appear regularly on book covers, musical albums, clothing, TV shows, movies, and even ecclesiastical stained-glass windows.

This booklet features thirteen representative topics with eye-catching images and thought-provoking discoveries. Selecting this set from the thousands of amazing Hubble observations was difficult, but these serve to highlight some of Hubble's greatest scientific achievements to date. Additional images and information can be found at <http://nasa.gov/hubble> and <http://hubblesite.org>.

Right: Astronaut Mike Good maneuvers to repair the Space Telescope Imaging Spectrograph during the final

Spectrograph during the final Hubble

Hubble servicing mission in May 2009. Periodic upgrades

servicing mission in May 2009. Periodic upgrades

have kept the telescope equipped with state-of-the-art instruments, which have given

have kept the telescope equipped with state-of-the-art instruments, which have given

astronomers increasingly better views of the cosmos.

astronomers increasingly better views of the cosmos.

Cover image: Veil Nebula

Cover image: Veil Nebula

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 2)

Retrieved Images



Each Voyager carries a Golden Record, providing potential finders with information about its creator, the location of and information about the home planet. Courtesy of NASA.

Query 16 of 20

User Query

Curiosity's radioisotope power system is described in detail in the text, and the rover's operational scale is visible in its self-portrait and the Gale Crater site image. Using the power system description alongside the mission timeline and data volume figures given in the text, construct a precise argument for why a nuclear RTG was the only viable power architecture for this mission — addressing operating lifespan, power budget at launch versus two years post-landing, terrain capability, and the communication data throughput achieved in the first two years.

Retrieved Text Context

[1] capability of landing within a target area only about 20 kilometers (12 miles) long. That precision, about a five-fold improvement on earlier Mars landings, made sites eligible this main destination in September 2014. In the low lay-ers of Mount Sharp during the rover's extended mission, that would otherwise be excluded for encompassing nearby unsuitable terrain. The Gale Crater landing site is so researchers anticipate finding further evidence about hab-itable past environments and about the evolution of the have been considered safe if the mission were not using Martian environment from a wetter past to a drier present. this improved precision.

Science findings began months before landing. Measure-ments that Curiosity made of natural radiation levels during and five times as heavy as NASA's twin Mars Exploration the flight from Earth to Mars will help NASA design for as-Rovers, Spirit and Opportunity, launched in 2003. It inher-tronaut safety on future human missions to Mars.

ited many design elements from them, including six-wheel In the first few weeks after landing, images from the rover showed that Curiosity touched down right in an area where water once coursed vigorously over the surface.

The evidence for stream flow was in rounded pebbles mixed with hardened sand in conglomerate rocks at and near the landing site. Analysis of Mars' atmospheric com-position early in the mission provided evidence that the planet has lost much of its original atmosphere by a pro-cess favoring loss from the top of the atmosphere rather than interaction with the surface.

In the initial months of the surface mission, the rover team (Source: mars-science-laboratory.pdf, page 2)


[2] National Aeronautics and Space Administration
Mars Science Laboratory/Curiosity
NASA's Mars Science Laboratory mission set
down a large, mobile laboratory — the rover
Curiosity — at Gale Crater, using precision

landing technology that made one of Mars' most intriguing regions a viable destination for the first time. Within the first eight months of a 23-month primary mission, Curiosity met its major objective of finding evidence of a past environment well suited to supporting microbial life. Curiosity touches down on Mars after being lowered by its Sky Crane.

The rover studies the geology and environment of selected areas in the crater and analyzes samples drilled from rocks or scooped from the surface. The Mars Science Laboratory spacecraft launched from Cape Canaveral Air Force Station, Florida, on Nov. 26, 2011. Mars rover of scientific gear ever used on Mars' surface, Curiosity landed successfully on the floor of Gale Crater on Aug. 6, 2012 Universal Time, at 4.6 degrees south latitude, 137.4 degrees east longitude, and minus 4,501 meters (2.8 miles) elevation. Its assignment: Investigate whether conditions have been favorable for microbial life and for preserving clues in the rocks about possible past life. Engineers designed the spacecraft to steer it- More than 400 scientists from around the world participate in the science operations. self during descent through Mars' atmosphere with a series of S-curve maneuvers similar to those used by astronauts piloting NASA space shuttles. During the three minutes before touchdown, the spacecraft slowed its descent with a parachute, then used retrorockets mounted (Source: mars-science-laboratory.pdf, page 1)

[3] samples of rocks and soil, distributing them to onboard test chambers inside analytical instruments. The principal investigator is Paul Mahaffy of NASA's Goddard Space Flight Center, Green- NASA's Jet Propulsion Laboratory (JPL), Pasadena, Calif., builder of the Mars Science Laboratory, engineered Curiosity. An X

Retrieved Images



favorable for microbial life and for preserving clues in the rocks about possible past life. More than 400 scientists from around the world participate in the science operations. The rover's landing site, Gale Crater, is about the size of Connecticut and Rhode Island combined.

Query 17 of 20

User Query

The multi-panel mosaic of planetary nebulas on the death-of-stars page shows at least a dozen distinct morphologies — pinwheels, hourglasses, butterflies, rings, and asymmetric lobes. Using the image grid and the text describing planetary nebula formation physics, explain why ground-based astronomy predicted simple spherical shells while Hubble revealed this diversity, what specific physical processes the non-spherical morphologies imply about the dying star's interior dynamics, and why this discovery is scientifically connected to Hubble's later observations of Supernova 1987A's three-ring structure.

Retrieved Text Context

[1] Documenting the Death Throes of Stars

Hubble has revealed unprecedented details in the appearance of Sun-like stars that have entered the death throes of their lives. Ground-based images suggested that many of these objects, called planetary nebulas (though they are unrelated to planets), have simple spherical shapes. Hubble has shown, however, that their shapes are much more varied and complex. Some look like pinwheels, others like butterflies, and still others like hourglasses. Such images yield insights into the complex dynamics that accompany a star's release of its outer gaseous layers before it collapses to form a white dwarf. Turning its sights to the tattered remains of a more massive star's explosive death, Hubble observations of Supernova 1987A revealed three mysterious rings of material encircling the doomed star. The telescope also spied bright spots on the middle ring's inner region caused by an expanding wave of matter slamming into it from the explosion. Likewise, Hubble's view of M1, the Crab Nebula, showed details never before seen about this mighty blast and the rapidly spinning pulsar that remains at its core.

Hubble has revealed the astounding variety and amazing complexity of planetary nebulas.
(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 14)

[2] objects as nearby and small as the collision of asteroids to distant star-forming galaxies that date back to when the universe was only three percent of its current age. In fact, Hubble observations have played a key role in discovering and characterizing the mysterious dark energy that now appears to permeate space. Results like these have changed our fundamental understanding of the cosmos. Well into its third operational decade, Hubble is still extremely productive. The orbiting telescope has taken over a million observations and provided data that astronomers have used to write more than 17,000 peer-reviewed scientific publications on a broad range of topics, from planet formation to gigantic black holes. These papers have been referenced in other publications approximately 900,000 times, and this total increases, on average, by more than 150 per day.

Every current astronomy textbook includes contributions from the observatory. Today's college undergraduates have not known

a time in their lives when astronomers were not actively making discoveries with Hubble data.

One of

One of Hubble

Hubble's most iconic images is of this portion of the Eagle Nebula

's most iconic images is of this portion of the Eagle Nebula

(M16). Dubbed the "Pillars of Creation," it shows star birth. Imbedded in

(M16). Dubbed the "Pillars of Creation," it shows star birth. Imbedded in

the tips of the finger-like protrusions at the top of the columns are dense,

the tips of the finger-like protrusions at the top of the columns are dense,

gaseous globules within which stars are being born.

gaseous globules within which stars are being born.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 2)

[3] Supernova 1987A as seen by

Hubble in 1994.

Jan. 5, 2003

Dec. 15, 2004

Dec. 6, 2006

A shock wave of material from the central, exploded star of SN 1987A is slamming into an existing ring of material around the doomed star. As it does, various spots in the ring are heated and begin to glow. The nature of the strangely shaped pink feature in the center of the ring is not well understood.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 15)

Retrieved Images



Supernova 1987A as seen by Hubble in 1994.

Query 18 of 20

User Query

The Voyager family portrait taken on February 14, 1990, is described as a mosaic of 60 images capturing six solar system planets. Using the image of the portrait mosaic and the surrounding text, explain the specific operational conditions that made this imaging session both uniquely possible and permanently terminal, what the phrase 'pale blue dot' refers to and which planet it identifies, and why the text describes these as the last pictures from either Voyager.

Retrieved Text Context

[1] On February 14, 1990, more than 12 years after it began its journey and shortly before its cameras were permanently turned off to conserve power, Voyager 1 spun around and pointed them back into the solar system. In a mosaic of 60 images, it was able to capture a “family portrait” of six of the solar system’s planets, including a pale blue dot called Earth. It was fitting that these were the last pictures from either Voyager.

(Source: Voyager Grand Tour PDF.pdf, page 4)

[2] Family portrait of six planets taken by Voyager 1 on February 14, 1990, when it was 6 B km from Earth. Not exactly close encounters

(Source: Voyager Grand Tour PDF.pdf, page 4)

[3] Astronomers used Hubble to take what might be the first-ever visible-light pictures of an extrasolar planet, named Fomalhaut b.


At the time of Hubble’s launch in 1990, astronomers had not found any planets outside our solar system. Scientists have now confirmed the existence of more than 4,000 extrasolar planets, most of them discovered by NASA’s Kepler space observatory and by ground-based telescopes. Hubble, however, has made some unique contributions to the planet hunt.

Neptune-sized planet due to extreme radiation from the star. This planet could explain the existence of so-called hot super-Earths, which might be stripped-down versions of this planet that have exposed a rocky core.

Using another technique called gravitational microlensing, astronomers have also used Hubble to confirm the existence of a Saturn-mass planet orbiting two small, faint stars in a tight orbit around each other. Gravitational microlensing occurs when the gravity of a moving foreground star bends and amplifies the light of a background star that temporarily aligns with it along our line of sight. Details in the character of the brightening reveal clues to the nature of the foreground star and any planets it may have.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 5)

Retrieved Images



Family portrait of six planets taken by Voyager 1 on February 14, 1990, when it was 6 B km from Earth. Not exactly close encounters

Query 19 of 20

User Query

The image of the Link rock outcrop and the image of the first drill hole at John Klein together form a two-stage visual argument for habitability. Using both images alongside the text describing what each sample revealed, explain what morphological features visible in the Link image constitute evidence for past water flow, why that visual evidence was scientifically necessary but not sufficient for a habitability claim, and how the drill hole image at John Klein represents the specific instrumental step that closed the evidentiary gap — including the two numerical results that no surface image could have produced.

Retrieved Text Context

[1] investigate an ancient river and fan system identified in The rover analyzed its first scoops of soil on the way to Yellowknife Bay. Once there, it collected the first samples of material ever drilled from rocks on Mars. Analysis of the first drilled sample, from a rock target called “John Klein,” provided the evidence of conditions favorable for life in Mars’ early history: geological and mineralogical evidence for sustained liquid water, other key elemental ingredients for life, a chemical energy source, and water not too acidic or too salty. On a subsequent drill sample, Curiosity A rock outcrop called Link shows signs of being formed by the deposition of water. was able to accomplish a first for measurements on another planet: determining the age of the rock. The lowered consideration of more than 30 Martian locations by measurements showed that the drilled material was more than 100 scientists participating in a series of open 4.2 billion years old and yet had been exposed at the workshops. The selection process benefited from examining surface for only 80 million years. ing candidate sites with NASA’s Mars Reconnaissance Orbiter and earlier orbiters, and from the rover mission’s In July 2013, Curiosity finished investigations in the Yellowknife Bay area and began a southwestward trek to 12 kilometers (12 miles) long. That precision, about a five-fold improvement on earlier Mars landings, made sites eligible this main destination in September 2014. In the low lay-out that would otherwise be excluded for encompassing the base of Mount Sharp during the rover’s extended mission, nearby unsuitable terrain. The Gale Crater landing site is so researchers anticipate finding further evidence about habitability. (Source: mars-science-laboratory.pdf, page 2)

[2] iterated many design elements from them, including six-wheel In the first few weeks after landing, images from the rover showed that Curiosity touched down right in an area where water once coursed vigorously over the surface. The evidence for stream flow was in rounded pebbles

mixed with hardened sand in conglomerate rocks at and near the landing site. Analysis of Mars' atmospheric composition early in the mission provided evidence that the planet has lost much of its original atmosphere by a process favoring loss from the top of the atmosphere rather than interaction with the surface.

In the initial months of the surface mission, the rover team drove Curiosity eastward toward "Yellowknife Bay" to Curiosity's first sample drilling, at a rock called "John Klein." Mars Science Laboratory/Curiosity

2

NASA Facts

(Source: mars-science-laboratory.pdf, page 2)

[3] capability of landing within a target area only about 20 kilometers (12 miles) long. That precision, about a five-fold improvement on earlier Mars landings, made sites eligible this main destination in September 2014. In the low lay-out that would otherwise be excluded for encompassing areas of Mount Sharp during the rover's extended mission, nearby unsuitable terrain. The Gale Crater landing site is so researchers anticipate finding further evidence about habitable past environments and about the evolution of the Martian environment from a wetter past to a drier present. this improved precision.

Science findings began months before landing. Measurements that Curiosity made of natural radiation levels during the flight from Earth to Mars will help NASA design for as-Rovers, Spirit and Opportunity, launched in 2003. It inherited many design elements from them, including six-wheel

In the first few weeks after landing, images from th

Retrieved Images



A rock outcrop called Link shows signs of being formed by the deposition of water. lowed consideration of more than 30 Martian locations by more than 100 scientists participating in a series of open workshops. The selection process benefited from examin

Query 20 of 20

User Query

The introduction page shows two images: the Pillars of Creation in M16 and astronaut Mike Good repairing the Space Telescope Imaging Spectrograph during the final servicing mission in 2009. Using both images alongside the introduction text citing Hubble's productivity metrics — over one million observations, more than 17,000 peer-reviewed publications, 900,000 citations growing at over 150 per day — explain how the Pillars image and the servicing image each represent a structurally distinct prerequisite for achieving that level of scientific output, and why neither prerequisite alone would have been sufficient without the two specific observational properties of Hubble's orbital position described in the text.

Retrieved Text Context

[1] Hubble's discoveries and memorable photos have also reinvigorated the public's interest in astronomy. Along with pictures of the telescope and the astronauts who launched and serviced it during six space shuttle missions, certain memorable science images have become cultural icons. They appear regularly on book covers, musical albums, clothing, TV shows, movies, and even ecclesiastical stained-glass windows.

This booklet features thirteen representative topics with eye-catching images and thought-provoking discoveries. Selecting this set from the thousands of amazing Hubble observations was difficult, but these serve to highlight some of Hubble's greatest scientific achievements to date. Additional images and information can be found at <http://nasa.gov/hubble> and <http://hubblesite.org>.

Right: Astronaut Mike Good maneuvers to repair the Space Telescope Imaging Spectrograph during the final

Spectrograph during the final Hubble

Hubble servicing mission in May 2009. Periodic upgrades servicing mission in May 2009. Periodic upgrades

have kept the telescope equipped with state-of-the-art instruments, which have given have kept the telescope equipped with state-of-the-art instruments, which have given astronomers increasingly better views of the cosmos.

astronomers increasingly better views of the cosmos.

Cover image: Veil Nebula

Cover image: Veil Nebula

(Source: [highlights_of_hubbles_exploration_of_the_universe.pdf](#), page 2)

[2] objects as nearby and small as the collision of asteroids to distant star-forming galaxies that date back to when the universe was only three percent of its current age. In fact, Hubble observations have played a key role in discovering and characterizing the mysterious dark energy that now appears to permeate space. Results like these have changed our fundamental understanding of the cosmos.

Well into its third operational decade, Hubble is still extremely productive. The orbiting telescope has taken over a million observations and provided data that astronomers have used to write more than 17,000 peer-reviewed scientific publications on a broad range of topics, from planet formation to gigantic black holes. These papers have been referenced in other publications approximately 900,000 times, and this total increases, on average, by more than

150 per day.

Every current astronomy textbook includes contributions from the observatory. Today's college undergraduates have not known

a time in their lives when astronomers were not actively making discoveries with Hubble data.

One of

One of Hubble

Hubble's most iconic images is of this portion of the Eagle Nebula

's most iconic images is of this portion of the Eagle Nebula

(M16). Dubbed the "Pillars of Creation," it shows star birth. Imbedded in

(M16). Dubbed the "Pillars of Creation," it shows star birth. Imbedded in

the tips of the finger-like protrusions at the top of the columns are dense,

the tips of the finger-like protrusions at the top of the columns are dense,

gaseous globules within which stars are being born.

gaseous globules within which stars are being born.

(Source: highlights_of_hubbles_exploration_of_the_universe.pdf, page 2)

[3] First photograph of the Earth-Moon system in a single frame (left), taken by Voyager 1 on September 18, 1977. Voyager spacecraft as it would appear in space (right). Images courtesy of NASA.

On their way to Jupiter, both Voyagers successfully passed through the Asteroid Belt without incident –

two Pioneer spacecraft had made the crossing safely in 1973 and 1974 as pathfinders, proving that it

was safe to do so. Voyager 1 arrived first, passing 217,000 miles above the giant planet's cloud tops on


March 5, 1979. Two months before the close encounter, Voyager 1 began taking photographs of Jupiter

and its moons and gathering other scientific data about the planet and its environment, a process that

continued for about a month after the fly-by. Voyager 2 followed not too far behind, beginning

scie

Retrieved Images



Left: Hubble's Ultra Deep Field is one of the most distant looks into space ever. The cumulative exposure time needed to capture the image was about a million seconds (11 days). Right: A sample of the faintest and farthest galaxies in the Hubble Ultra Deep Field shows that they were irregularly shaped and frequently interacting in the distant past.