

DATA70121: Statistics and Machine Learning 1

Lecture 5: Regression I

Diego Perez Ruiz
Department of Social Statistics
School of Social Sciences
University of Manchester

26 October 2023

this lecture

modelling numerical and categorical variables

we will be looking at some models

- ▶ simple linear regression
numerical response and one predictor
- ▶ multiple linear regression
numerical response and multiple predictors
- ▶ generalized linear models (glm)
categorical response and predictor(s) that may be non-linear or
have complicated dependence structures
 - ▶ logistic regression
 - ▶ Poisson regression

introduction and notation

► Y

the dependent variable

or outcome

or regressand

or left-hand-side variable

or response

or endogenous variable

► X

the independent variable

or explanatory variable

or regressor

or right-hand-side variable

or treatment

or predictor

or covariate

or exogenous variable

introduction and notation

► Y

the dependent variable

or outcome

or regressand

or left-hand-side variable

or response

or endogenous variable

► X

the independent variable

or explanatory variable

or regressor

or right-hand-side variable

or treatment

or predictor

or covariate

or exogenous variable

generally our goal is to understand how Y varies as a function of X :

$$Y = f(X) + \text{error}$$

why regression?

roughly, we can distinguish between three uses for regression analysis:

1. **description** - parsimonious summary of the data
2. **prediction/estimation/inference** - learn about parameters of the joint distribution of the data
3. **causal inference** - evaluate counterfactuals

how regression?

- ▶ regression quantifies how an outcome variable Y varies as a function of one or more predictor variables X
- ▶ the common idea: **conditioning** on X
- ▶ goal is to characterize $f(Y|X)$, the conditional probability distribution of Y for different levels of X
- ▶ instead of modelling the whole conditional density of Y given X , in regression we usually only model **the conditional mean** of Y given X :

$$E[Y|X = x]$$

- ▶ our key goal is to approximate **the conditional expectation function** $E[Y|X]$, which summarizes how the average of Y varies across all possible levels of X (also called **the population regression function**)
- ▶ once we have estimated $E[Y|X]$, we can use it for **prediction** and/or **causal inference**

linear regression

- ▶ linear regression works by assuming linear parametric form for the conditional expectation function:

$$E[Y|X] = \beta_0 + X\beta_1$$

- ▶ conditional expectation defined by only two coefficients which are estimated from the data:
 - ▶ β_0 is called the intercept or constant
 - ▶ β_1 is called the slope coefficient
- ▶ notice that the linear functional form imposes a constant slope

interpreting the regression intercept and slope

$$E[Y|X] = \beta_0 + X\beta_1$$

when we model the regression function as a line, we can interpret the parameters of the line in appealing ways:

► **intercept**

the average outcome among units with $X = 0$ is β_0

$$E[Y|X = 0] = \beta_0 + \beta_1(0) = \beta_0$$

► **slope**

a one-unit change in X is associated with a β_1 change in Y

$$\begin{aligned} E[Y|X = x + 1] - E[Y|X = x] &= (\beta_0 + \beta_1(x + 1)) - (\beta_0 + \beta_1x) \\ &= \beta_0 + \beta_1x + \beta_1 - \beta_0 - \beta_1x \\ &= \beta_1 \end{aligned}$$

linear regression

note. the model will not always be a good fit for the data
(even though it really wants to be)



linear regression always returns a **line** regardless of the data

simple linear regression

numerical response and one predictor

correlation


- ▶ strength of the **linear** association between two variables (X, Y)
- ▶ the population correlation coefficient is denoted ρ
- ▶ the sample correlation coefficient is denoted r (or R)
- ▶ values between -1 (perfect negative) and +1 (perfect positive)
- ▶ value 0 indicate no linear association

simple linear regression

numerical response and one predictor

correlation

- ▶ strength of the **linear** association between two variables (X, Y)
- ▶ the population correlation coefficient is denoted ρ
- ▶ the sample correlation coefficient is denoted r (or R)
- ▶ values between -1 (perfect negative) and +1 (perfect positive)
- ▶ value 0 indicate no linear association



from <http://en.wikipedia.org/wiki/Correlation>

defining correlation

covariance between X and Y

covariance

- ▶ a generalization of variance to two random variables

$$\text{Cov}(X, Y) = \frac{1}{n} \sum_{i=1}^n (x_i - \mu_X)(y_i - \mu_Y)$$

- ▶ is not a measure of uncertainty
- ▶ a measure of how the two variables vary in relation to each other

defining correlation

covariance between X and Y

covariance

- ▶ a generalization of variance to two random variables

$$\text{Cov}(X, Y) = \frac{1}{n} \sum_{i=1}^n (x_i - \mu_X)(y_i - \mu_Y)$$

- ▶ is not a measure of uncertainty
- ▶ a measure of how the two variables vary in relation to each other

properties of covariance:

- ▶ $\text{Cov}(X, X) = \text{Var}(X, X)$
- ▶ $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶ $\text{Cov}(X, Y) = 0$ if X and Y are independent
- ▶ $\text{Cov}(X, c) = 0$

defining correlation

the Pearson correlation coefficient is given by

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y} \quad -1 \leq \rho(X, Y) \leq 1$$

- ▶ X and Y are independent $\Rightarrow \text{Cov}(X, Y) = \rho(X, Y) = 0$
- ▶ $\text{Cov}(X, Y) = \rho(X, Y) = 0 \not\Rightarrow X$ and Y are independent


defining correlation

the Pearson correlation coefficient is given by

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y} \quad -1 \leq \rho(X, Y) \leq 1$$



- ▶ X and Y are independent $\Rightarrow \text{Cov}(X, Y) = \rho(X, Y) = 0$
- ▶ $\text{Cov}(X, Y) = \rho(X, Y) = 0 \not\Rightarrow X$ and Y are independent

which scatterplot shows the strongest correlation?




Source: Wikipedia

correlation \neq causation




Source: <http://tylervigen.com/>

correlation \neq causation




finding the best linear fit

example




finding the best linear fit

example



finding the best linear fit

residuals



residual is the difference between the observed and predicted y

$$e_i = y_i - \hat{y}_i$$

ordinary least squares (ols)

we wish to find the line that minimises the sum of squared residuals

$$e_1^2 + e_2^2 + \cdots + e_n^2$$

ordinary least squares (ols)

we wish to find the line that minimises the sum of squared residuals

$$e_1^2 + e_2^2 + \cdots + e_n^2$$

simple linear regression model has the following regression line

$$Y_i = \beta_0 + \beta_1 X_i$$

which is by ols estimated to

$$\hat{y}_i = b_0 + b_1 x_i$$

ordinary least squares (ols)

we wish to find the line that minimises the sum of squared residuals

$$e_1^2 + e_2^2 + \dots + e_n^2$$

simple linear regression model has the following regression line

$$Y_i = \beta_0 + \beta_1 X_i$$

which is by ols estimated to

$$\hat{y}_i = b_0 + b_1 x_i$$

intercept

- ▶ parameter β_0
- ▶ point estimate b_0

slope

- ▶ parameter β_1
- ▶ point estimate b_1

ordinary least squares (ols)

we wish to find the line that minimises the sum of squared residuals

$$e_1^2 + e_2^2 + \dots + e_n^2$$

simple linear regression model has the following regression line

$$Y_i = \beta_0 + \beta_1 X_i$$

which is by ols estimated to

$$\hat{y}_i = b_0 + b_1 x_i$$

intercept

- ▶ parameter β_0
- ▶ point estimate b_0

slope

- ▶ parameter β_1
- ▶ point estimate b_1

for simple linear regression it is easy to calculate b_0 and b_1 by hand

ordinary least squares (ols)

the slope of the regression is estimated by

$$b_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} = \frac{\text{sample covariance between } X \text{ and } Y}{\text{sample variance of } X}$$
$$= \frac{s_y}{s_x} r$$

where


$$r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}}$$

and can be used to estimate the intercept b_0 by

$$b_0 = \bar{y} - b_1 \bar{x}$$


ordinary least squares (ols)

ols is an estimator:



ordinary least squares (ols)


ols is an estimator:




- ▶ just like sample mean, sample difference in means, or sample variance
- ▶ it has a sampling distribution, with a sampling variance/standard error, etc.

sampling distribution of ols

Sampling distribution of intercepts




Sampling distribution of slopes



the estimated slopes and intercepts vary from sample to sample, but on average the lines looks about right

example



the estimated model is

$$\hat{y}_i = 101.75 + 0.14x_i$$

what's the interpretation?

assessing the fit of the model

coefficient of determination

- ▶ denoted r^2 (or R^2) where $0 \leq r^2 \leq 1$
- ▶ what % of variability in response variable is explained by model (so optimal scenario is $r^2 = 1$)
- ▶ remainder is explained by variables not included in model

assessing the fit of the model

coefficient of determination

- ▶ denoted r^2 (or R^2) where $0 \leq r^2 \leq 1$
- ▶ what % of variability in response variable is explained by model (so optimal scenario is $r^2 = 1$)
- ▶ remainder is explained by variables not included in model

for simple linear regression

determination coefficient is the square of correlation coefficient

assessing the fit of the model

coefficient of determination

- ▶ denoted r^2 (or R^2) where $0 \leq r^2 \leq 1$
- ▶ what % of variability in response variable is explained by model (so optimal scenario is $r^2 = 1$)
- ▶ remainder is explained by variables not included in model

for simple linear regression

determination coefficient is the square of correlation coefficient

example

the correlation between birth weight and mom's weight is $r = 0.156$

then the determination coefficient is $r^2 = 0.156^2 = 0.024$

output for example

baby weight predicted by mom's weight

$$\hat{y}_i = 101.75 + 0.14x_i$$

Call:

```
lm(formula = birth.weight ~ mom.weight)
```

Residuals:

Min	1Q	Median	3Q	Max
-66.065	-10.943	0.333	11.048	56.075

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	101.74736	3.32162	30.632	< 2e-16 ***
mom.weight	0.13798	0.02552	5.406	7.82e-08 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 18.12 on 1170 degrees of freedom

Multiple R-squared: 0.02437, Adjusted R-squared: 0.02353

F-statistic: 29.22 on 1 and 1170 DF, p-value: 7.819e-08

output for example

baby weight predicted by mom's weight

$$\hat{y}_i = 101.75 + 0.14x_i$$

Call:

lm(formula = birth.weight ~ mom.weight)

Residuals:

Min	1Q	Median	3Q	Max
-66.065	-10.943	0.333	11.048	56.075

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	101.74736	3.32162	30.632	< 2e-16 ***
mom.weight	0.13798	0.02552	5.406	7.82e-08 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 18.12 on 1170 degrees of freedom

Multiple R-squared: 0.02437, Adjusted R-squared: 0.02353

F-statistic: 29.22 on 1 and 1170 DF, p-value: 7.819e-08

output for example

baby weight predicted by mom's weight

$$\hat{y}_i = 101.75 + 0.14x_i$$

Call:

```
lm(formula = birth.weight ~ mom.weight)
```

Residuals:

Min	1Q	Median	3Q	Max
-66.065	-10.943	0.333	11.048	56.075

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	101.74736	3.32162	30.632	< 2e-16	***
mom.weight	0.13798	0.02552	5.406	7.82e-08	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 18.12 on 1170 degrees of freedom

Multiple R-squared: 0.02437, Adjusted R-squared: 0.02353

F-statistic: 29.22 on 1 and 1170 DF, p-value: 7.819e-08

output for example

baby weight predicted by mom's weight

$$\hat{y}_i = 101.75 + 0.14x_i$$

Call:

```
lm(formula = birth.weight ~ mom.weight)
```

Residuals:

Min	1Q	Median	3Q	Max
-66.065	-10.943	0.333	11.048	56.075

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	101.74736	3.32162	30.632	< 2e-16 ***
mom.weight	0.13798	0.02552	5.406	7.82e-08 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 18.12 on 1170 degrees of freedom

Multiple R-squared: 0.02437, Adjusted R-squared: 0.02353

F-statistic: 29.22 on 1 and 1170 DF, p-value: 7.819e-08

coefficient of determination



$$R^2 = 1 - \frac{RSS}{TSS} \quad RSS = \sum_{i=1}^n e_i^2 \quad TSS = \sum_{i=1}^n (y_i - \bar{y})^2$$

- ▶ Caveat: when a covariate is added, R^2 increases

coefficient of determination



$$R^2 = 1 - \frac{RSS}{TSS} \quad RSS = \sum_{i=1}^n e_i^2 \quad TSS = \sum_{i=1}^n (y_i - \bar{y})^2$$

- ▶ Caveat: when a covariate is added, R^2 increases
- ▶ Adjusted R^2 , or R_{adj}^2

$$R_{adj}^2 = 1 - (1 - R^2) \frac{n - 1}{n - p - 1}$$

- ▶ p - number of covariates in the model

Part II

multiple linear regression

same underlying idea as before but with multiple predictors

$$Y_i = \beta_0 + \beta_1 X_{i1} + \beta_2 X_{i2} + \cdots + \beta_p X_{ip}$$

which is by ols estimated to (requires matrix algebra)

$$\hat{y}_i = b_0 + b_1 x_{i1} + b_2 x_{i2} + \cdots + b_p x_{ip}$$

multiple linear regression

same underlying idea as before but with multiple predictors

$$Y_i = \beta_0 + \beta_1 X_{i1} + \beta_2 X_{i2} + \cdots + \beta_p X_{ip}$$

which is by ols estimated to (requires matrix algebra)

$$\hat{y}_i = b_0 + b_1 x_{i1} + b_2 x_{i2} + \cdots + b_p x_{ip}$$

```
Call:
lm(formula = birth.weight ~ gestation + mom.smokes)

Residuals:
    Min       1Q   Median       3Q      Max
-50.553 -10.855  -0.178  10.013  50.495

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept) -13.64855    8.69303  -1.570   0.117
gestation     0.48809    0.03095  15.771 <2e-16 ***
mom.smokes    -8.17175    0.96916  -8.432 <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 16.15 on 1169 degrees of freedom
Multiple R-squared:  0.2259,    Adjusted R-squared:  0.2246
F-statistic: 170.6 on 2 and 1169 DF,  p-value: < 2.2e-16
```

for interpretation we must use **'all else held constant'**

conditions for inference


model assumptions must be met:

- ▶ linearity
- ▶ constant residual variance (homoscedasticity and no autocorrelation)
- ▶ approximately normally distributed residuals

conditions for inference

model assumptions must be met:


- ▶ **linearity**
- ▶ **constant residual variance (homoscedasticity and no autocorrelation)**
- ▶ approximately normally distributed residuals



conditions for inference

model assumptions must be met:

- ▶ linearity
- ▶ constant residual variance (homoscedasticity and no autocorrelation)
- ▶ **approximately normally distributed residuals**



conditions for inference

model assumptions must be met:

- ▶ linearity
- ▶ constant residual variance (homoscedasticity and no autocorrelation)
- ▶ approximately normally distributed residuals

and most importantly

- ▶ the response is continuous variable that is normally distributed
...but what if it isn't?

generalized linear models (glm)

a glm has the following three components:

1. a probability distribution describing the response variable Y that should belong to **the exponential family**:
 - ▶ normal
 - ▶ binomial
 - ▶ Poisson
 - ▶ \vdots
2. a linear function of the regressors, called **linear predictor**

$$\eta_i = \beta_0 + \beta_1 X_{1i} + \dots + \beta_k X_{ki}$$

on which the expected value μ_i of Y_i depend

3. a invertible **link function** which transforms the expectation of the response to the linear predictor

$$g(\mu_i) = \eta_i \quad \text{or} \quad g^{-1}(\eta_i) = \mu_i$$

logistic regression

- ▶ assume a binomial distribution produced the outcome variable
- ▶ want to model $\mu = p$ (probability of success) given set of predictors

the logistic model is specified when a link function connects η to p

logistic regression

- ▶ assume a binomial distribution produced the outcome variable
- ▶ want to model $\mu = p$ (probability of success) given set of predictors

the logistic model is specified when a link function connects η to p

the most commonly used is the logit function:

$$\text{logit}(p) = \log \left(\frac{p}{1-p} \right) \quad \text{for } 0 \leq p \leq 1$$

the logit function takes a value between 0 and 1 and maps it to a value between $-\infty$ and ∞

the inverse of the logit function:

$$g^{-1}(x) = \frac{\exp(x)}{1 + \exp(x)} = \frac{1}{1 + \exp(-x)}$$

the inverse logit function takes a value between $-\infty$ and ∞ and maps it to a value between 0 and 1

logistic regression

the three glm criteria give us

1. $y_i \sim \text{Binomial}(p_i, n)$
2. $\eta_i = \beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki}$
3. $\text{logit}(p_i) = \eta_i$

which gives us

$$\log \left(\frac{p_i}{1 - p_i} \right) = \beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki}$$

and

$$p_i = \frac{\exp(\beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki})}{1 + \exp(\beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki})}$$

logistic regression

the three glm criteria give us

1. $y_i \sim \text{Binomial}(p_i, n)$
2. $\eta_i = \beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki}$
3. $\text{logit}(p_i) = \eta_i$

which gives us

$$\log \left(\frac{p_i}{1 - p_i} \right) = \beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki}$$

and

$$p_i = \frac{\exp(\beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki})}{1 + \exp(\beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki})}$$

logistic regression

the three glm criteria give us

1. $y_i \sim \text{Binomial}(p_i, n)$
2. $\eta_i = \beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki}$
3. $\text{logit}(p_i) = \eta_i$

which gives us

$$\log \left(\frac{p_i}{1 - p_i} \right) = \beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki}$$

and

$$p_i = \frac{\exp(\beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki})}{1 + \exp(\beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki})}$$

logistic regression

odds

odds are another way to quantify the probability of an event

logistic regression

odds

odds are another way to quantify the probability of an event

for some event E

$$\text{odds}(E) = \frac{P(E)}{P(E^c)} = \frac{P(E)}{1 - P(E)}$$

logistic regression

odds

odds are another way to quantify the probability of an event

for some event E

$$\text{odds}(E) = \frac{P(E)}{P(E^c)} = \frac{P(E)}{1 - P(E)}$$

if we are told the odds of E are x to y , then

$$\text{odds}(E) = \frac{x}{y} = \frac{x/(x+y)}{y/(x+y)}$$

$$\implies P(E) = x/(x+y), \quad P(E^c) = y/(x+y)$$

logistic regression

odds

odds are another way to quantify the probability of an event

for some event E

$$\text{odds}(E) = \frac{P(E)}{P(E^c)} = \frac{P(E)}{1 - P(E)}$$

if we are told the odds of E are x to y , then

$$\text{odds}(E) = \frac{x}{y} = \frac{x/(x+y)}{y/(x+y)}$$

$$\implies P(E) = x/(x+y), \quad P(E^c) = y/(x+y)$$

useful when interpreting coefficient estimates of a logistic regression

logistic regression

example. lab 5

we want to create a spam filter based on

- ▶ 3921 observations/emails
- ▶ properties of the emails (more details during class)

logistic regression

example. lab 5

we want to create a spam filter based on

- ▶ 3921 observations/emails
- ▶ properties of the emails (more details during class)

simple model: use binary predictor 'winner'

$$\log \left(\frac{p_i}{1 - p_i} \right) = \beta_0 + \beta_{1i} \times \text{winner}$$

which is estimated to

$$\log \left(\frac{\hat{p}_i}{1 - \hat{p}_i} \right) = -2.31 + 1.53 \times \text{winner}$$

logistic regression

example. lab 5

$$\log \left(\frac{\hat{p}_i}{1 - \hat{p}_i} \right) = -2.31 + 1.53 \times \text{winner}$$

logistic regression

example. lab 5

$$\log \left(\frac{\hat{p}_i}{1 - \hat{p}_i} \right) = -2.31 + 1.53 \times \text{winner}$$

the odds of an email being spam if 'winner' = 0:

$$\frac{\hat{p}_i}{1 - \hat{p}_i} = \exp(-2.31) = 0.10$$

logistic regression

example. lab 5

$$\log \left(\frac{\hat{p}_i}{1 - \hat{p}_i} \right) = -2.31 + 1.53 \times \text{winner}$$

the odds of an email being spam if 'winner' = 0:

$$\frac{\hat{p}_i}{1 - \hat{p}_i} = \exp(-2.31) = 0.10$$

the probability of an email being spam if 'winner' = 0:

$$\hat{p}_i = \frac{0.1}{1.1} = 0.09$$

logistic regression

example. lab 5

$$\log \left(\frac{\hat{p}_i}{1 - \hat{p}_i} \right) = -2.31 + 1.53 \times \text{winner}$$

the odds of an email being spam if 'winner' = 0:

$$\frac{\hat{p}_i}{1 - \hat{p}_i} = \exp(-2.31) = 0.10$$

the probability of an email being spam if 'winner' = 0:

$$\hat{p}_i = \frac{0.1}{1.1} = 0.09$$

what are the odds and probability if 'winner' = 1?

logistic regression

example. lab 5

$$\log \left(\frac{\hat{p}_i}{1 - \hat{p}_i} \right) = -2.31 + 1.53 \times \text{winner}$$

the odds of an email being spam if 'winner' = 0:

$$\frac{\hat{p}_i}{1 - \hat{p}_i} = \exp(-2.31) = 0.10$$

the probability of an email being spam if 'winner' = 0:

$$\hat{p}_i = \frac{0.1}{1.1} = 0.09$$

what are the odds and probability if 'winner' = 1?

answer: odds is 0.45 and probability is 0.31

output for spam filter example

```
Call:
glm(formula = spam ~ winner, family = binomial, data = email)

Deviance Residuals:
    Min       1Q   Median       3Q      Max
-0.8657  -0.4342  -0.4342  -0.4342   2.1947

Coefficients:
              Estimate Std. Error z value Pr(>|z|)
(Intercept) -2.31405     0.05627 -41.121  < 2e-16 ***
winneryes    1.52559     0.27549   5.538 3.06e-08 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

    Null deviance: 2437.2  on 3920  degrees of freedom
Residual deviance: 2412.7  on 3919  degrees of freedom
AIC: 2416.7

Number of Fisher Scoring iterations: 5
```

output for spam filter example

Call:

glm(formula = spam ~ winner, family = binomial, data = email)

Deviance Residuals:

Min	1Q	Median	3Q	Max
-0.8657	-0.4342	-0.4342	-0.4342	2.1947

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-2.31405	0.05627	-41.121	< 2e-16 ***
winneryes	1.52559	0.27549	5.538	3.06e-08 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

Null deviance: 2437.2 on 3920 degrees of freedom
Residual deviance: 2412.7 on 3919 degrees of freedom
AIC: 2416.7

Number of Fisher Scoring iterations: 5

output for spam filter example

Call:

glm(formula = spam ~ winner, family = binomial, data = email)

Deviance Residuals:

Min	1Q	Median	3Q	Max
-0.8657	-0.4342	-0.4342	-0.4342	2.1947

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
<u>(Intercept)</u>	<u>-2.31405</u>	0.05627	-41.121	< 2e-16 ***
<u>winnersyes</u>	<u>1.52559</u>	0.27549	5.538	3.06e-08 ***


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

Null deviance: 2437.2 on 3920 degrees of freedom
Residual deviance: 2412.7 on 3919 degrees of freedom
AIC: 2416.7


Number of Fisher Scoring iterations: 5

output for spam filter example




linear versus logistic regression

Linear Regression



Logistic Regression



Poisson regression

suitable for when the response is count data
the three glm criteria give us

1. $y_i \sim \text{Poisson}(\mu_i)$
2. $\eta_i = \beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki}$
3. $\log(\mu_i) = \eta_i$

which gives us

$$\log(\mu_i) = \beta_0 + \beta_1 X_{1i} + \cdots + \beta_k X_{ki}$$

and

$$\mu_i = \exp(\beta_0 + \beta_1 X_{1i} + \cdots + \beta_p X_{ki})$$

Poisson regression

example in r

y_i is number of spam emails received, $y_i \sim \text{Poisson}(\mu_i)$

predict y_i using total nr of emails received

$$\log(\mu_i) = \beta_0 + \beta_1 \times \text{nr.emails}$$


Poisson regression

example in r

y_i is number of spam emails received, $y_i \sim \text{Poisson}(\mu_i)$

predict y_i using total nr of emails received

$$\log(\mu_i) = \beta_0 + \beta_1 \times \text{nr.emails}$$



Poisson regression

example in r

```
Call:
glm(formula = nr.spam ~ nr.emails, family = poisson, data = MyData)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-2.75498	-0.98292	-0.07578	0.53401	2.35312

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-0.253771	0.143496	-1.768	0.077 .
nr.emails	0.029698	0.001889	15.724	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for poisson family taken to be 1)

Null deviance: 424.24 on 100 degrees of freedom
Residual deviance: 123.01 on 99 degrees of freedom
AIC: 417.35

Number of Fisher Scoring iterations: 5

Poisson regression

example in r

Call:

```
glm(formula = nr.spam ~ nr.emails, family = poisson, data = MyData)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-2.75498	-0.98292	-0.07578	0.53401	2.35312

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-0.253771	0.143496	-1.768	0.077 .
nr.emails	0.029698	0.001889	15.724	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for poisson family taken to be 1)

Null deviance: 424.24 on 100 degrees of freedom
Residual deviance: 123.01 on 99 degrees of freedom
AIC: 417.35

Number of Fisher Scoring iterations: 5

Poisson regression

example in r

Call:

```
glm(formula = nr.spam ~ nr.emails, family = poisson, data = MyData)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-2.75498	-0.98292	-0.07578	0.53401	2.35312

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
<u>(Intercept)</u>	<u>-0.253771</u>	0.143496	-1.768	0.077 .
<u>nr.emails</u>	<u>0.029698</u>	0.001889	15.724	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for poisson family taken to be 1)

Null deviance: 424.24 on 100 degrees of freedom
Residual deviance: 123.01 on 99 degrees of freedom
AIC: 417.35

Number of Fisher Scoring iterations: 5

Poisson regression

example in r

estimated model is $\hat{\mu}_i = \exp(-0.25 + 0.03 \times \text{nr.emails})$

Poisson regression

example in r

estimated model is $\hat{\mu}_i = \exp(-0.25 + 0.03 \times \text{nr.emails})$

assume you receive 30 emails, how many are expected to be spam?

$$\hat{\mu}_i = \exp(-0.25 + 0.03 \times 30) = 1.91 \approx 2$$


Poisson regression

example in r

estimated model is $\hat{\mu}_i = \exp(-0.25 + 0.03 \times \text{nr.emails})$

assume you receive 30 emails, how many are expected to be spam?

$$\hat{\mu}_i = \exp(-0.25 + 0.03 \times 30) = 1.91 \approx 2$$



Poisson regression

example in r

- ▶ in Poisson distribution, $E(X) = Var(X) = \mu$
- ▶ problem of overdispersion $E(X) < Var(X)$

Poisson regression

example in r

- ▶ in Poisson distribution, $E(X) = Var(X) = \mu$
- ▶ problem of overdispersion $E(X) < Var(X)$
- ▶ `family=quasipoisson`

Poisson regression

example in r

- ▶ in Poisson distribution, $E(X) = Var(X) = \mu$
- ▶ problem of overdispersion $E(X) < Var(X)$
- ▶ `family=quasipoisson`
- ▶ negative binomial distribution

different link functions

link name	$\eta_i = g(\mu_i)$	$\mu_i = g^{-1}(\eta_i)$
identity	μ_i	η_i
log	$\log(\mu_i)$	$\exp(\eta_i)$
logit	$\log\left(\frac{\mu_i}{1-\mu_i}\right)$	$\frac{1}{1+\exp(-\eta_i)}$
\vdots	\vdots	\vdots

different link functions

link name	$\eta_i = g(\mu_i)$	$\mu_i = g^{-1}(\eta_i)$
identity	μ_i	η_i
log	$\log(\mu_i)$	$\exp(\eta_i)$
logit	$\log\left(\frac{\mu_i}{1-\mu_i}\right)$	$\frac{1}{1+\exp(-\eta_i)}$
\vdots	\vdots	\vdots

family	canonical link	range of Y_i
Gaussian (normal)	identity	$(-\infty, \infty)$
binomial	logit	$\{0, 1\}$
Poisson	log	$0, 1, 2, \dots$
\vdots	\vdots	\vdots

estimation of glm

- ▶ glms are estimated via **maximum likelihood estimation** (mle)
- ▶ **most likely** values of parameters **given the data** we observed
- ▶ applied to linear regression \implies ols=mle

estimation of glm

- ▶ glms are estimated via **maximum likelihood estimation** (mle)
- ▶ **most likely** values of parameters **given the data** we observed
- ▶ applied to linear regression \implies `ols=mle`

model fit

conditions for inference are the same as for linear regression
(recall that linear regression is just a special case of glm)

goodness of fit measures:

- ▶ Pearson chi-square statistic
- ▶ deviance: $-2(\log L_m - \log L_s)$
`(anova(model1, test=' 'chisq' '))`
- ▶ likelihood ratio tests (`epicalc::lrtest(model1, model2)`)

estimation of glm

- ▶ glms are estimated via **maximum likelihood estimation** (mle)
- ▶ **most likely** values of parameters **given the data** we observed
- ▶ applied to linear regression \Rightarrow ols=mle

model fit

conditions for inference are the same as for linear regression
(recall that linear regression is just a special case of glm)

goodness of fit measures:

- ▶ Pearson chi-square statistic
- ▶ deviance: $-2(\log L_m - \log L_S)$

```
(anova(modell, test=' 'chisq' '))
```

	NULL		Df	Deviance	Resid.	Df	Resid.	Dev	Pr(>Chi)
winner	1	24.506	3920	2437.2	3919	2412.7	7.41e-07	***	

- ▶ likelihood ratio tests (`epicalc::lrtest(modell, model2)`)

estimation of glm

- ▶ glms are estimated via **maximum likelihood estimation** (mle)
- ▶ **most likely** values of parameters **given the data** we observed
- ▶ applied to linear regression \implies `ols=mle`

model fit

conditions for inference are the same as for linear regression
(recall that linear regression is just a special case of glm)

goodness of fit measures:

- ▶ Pearson chi-square statistic
- ▶ deviance: $-2(\log L_m - \log L_s)$
`(anova(modell, test=' 'chisq' '))`
- ▶ likelihood ratio tests (`epicalc::lrtest(modell, model2)`)
- ▶ Akaike Information Criterion (AIC) – Lecture 9

reading


Agresti A., 2018, Statistical Methods for the Social Sciences, Fifth Edition, **Chapters 9, 11, 14.4, 15.1**

link to the book via Manchester library

reading

Agresti A., 2018, Statistical Methods for the Social Sciences, Fifth Edition, **Chapters 9, 11, 14.4, 15.1**

link to the book via Manchester library




Source: Wikipedia

reading

Agresti A., 2018, Statistical Methods for the Social Sciences, Fifth Edition, **Chapters 9, 11, 14.4, 15.1**

link to the book via Manchester library



Source: Wikipedia

$$r = 0. - 0.06$$