

THE SHARP SOBOLEV INEQUALITY AND THE BANCHOFF-POHL INEQUALITY ON SURFACES

RALPH HOWARD
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF SOUTH CAROLINA
COLUMBIA, S.C. 29208
HOWARD@MATH.SC.EDU

ABSTRACT. Let (M, g) be a complete two dimensional simply connected Riemannian manifold with Gaussian curvature $K \leq -1$. If f is a compactly supported function of bounded variation on M then f satisfies the Sobolev inequality

$$4\pi \int_M f^2 dt + \left(\int_M |f| dA \right)^2 \leq \left(\int_M \|\nabla f\| dA \right)^2.$$

Conversely letting f be the characteristic function of a domain $D \subset M$ recovers the sharp form $4\pi A(D) + A(D)^2 \leq L(\partial D)^2$ of the isoperimetric inequality for simply connected surfaces with $K \leq -1$. Therefore this is the Sobolev inequality “equivalent” to the isoperimetric inequality for this class of surfaces. This is a special case of a result that gives the equivalence of more general isoperimetric inequalities and Sobolev inequalities on surfaces.

Under the same assumptions on (M, g) if $c: [a, b] \rightarrow M$ is a closed curve and $w_c(x)$ is the winding number of c about x then the Sobolev inequality implies

$$4\pi \int_M w_c^2 dA + \left(\int_M |w_c| dA \right)^2 \leq L(c)^2$$

which is an extension of the Banchoff-Pohl inequality to simply connected surfaces with curvature ≤ -1 .

1991 *Mathematics Subject Classification*. Primary 53C42 Secondary 53A04, 53C65.

Key words and phrases. Isoperimetric inequalities, Sobolev inequalities, Banchoff-Pohl inequality.