

# On fine differentiability properties of horizons and applications to Riemannian geometry

Piotr T. Chruściel\*  
Département de Mathématiques  
Faculté des Sciences  
Parc de Grandmont  
F37200 Tours, France

Joseph H. G. Fu†  
Department of Mathematics  
University of Georgia  
Athens, GA 30602, USA

Gregory J. Galloway‡  
Department of Mathematics  
University of Miami  
Coral Gables FL 33124, USA

Ralph Howard§  
Department of Mathematics  
University of South Carolina  
Columbia S.C. 29208, USA

November 21, 2000

## Abstract

We study fine differentiability properties of horizons. We show that the set of end points of generators of a  $n$ -dimensional horizon  $\mathcal{H}$  (which is included in a  $(n+1)$ -dimensional space-time  $M$ ) has vanishing  $n$ -dimensional Hausdorff measure. This is proved by showing that the set of end points of generators at which the horizon is differentiable has the same property. For  $1 \leq k \leq n+1$  we show (using deep results of Alberti) that the set of points where the convex hull of the set of generators leaving the horizon has dimension  $k$  is “almost a  $C^2$  manifold of dimension  $n+1-k$ ”: it can be covered, up to a set of vanishing  $(n+1-k)$ -dimensional Hausdorff measure, by a countable number of  $C^2$  manifolds. We use our Lorentzian geometry results to derive information about the fine differentiability properties of the distance function and the structure of cut loci in Riemannian geometry.