



**PARUL UNIVERSITY**  
**Faculty of Engineering & Technology**  
**Department of Applied Sciences and Humanities**  
**1ST SEMESTER B.Tech PROGRAMME (CSE, IT)**  
**CALCULUS(03019101BS01)**  
**ACADEMIC YEAR – 2025-26**

### Assignment-04

1.	Evaluate the following integrals: (i) $\int_0^3 \int_1^4 (2x + 3y) dx dy$ (ii) $\int_1^4 \int_2^5 (xy + 3) dx dy$ (iii) $\int_0^1 \int_1^3 (e^{x+y}) dx dy$ (iv) $\int_0^3 \int_1^4 (x + y^2) dx dy$
2.	Evaluate the following integrals: (i) $\int_0^2 \int_0^x xy^2 dy dx$ (ii) $\int_0^3 \int_0^{\frac{x}{3}} (x^2 - y^2) dy dx$ (iii) $\int_0^2 \int_0^{2-x} (x^2 - y^2) dy dx$ (iv) $\int_0^2 \int_0^{2-x} (x^2 + y^2) dy dx$
3.	Evaluate the $\iint_R (x^2 + y^2) dA$ , where R is the region bounded by the line $y = 2x$ and the parabola $y = x^2$ .
4.	Evaluate the $\iint_R (3x - y) dA$ . where R is bounded by $x = y^2$ and the vertical line $x = 4$ .
5.	Evaluate the $\iint_R (x^2 + y) dA$ , where R is bounded by $y = 0$ and $y = \sqrt{x}$ for $0 \leq x \leq 1$ .
6.	Compute the double integral of the function $f(x, y) = 6 - x + 2y$ over the region bounded by the curves $x = y^2$ and $y = 2 - x$ in the $x$ - $y$ plane.
7.	Evaluate $\iint_R (6x + 2y^2) dA$ , where $R$ is the region enclosed by the parabola $x = y^2$ and the line $x + y = 2$ .
8.	Evaluate $\iint_R \sin(y^2) dA$ , where R is the region bounded by the lines $y = x, y = \pi, x = 0$ .
9.	Change the order of integration of the following integrals: (i) $\int_0^a \int_y^a f(x, y) dy dx$ (ii) $\int_0^a \int_x^a f(x, y) dy dx$ (iii) $\int_0^1 \int_{y^2}^{\frac{1}{y^3}} f(x, y) dx dy$ (iv) $\int_0^1 \int_x^{\sqrt{x}} f(x, y) dy dx$
10.	Change the order of integration in $\int_0^a \int_y^a \frac{x}{x^2 + y^2} dx dy$ and evaluate the same.
11.	Change the order of integration in the following integral and evaluate $\int_0^{4a^2} \int_{\frac{x^2}{4a}}^{2\sqrt{ax}} dy dx$
12.	Change the order of integration and evaluate $\int_0^1 \int_{x^2}^{2-x} xy dy dx$ .
13.	Change the order of integration and evaluate $\int_0^a \int_x^a (x^2 + y^2) dy dx$ .
14.	Change the order of integration and evaluate $\int_0^\pi \int_x^\pi \frac{\sin y}{y} dy dx$ .
15.	Change the order of integration and evaluate $\int_0^\infty \int_x^\infty \frac{e^{-y}}{y} dy dx$ .

16.	Change the order of integration and evaluate $\int_0^\pi \int_x^{\frac{\cos y}{y}} dy dx$ .
17.	Evaluate the following integrals: (i) $\int_0^{\frac{\pi}{4}} \int_0^1 r dr d\theta$ (ii) $\int_0^\pi \int_0^{\sin \theta} r dr d\theta$ (iii) $\int_0^{\frac{\pi}{4}} \int_0^1 r dr d\theta$ (iv) $\int_0^{2\pi} \int_0^a r^3 dr d\theta$ (v) $\int_0^{\frac{\pi}{2}} \int_0^{a \cos \theta} r \sin \theta dr d\theta$
18.	If $u = x + y, v = x - y$ , find the Jacobian $\frac{\partial(u,v)}{\partial(x,y)}$ .
19.	If $u = x^2, v = y^2$ , find the Jacobian $\frac{\partial(u,v)}{\partial(x,y)}$ .
20.	If $x = (u + v)/2, y = (u - v)/2$ , find the Jacobian $\frac{\partial(x,y)}{\partial(u,v)}$ .
21.	Evaluate $\iint_R (y - x) dx dy$ over the region R in the xy-plane bounded by the straight lines $y = x - 3, y = x + 1, 3y + x = 5, 3y + x = 7$ .
22.	Evaluate $\iint_R \sqrt{x + y} dx dy$ where R in the parallelogram bounded by the lines $x + y = 0, x + y = 1, 2x - 3y = 0, 2x - 3y = 4$ .
23.	Evaluate $\iint_R r^3 dr d\theta$ , over the area bounded between the circles $r = 2 \cos \theta$ & $r = 4 \cos \theta$ .
24.	Evaluate $\iint_R \frac{r dr d\theta}{\sqrt{a^2 + r^2}}$ over one loop of lemniscate $r^2 = a^2 \cos 2\theta$ .
25.	Evaluate $\int_0^\infty \int_0^\infty e^{-(x^2+y^2)} dx dy$ by changing into polar co-ordinates.
26.	Evaluate $\iint_R (x^2 + y^2) dA$ , where R is the annular region between the two circles $x^2 + y^2 = 1$ and $x^2 + y^2 = 5$ , by changing into polar coordinates.
27.	Evaluate $\int_0^1 \int_x^{\sqrt{2-x^2}} (x^2 + y^2) dy dx$ by changing into polar co-ordinates.
28.	Find the area of the region that lies inside the cardioid $r = 1 + \cos \theta$ and outside the circle $r = 1$ .
29.	Evaluate $\iint \sqrt{\frac{1-x^2-y^2}{1+x^2+y^2}} dx dy$ over the first quadrant of the circle $x^2 + y^2 = 1$ .
30.	Find the volume of the region bounded by the surface $x = 0, y = 0, z = 0$ and $2x + 3y + z = 6$ .
31.	Find the volume of the solid bounded by the sphere $x^2 + y^2 + z^2 = 6$ and the paraboloid $z = x^2 + y^2$ .
32.	Evaluate the following integrals: (i) $\int_0^{\frac{\pi}{2}} \int_0^1 \int_0^r r dz dr d\theta$ (ii) $\int_0^1 \int_0^1 \int_0^1 xyz dz dy dx$ (iii) $\int_0^{2\pi} \int_0^2 \int_0^1 r dz dr d\theta$ (iv) $\int_0^{2\pi} \int_0^2 \int_0^3 r^3 dz dr d\theta$ (v) $\int_0^1 \int_0^2 \int_0^e dy dx dz$
33.	Evaluate the following integrals: (i) $\int_0^1 \int_0^{2-x} \int_0^{x-y} dz dy dx$ (ii) $\int_0^1 \int_0^{\sqrt{z}} \int_0^{2\pi} (r^2 \theta + z^2) r d\theta dr dz$ (iii) $\int_0^a \int_0^{\sqrt{a^2-x^2}} \int_0^{\sqrt{a^2-x^2-y^2}} xyz dx dy dz$ .
34.	Find the volume bounded by the cylinder $x^2 + y^2 = 4$ and the planes $y + z = 4$ and $z = 0$ .
35.	Use triple integration to find the volume of the solid within the cylinder $x^2 + y^2 = 9$ between the planes $z = 1$ and $x + z = 1$ .
36.	Find the gradient of $\phi = 3x^2y - y^3z^2$ at the point $(1, -2, 1)$ .
37.	If $r(t) = \cos t \hat{i} + \sin t \hat{j} + t \hat{k}$ , then find velocity $v(t)$ and acceleration $a(t)$ at $t = \frac{\pi}{2}$ .

38.	A particle moves along a curve with position $r(t) = (2t + 1)i + t^2j - 3tk$ . Find the velocity at $t = 3$ .
39.	Evaluate $\nabla e^{(r^2)}$ , where $r^2 = x^2 + y^2 + z^2$ .
40.	Find a unit normal vector to the surface $x^2 + y^2 + z^2 = a^2$ at the point $\left(\frac{a}{\sqrt{3}}, \frac{a}{\sqrt{3}}, \frac{a}{\sqrt{3}}\right)$ .
41.	Find the directional derivative of $\phi = 6x^2y + 24y^2z - 8z^2x$ at $(1,1,1)$ in the direction of $v = 2i - 2j + k$ . Hence, find the maximum value.
42.	Find the directional derivative of $\phi(x, y, z) = xy + yz + zx$ at the point $(2, -1, 1)$ in the direction of the vector $i + 2j + 2k$ .
43.	In what direction from $(2, 1, -1)$ is the directional derivative of $\phi = x^2yz^3$ a maximum? Find also the magnitude of this maximum.
44.	The temperature of the points in space is given by $\phi = x^2 + y^2 - z$ . A mosquito located at the point $(1, 1, 2)$ desires to fly in such a direction that it will get warm as soon as possible. In what direction should it move.
45.	If $F = x^2z\hat{i} - 2y^3z^3\hat{j} + xy^2z\hat{k}$ then, find divergence of $F$ at $(1, -1, 1)$ .
46.	Determine the constant $a$ such that $A = (ax^2y + yz)\hat{i} + (xy^2 + xz^2)\hat{j} + (2xyz - 2x^2y^2)\hat{k}$ is solenoidal.
47.	Find $\text{div } \vec{F}$ , where $\vec{F} = \text{grad}(x^3 + y^3 + z^3 - 3xyz)$ .
48.	If $F = xz^3\hat{i} - 2x^2yz\hat{j} + 2yz^4\hat{k}$ . then, find curl of $F$ at $(1, -1, 1)$
49.	Find $\text{div}(\text{grad}\phi)$ and $\text{curl}(\text{grad}\phi)$ at $(1, 1, 1)$ for $\phi = x^2y^3z^4$ .
50.	Find curl of $A = e^{xyz}(i + j + k)$ at the point $(1, 2, 3)$ .
51.	Find curl of $F(x, y, z) = (y^2 - z^2)i + (z^2 - x^2)j + (x^2 - y^2)k$ at the point $(1, -1, 2)$ .
52.	Determine the constant $a$ and $b$ such that $\text{curl of } A = (2xy + 3yz)\hat{i} + (x^2 + axz - 4z^2)\hat{j} + (3xy + 2byz)\hat{k}$ is zero.
53.	Show that $F = (y^2 - z^2 + 3yz - 2x)\hat{i} + (3xz + 2xy)\hat{j} + (3xy - 2xz + 2z)\hat{k}$ is both Solenoidal and irrotational.
54.	Write the parametric equations of a circle of radius $r$ centered at the origin in the $xy$ -plane.
55.	Find the parametric form of the line passing through the points $(1, 2, 3)$ and $(4, 5, 6)$ .
56.	Express the surface of a sphere of radius $a$ centered at the origin in parametric form.
57.	Evaluate the line integral $\int_C (x^2y dx + y^2 dy)$ , where $C$ is the line segment from $(0, 0)$ to $(1, 2)$ .
58.	Find the work done by the force field $\vec{F} = (y^2, 2xy)$ in moving a particle along the curve $y = x^2$ from $(0, 0)$ to $(1, 1)$ .
59.	Compute $\int_C (3x - y)dx + (x + y)dy$ where $C$ is the boundary of the square with vertices $(0, 0), (1, 0), (1, 1), (0, 1)$ traversed counter-clockwise.
60.	Evaluate the line integral of $\vec{F} = (y, -x)$ around the circle $x^2 + y^2 = a^2$ .

61.	Find the flux of $F = yzj + z^2k$ outward through the surface $S$ cut from the cylinder $y^2 + z^2 = 1, z \geq 0$ by the planes $x = 0$ and $x = 1$ .
62.	Find the flux of the vector field $F = zi + yj + xk$ across the unit sphere $x^2 + y^2 + z^2 = 1$ .
63.	Evaluate $\iint_S F \cdot \hat{n} dS$ when $F = x^2i + 3y^2k$ and $S$ is the portion of the plane $x + y + z = 1$ in the first octant.
64.	Use Green's theorem to evaluate $\int_C (y^3 dx + x^3 dy)$ where $C$ is the boundary of the region enclosed by $x = 0, y = 0, x = 1, y = 1$ .
65.	Apply Green's theorem to compute $\int_C (2x^2 - y)dx + (x + y^2)dy$ where $C$ is the square with vertices $(0,0), (2,0), (2,2), (0,2)$ .
66.	Verify Green's theorem for $\int_C (ydx - xdy)$ where $C$ is the unit circle $x^2 + y^2 = 1$ .
67.	Use Green's theorem to evaluate $\int_C x^2ydx + xy^2dy$ where $C$ is the rectangle bounded by $x = 0, x = 1, y = 0, y = 2$ .
68.	Find the area enclosed by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ using Green's theorem.
69.	Use Green's theorem to evaluate the integral $\int_C y^2dx + x^2dy$ where $C$ is the triangle bounded by $x = 0, x + y = 1, y = 0$ .
70.	Use Green's theorem to evaluate the integral $\int_C (x - y)dx + 3xydy$ , where $C$ is the boundary of the region bounded by the parabolas $x^2 = 4y$ and $y^2 = 4x$ .
71.	Evaluate $\iint_S \vec{F} \cdot \hat{n} dS$ where $\vec{F} = xi + yj + zk$ and $S$ is the surface of the cube bounded by $x = 0, x = 1, y = 0, y = 1, z = 0, z = 1$ .
72.	Use the divergence theorem to evaluate $\iiint_S \vec{F} \cdot \hat{n} dS$ for $\vec{F} = x^2i + y^2j + z^2k$ where $S$ is the surface of the sphere $x^2 + y^2 + z^2 = a^2$ .
73.	Verify the divergence theorem for $\vec{F} = xzi + yzj + z^2k$ over the volume bounded by $z = 0$ and $z = 1 - x^2 - y^2$ .
74.	Use Gauss's divergence theorem to evaluate $\iiint_S \vec{F} \cdot \hat{n} dS$ , where $\vec{F} = xi + yj + z^3k$ , and $S$ is the closed surface bounded by $x = 0, x = 1, y = 0, y = 1, z = 0, z = 1$ .
75.	Evaluate $\iiint_V \text{div } F dV$ , where $F = (2x^2 - 4z)i - 2xyj - 8x^2k$ and $V$ is bounded by the planes $x = 0, y = 0, z = 0, x + y + z = 1$ .
76.	Use Stokes' theorem to evaluate $\int_C F \cdot dr$ if $F = (x - y)i + (2x - z)j + (y + z)k$ and $C$ is the boundary of the triangle $(2,0,0), (0,3,0)$ and $(0,0,6)$ .
77.	Use Stokes' theorem to evaluate $\int_C F \cdot dr$ for $F = 2yi - xj + zk$ over the triangular surface with vertices $(0,0,0), (1,0,0)$ and $(0,1,0)$ .
78.	Verify Stokes' theorem for $F = (x + y)i + (y + z)j - xk$ and $S$ is the surface of the plane $2x + y + z = 2$ in the first octant.
79.	Evaluate by Stokes' theorem $\oint_C (e^x dx + 2y dy - dz)$ where $C$ is the curve $x^2 + y^2 = 4, z = 2$ .