

Tribhuvan University
Faculty of Humanities & Social Sciences
OFFICE OF THE DEAN

2019

Bachelor in Computer Applications Full Marks: 60
Course Title: Numerical Methods Pass Marks: 24
Code No: CACS252 Time: 3 hours
Semester: IV

Candidates are required to answer the questions in their own words as far as possible.

Tribhuvan University
Faculty of Humanities & Social Sciences
OFFICE OF THE DEAN
2019

er Applications Full Marks: 60
eal Methods Pass Marks: 24
Tize: 3 hours

ired to answer the questions in their own words as lar as possible,

Group B

pstions. 165 = 30|

£ zorors important to & computational scientist? Differentiate between inherent

UIS- www.bcanotesnepal.com

quation $x^* - 43 - 10 = 0$ using bisection method where root lies between 5 and 6.

oot of $3/5$ using second order Lagrange interpolation polynomial using the
q■m

integral using Trapezoidal Rule,—

using Gauss-Jordan Method.

0

¥(0.2) when $y'(x) = x^* + y$? with $y(0) = 0$ and $h = 0.2$.

the square domain $0 < x < J$ and $0 < y < 3$ with $f=0$

esnepal.com

[2x10 = 20)

Tribhuvan University

Faculty of Humanities & Social Sciences

OFFICE OF THE DEAN

2019

er Applications Full Marks: 60

eal Methods Pass Marks: 24

Tize: 3 hours

ired to answer the questions in their own words as lar as possible,

Group B

pstions. 165 = 30|

£ zorors important to & computational scientist? Differentiate between inherent

UIS- www.bcanotesnepal.com

quation $x^* - 43 - 10 = 0$ using bisection method where root lies between 5 and 6.

oot of $3/5$ using second order Lagrange interpolation polynomial using the

q■m

integral using Trapezoidal Rule,—

using Gauss-Jordan Method.

0

¥(0.2) when $y'(x) = x^* + y$? with $y(0) = 0$ and $h = 0.2$.

the square domain $0 < x < J$ and $0 < y < 3$ with $f = 0$

esnepal.com

[2x10 = 20)

Tribhuvan University

Faculty of Humanities & Social Sciences

OFFICE OF THE DEAN

2019

er Applications Full Marks: 60

Real Methods Pass Marks: 24

Time: 3 hours

Required to answer the questions in their own words as far as possible,

Group B

Questions. 165 = 30

Factors important to & computational scientist? Differentiate between inherent

UIS- www.bcanotesnepal.com

Equation $x^* - 43 - 10 = 0$ using bisection method where root lies between 5 and 6.

Root of $3/5$ using second order Lagrange interpolation polynomial using the
question

Integral using Trapezoidal Rule,—

using Gauss-Jordan Method.

0

$y(0.2)$ when $y'(x) = x^* + y$ with $y(0) = 0$ and $h = 0.2$.

the square domain $0 < x < 1$ and $0 < y < 3$ with $f = 0$

www.bcanotesnepal.com

$[2 \times 10 = 20)$

Tribhuvan University

Faculty of Humanities & Social Sciences

OFFICE OF THE DEAN

2019

Real Applications Full Marks: 60

Real Methods Pass Marks: 24

Time: 3 hours

Required to answer the questions in their own words as far as possible,

Group B

Questions. 165 = 30

Factors important to & computational scientist? Differentiate between inherent

UIS- www.bcanotesnepal.com

equation $x^3 - 43x - 10 = 0$ using bisection method where root lies between 5 and 6.

root of $3/5$ using second order Lagrange interpolation polynomial using the
q■m

integral using Trapezoidal Rule,—

using Gauss-Jordan Method.

0

$y(0.2)$ when $y'(x) = x^2 + y$ with $y(0) = 0$ and $h = 0.2$.

the square domain $0 < x < 1$ and $0 < y < 3$ with $f = 0$

esnepal.com

$[2 \times 10 = 20)$