

Clinical Study Report

Protocol DZ-204: A Randomized, Double-Blind, Phase 3 Study
Evaluating the Efficacy and Safety of Dazomab in Adults

Table of Contents

1	Title Page
2	Synopsis
3	Table of Contents
4	List of Abbreviations and Definitions of Terms
5	Ethics
6	Investigators and Study Administrative Structure
7	Introduction
8	Study Objectives
9	Investigational Plan
10	Study Patients
11	Efficacy Evaluation
13	Discussion and Overall Conclusions
14	Tables, Figures and Graphs Referred to but Not Included in Text
15	Reference List
16	Appendices

10 Study Patients

10.1 Disposition of Patients

Patients were enrolled across four clinical sites in the United States between March 2025 and September 2025. Eligible adults were randomized 1:1 to receive either dazomab or matching placebo in addition to standard of care.

A total of 250 patients were enrolled in the study. Enrollment by site is summarized in Table 14.1. All randomized patients who received at least one dose were included in the safety population.

Of the enrolled patients, 231 (92.4%) completed the 24-week treatment period. The most common reason for discontinuation was withdrawal of consent.

14 Tables, Figures and Graphs Referred to but Not Included in Text

Table 14.1 Patient Enrollment by Site

Site	Patients Enrolled (N)
Site 01 — Boston	62
Site 02 — Chicago	58
Site 03 — Houston	49
Site 04 — Seattle	78
Total	247