



# Caveolae: Formation, dynamics, and function

Robert G. Parton<sup>1,2</sup>, Kerrie-Ann McMahon<sup>1</sup> and Yeping Wu<sup>1</sup>

## Abstract

Caveolae are abundant surface pits formed by the assembly of cytoplasmic proteins on a platform generated by caveolin integral membrane proteins and membrane lipids. This membranous assembly can bud off into the cell or can be disassembled releasing the cavin proteins into the cytosol. Disassembly can be triggered by increased membrane tension, or by stress stimuli, such as UV. Here, we discuss recent mechanistic studies showing how caveolae are formed and how their unique properties allow them to function as multifunctional protective and signaling structures.

## Addresses

<sup>1</sup> The University of Queensland, Institute for Molecular Bioscience, Brisbane, Queensland, 4072, Australia

<sup>2</sup> The University of Queensland, Centre for Microscopy and Microanalysis, Brisbane, Queensland, 4072, Australia

Corresponding authors: Parton, Robert G ([R.Parton@imb.uq.edu.au](mailto:R.Parton@imb.uq.edu.au)); McMahon, Kerrie-Ann ([K.Mcmahon@imb.uq.edu.au](mailto:K.Mcmahon@imb.uq.edu.au))

Current Opinion in Cell Biology 2020, 65:8–16

This review comes from a themed issue on **Membrane Trafficking**

Edited by **Frances M. Brodsky** and **Jennifer L. Stow**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 6 March 2020

<https://doi.org/10.1016/j.ceb.2020.02.001>

0955-0674/© 2020 Elsevier Ltd. All rights reserved.

## Keywords

Caveolae, Caveolin, Cavin, Membrane traffic, Lipid, EHD2.

## Introduction

Caveolae cover the surface of many mammalian cells. These invaginations, of approximately 70-nm diameter, show a characteristic flask-shape as seen by conventional thin section electron microscopy and lack the classical fuzzy coat of clathrin-coated pits. Caveolae are highly abundant in muscle, adipocytes, endothelia, and primary fibroblasts but are scarce in other cell types. Caveolae have been linked to numerous disease conditions including diabetes (reviewed by Stralfors [1]), cardiovascular disease [2,3], pulmonary fibrosis [4], cancer [5–8] and a variety of degenerative muscular dystrophies [9\*–11], making their mechanistic analysis of great biomedical importance. In this short review, we will focus on recent advances in three areas; the formation of caveolae, their dynamics, and their role in

specific cellular signaling processes. For other aspects of caveola biology including caveola-independent functions of caveolar proteins, regulation of lipids by caveolae, and the dynamic interplay with focal contacts and the actin cytoskeleton, readers are referred to recent reviews [12–18].

## Formation of caveolae: an interplay between integral membrane proteins, cytoplasmic proteins, and lipids

Caveolae are formed through the coassembly of two distinct sets of proteins that reach the plasma membrane (PM) by different mechanisms. Caveolins, hairpin-like integral membrane proteins, are synthesized in the endoplasmic reticulum (ER), and then pass through the Golgi complex *en route* to the PM, forming oligomeric assemblies as they progress through the secretory pathway [19]. At the PM, they meet accessory proteins, such as cavins, that are synthesized in the cytosol and then recruited to the caveolin-enriched surface nanodomain by interaction with lipids and, presumably, caveolin. Both caveolins and cavins have distinct properties that are important for generation of the caveolar domain, and these properties have been dissected using a number of model systems. Caveolin-1 (CAV1) can generate caveolar structures when expressed without other accessory proteins in model systems in a process that requires stable insertion of a large portion of the protein (residues 81 to 147) in the membrane and oligomerization [20–22]. In addition, *in vitro* studies suggest that caveolins can cluster specific lipids, such as cholesterol and phosphatidylserine (PtdSer) [23,24]. The cytoplasmic cavin proteins also have lipid binding activity, undergo oligomerization (forming trimers and then higher-order homo- and hetero-oligomers), and have the ability to sculpt membranes *in vitro*. Cavin1, which is essential for caveola formation, binds to phosphatidylinositol (4,5) bisphosphate (PtdIns(4,5)P<sub>2</sub>) and to PtdSer through distinct domains. The cavins are elongated proteins with a very characteristic domain organization; two positively charged helical regions (HR1, crystal structure solved [25] and HR2 based on predictions) are flanked by acidic disordered regions. The HR1 region of all cavins forms a trimeric coiled coil domain and contains a region of positively charged amino acids that mediate binding to PtdIns(4,5)P<sub>2</sub> [25]. The HR2 region of cavin1 differs from the other cavins in containing a unique undecad (11 amino acid) repeat region, termed the UC1 domain,

that binds PtdSer [26\*\*]. The PtdIns(4,5)P<sub>2</sub> binding region of the cavins is not absolutely essential for caveola formation [25] but plays an important role as a sensor of membrane association [27]. When bound to the cytoplasmic surface of caveolae via PtdIns(4,5)P<sub>2</sub>, this domain is masked but when free in the cytosol this region is ubiquitinated and the cavin protein is degraded by the proteasome. In contrast, the PtdSer-binding UC1 domain of cavin1 plays an important role in caveola formation and stability [26\*\*].

Taken together, these and other studies (reviewed in [15]) suggest a model for caveola formation as follows (Figure 1). Caveolins reach the PM, arriving from the Golgi complex as oligomeric cholesterol-rich structures, termed scaffolds [28\*] to generate nanodomains enriched in a specific subset of membrane lipids. These domains act as platforms to recruit oligomers of cytoplasmic cavin proteins, in a process that depends on the affinity of the elongated cavin oligomers for specific lipids and possibly the special architecture (the target lipids within a curved membrane domain) of the caveolin-rich nanodomains of the PM.

This model may represent the simplest explanation for the coordinated role of caveolins and cavin1 in caveola formation, but it is clearly an oversimplification. While caveolins (specifically CAV1 in nonmuscle cells, and CAV3 in striated muscle) and cavin1 are essential for caveola formation in vertebrate cells, other factors are clearly involved. We will consider two recent examples, the transmembrane orphan receptor, ROR1, and the muscle-specific pacsin/syndapin protein, pacsin3. ROR1 is expressed in a number of embryonic tissues but not in all adult tissues (including those with abundant caveolae) yet in some cell types ROR1 is required for caveola formation [29]. This involves binding of ROR1 to both CAV1 and cavin1 through distinct protein domains in the cytoplasmic tail of the transmembrane protein. It seems unlikely that ROR1 is an essential ubiquitous factor in caveola formation and the stoichiometry with respect to other caveolar components is also unclear. An additional question is how this transmembrane protein is incorporated into the caveolar domain when other transmembrane proteins are excluded [30]. A second example has come from studies of mice lacking pacsin3. Pacsins are associated with caveolae and other membrane domains, and have been linked to caveola endocytosis. Intriguingly, skeletal muscle lacking pacsin3 shows a lack of caveolar invaginations but normal levels of surface CAV3 and cavin1 [31\*\*]. This demonstrates that in this system pacsin3 is essential for caveola formation, and that CAV3 and cavin1 alone are not sufficient, and also that cavins can still associate with the PM in the absence of apparent curvature.

The other components that integrate into the caveolin-cavin assembly are the other cavin family proteins,


cavin2 and cavin3 (and in muscle cavin4). These proteins co-oligomerize with cavin1 via the HR1 domain, forming a trimeric coiled coil but cavin2 and cavin3 do not co-oligomerize [32,33]. This results in distinct striations on the cytoplasmic face of caveolae made up of cavin1/2 and cavin1/3 oligomers [32]. EHD2 is a large mechanical ATPase that specifically associates with the neck of caveolae forming rings made up of 14 monomers. This involves binding of the N-terminus of EHD2 that is important for regulating targeting and stability of EHD2 to caveolae [34]. *In vitro*, EHD2 assembles in an ATP-dependent manner to form ring-like oligomers that induce the formation of tubular liposomes with an inner diameter of 20 nm [35].

A single mature assembled caveola comprises approximately 140 CAV1 molecules [36], 50–80 cavin1 molecules [19,32,37], other cavins (approx. 20) and the assembled EHD2 oligomeric ring structure (approx. 40; [35]). These numbers provide a starting point for comparative studies with published proteomic studies that have determined the copy numbers of each of these proteins in a single cell (Table 1; Quantitative aspects of caveolae). Such comparisons can provide insights into the number of caveolae per cell if one makes the assumption that all the protein is associated with caveolae. As Table 1 indicates, this is clearly an oversimplification but, nevertheless, an interesting starting point for understanding the caveolar system in quantitative terms.

### Dynamics of caveolae: trafficking and disassembly of a metastable domain

We now have a picture of how this assembly of membrane proteins, lipids and cytoplasmic proteins is brought together to make a functional unit. Assembly of caveolae only occurs at the PM showing that the caveolin synthesized in the ER, and while transiting through the secretory pathway, cannot recruit cavins and other proteins. This suggests that caveolae are not formed by simple heterotypic protein–protein interactions between, for example, cavins and caveolins, but that the unique environment of the PM, presumably its specialized lipid composition, is required for caveola formation. The dynamic structure formed in this way is therefore poised to disassemble. The first studies to dissect the process of caveola disassembly involved the use of mechanical stress [38]. Increased membrane tension, through use of hypotonic media or direct physical stretch, caused flattening of caveolae in an energy-independent physical process. The flattening of caveolae can protect cells against physical damage by allowing a cell to change shape — the caveolae acting as springs within the membrane to buffer membrane tension. This has now been shown in muscle [9\*,39,40], in endothelia [41], and in the notochord of the zebrafish, a mechanically important embryonic structure [42,43]. It

Figure 1



is also likely that the flattening of caveolae can be controlled by cavins to alter their sensitivity to physical forces. Progressive loss of the repeats within the PtdSer-binding UC1 region of cavin1 makes caveolae more sensitive to mechanical stimuli, and the variability in the number of repeats observed in evolutionary analyses may indicate the need to cope with forces of differing strength in different tissues [26,43].

As well as being mechanoprotective, caveolae can also transmit mechanical signals and so act as mechanosensors (Figure 1). This can happen through a number of different mechanisms. First, the disassembly process can trigger the release of cavins into the cytosol. Cavins are released as subcomplexes of cavin1/cavin2, cavin1/cavin3 [32] or cavin3 monomers [44\*]. The released proteins can then potentially interact with, and regulate, intracellular proteins. A second example is shown by studies of EHD2. EHD2 is released from caveolae as membrane tension increases, translocates to the nucleus, and regulates transcription as part of the cellular response to external forces [45\*\*]. Targets include distinct gene sets related to cell cycle, cell division, and cell-cycle checkpoints were all negatively regulated [45\*\*]. Intriguingly, a recent study demonstrates that a third mechanism can also operate, in this case intact caveolae serve as indicators of lower membrane tension. Cells rapidly migrating in a rigid matrix must retract the rear of the cell as they move forward. The lower tension in the rear of the cell allows caveola formation, and these intact caveolae specifically recruit the Rho GTP exchange factor, Ect2 [46\*\*]. Recruitment of Ect2 hyperactivates RhoA and the actomyosin machinery to cause rear retraction and facilitate forward migration [46\*\*]. In contrast to the caveolar disassembly triggered by high membrane tension and the subsequent release of signaling proteins, in this case it is the intact caveolae which are the indicators of the level of (low) membrane tension.

While these studies emphasize the effect of mechanical stress on the disassembly of caveolae, under physiological conditions cavins are tightly associated with caveolae as shown by photobleaching recovery studies [19]. But caveolae can bud from the PM with their associated caveolins and cavins (Figure 1). EHD2 negatively regulates this budding process; loss of EHD2 causes increased endocytosis [47,48], whereas loss of cavin3 decreases caveola dynamics [49,50]. Caveolae then fuse with the early endosome, in a Rab5-dependent process [51]. Caveolae can then cycle back

to the PM, and this can regulate surface caveola density, as seen, for example, during mitosis [52]. However, caveola endocytosis might not be a major pathway for uptake of cargo proteins [53] but allow a cell to fine tune the number of caveolae on the cell surface able to protect, or signal, in response to membrane force changes.

## Caveolae and stress signaling

The proposed role of caveolae in mechanoprotection and mechanosignaling is consistent with the observation that some of the cell types with the highest density of caveolae, particularly skeletal muscle [9\*,40] and the zebrafish notochord [42,43], are all subjected to considerable mechanical stress *in vivo*. However, many other cell types have significant levels of caveolae, and it is clear that the functions of caveolae are not only restricted to mechanical stimuli. Numerous signaling pathways have been linked to caveolae, and these pathways have generally been linked to a direct

**Table 1**

### Quantitative aspects of caveolae.

Protein	Copy number per HeLa cell <sup>a</sup>	Approximate number per caveolae <sup>b</sup>	Estimated number of caveolae per HeLa cell (based on column 1 and 2) <sup>c</sup>
Caveolin-1 (CAV1)	28,069	105–189	267–148
Caveolin-2 (CAV2)	5592	29*	–
Cavin1	39,779	40–50	796–994
Cavin2	1427	20–25	–
Cavin3	4333	20–25	–
Cavins total	45,539	60–80	569–759
Pacsin2	256,595	–	–
EHD2	17,104	40 <sup>#</sup>	428

\*Based on ratio of CAV1/CAV2 copy numbers. <sup>#</sup>Approximation based on preference of EHD2 for 18-nm diameter tubules *in vitro* and predicted number of dimers predicted to form a ring of this size [35].

<sup>a</sup> From [72].

<sup>b</sup> From [32,35–37].

<sup>c</sup> Note that the estimated number of caveolae per cell assumes that all of that particular protein is associated with surface caveolae; a simplification for the purpose of this comparison. The surface area of the PM occupied by one caveolae is approximately 0.008  $\mu\text{m}^2$  (8000  $\text{nm}^2$ ); 500 caveolae equates to approximately 4  $\mu\text{m}^2$  or 0.14% of the surface of a HeLa cell (surface area approximately 2800  $\mu\text{m}^2$ ).


**Caveolar formation and dynamics.** The caveola-forming protein CAV1 is synthesized in the ER. The hydrophobic domain allows the accumulation of CAV1 on lipid droplets budding from the ER. Synthesized CAV1 undergoes transport to the Golgi complex, where CAV1 assembles into a cholesterol-rich complex and is then transferred to the PM. The insertion of CAV1 into the PM results in lipid clustering and the recruitment of other caveolar components such as the cavin coat proteins and EHD2. CAV1 can be internalized through caveolar endocytosis and delivered to early and late endosomes. In response to mechanical force or signal stimulation, caveolae can flatten and release cavins and EHD2 for intracellular targeting, whereas CAV1 is redistributed into the bulk PM. ER, endoplasmic reticulum; PM, plasma membrane.



interaction between the caveolin membrane protein and specific signaling proteins. Here, we will consider the emerging evidence that a specific set of signaling pathways, specifically related to stress signaling, are linked to caveolae and, particularly, through the cavin proteins.

It has long been appreciated that caveolae are sensitive to a number of stresses, as indicated by the phosphorylation of CAV1 on a key tyrosine residue close to the N-terminus, Tyr14. These include mechanical stress [54], UV, oxidative stress [21\*,55], and microgravity [56]. In recent years, it has become increasingly apparent that

Figure 2



**Caveolar signaling under stress conditions.** Caveola-related signal transduction under stress conditions is mediated by the intracellular redistribution of caveolar components involving CAV1, EHD2 and the cavin proteins. Mechanical stress-induced caveola disassembly is responsible for the nuclear translocation of EHD2 and the subsequent transcriptional inhibition on caveola-related genes. Phosphorylation of CAV1 at tyrosine 14 is required for the redistribution of CAV1 from caveolae to the early endosome in response to mechanical stress, as well as nonmechanical stimuli including irradiation and oxidative stress. However, the mechanisms for the redistribution of CAV1 to the endosome under these stress conditions is still unclear. The molecular roles of the cavin proteins have been mainly linked to nonmechanical stress-induced signaling. Cavin1, in response to insulin stimulation, translocates to the nucleus to regulate the transcription of ribosomal RNA (rRNA) through interaction with TTF-I. Tyrosine 158 phosphorylation may be required for the nuclear import of cavin1. Upon UV irradiation, cavin3 can be released into the nucleus where it inhibits PP1 alpha phosphatase activity to promote apoptosis.

many, if not all, of these stresses may cause caveola disassembly and release of caveolar proteins, particularly cavin3, into the cell (Figure 2). An example of such a pathway from recent work is the response to UV. A brief UV treatment causes redistribution of cavin3 from surface caveolae to the cytosol and partial loss of morphological caveolae, as judged by electron microscopy, indicative of partial disassembly of caveolae [44\*\*]. The released cavin3 protein then directly interacts with the phosphatase, PP1 alpha, inhibiting its catalytic activity and increasing phosphorylation of H2AX, a key DNA damage indicator, triggering apoptosis. In this instance, the caveolae appear to act as a safety mechanism to indicate severe damage, triggering cell suicide, and preventing survival of cells harboring deleterious DNA damage. Interestingly, loss of cavin3 expression through promoter methylation is tightly linked to cancer progression [57–63].

This pathway (Figure 2) may be illustrative of just one of a series of interactions that can occur following release of proteins from caveolae. Proximity interaction analyses using a model cell system that mimics release of cavin3 from caveolae and direct *in vitro* interaction studies suggest that cavin3 can interact with a host of different cellular proteins in a range of pathways, including Period circadian regulator 2 (PER2) and Cytochrome circadian regulator 2 (CRY2), two key components of circadian rhythm regulation [64,65], the guardian of the genome p53, and several ribosomal RNA binding proteins such as DExD-Box helicase 21 (DDX21), ELAV like RNA binding protein 1 (ELAVL1) and Myb binding protein 1a (Mybbp1a) [44\*\*]. Whether these interactions with specific subsets of proteins only occur in response to particular upstream signals is not clear but the release of different oligomers of cavin3 (eg. cavin1/cavin2 or cavin1/cavin3; [32]) certainly expands the number of possible interactions. It is also likely that the phosphorylation of the cavin3 themselves will further modulate these interactions further tuning the specificity. In fact, phosphorylation of cavin1 may be sufficient to trigger dissociation from caveolae. Insulin stimulation of differentiating adipocytes causes cavin1 phosphorylation, cavin1 release from caveolae, and translocation to the nucleus [66]. In this pathway, nuclear cavin1 upregulates ribosomal DNA transcription in response to metabolic challenges to increase ribosomal biosynthesis essential for the adipocyte differentiation process. This pathway may be linked to the lipodystrophic phenotype associated with loss of cavin1. In other pathways, phosphorylation may further modulate the specificity/activity of released cavin3.

These pathways highlight the emerging role of cavin3 in regulated interactions with intracellular proteins. Caveolae can be envisaged as a reservoir of these signaling proteins that can be released into the cell when caveolae are disrupted or in response to post-

translational modifications. In parallel to these events, Tyr14 phosphorylated CAV1 generated in response to growth factors or stress stimuli, can recruit specific proteins. For example, oxidative stress causes CAV1 phosphorylation and interaction with the adapter protein, TRAF2 [21,67,68]. Phosphorylation of CAV1 in response to cyclic stretch is required for transcriptional upregulation of caveolar proteins via Egr1 [54]. In addition, caveolar disassembly can impact upon specific lipid-anchored signaling pathways through changes in the nanoscale organization of lipids at the PM [69]. Thus, at least three different effects of caveolar modification, involving cavin3, caveolin, and lipids, result in downstream effects on signaling processes.

## Perspectives

We speculate that caveolar assembly, built upon multiple low affinity interactions between proteins and lipids and dependent on the curvature of the domain, is poised to disassemble in response to changes in curvature, changes in lipid composition, stress conditions and/or phosphorylation of key components. This can release caveolar proteins into the cell to mediate their biological effects, as exemplified by transcriptional regulation by EHD2, regulation of protein phosphorylation networks by cavin3, or regulation of ribosomal biogenesis by cavin1.

Despite the extensive literature linking caveolae to stress, the upstream mechanisms involved in this response are unclear. For example, it remains to be elucidated how UV or other nonmechanical stresses cause cavin release and caveola disassembly and whether this might be linked to Caveolin-1 phosphorylation (and, if so, how). One interesting possibility, in view of the proposed unique lipid composition of caveolae, is that lipid peroxidation by reactive oxygen species might trigger the disassembly process. A further question relates to how the complement of released cavin3 allow the correct physiological response, such as the different responses to disassembly of caveolae in muscle as compared to severe stress leading to apoptosis in other cell types. In view of the phosphorylation of cavin3 proteins [70], it is likely that stress-specific post-translational modifications might further tune the specific responses to intracellular cavin3.

Finally, how are these new concepts related to the multiple signaling pathways previously linked to caveolae? For example, are membrane proteins actually excluded from caveolae, as suggested by studies of model membrane proteins [30], or can they be incorporated during formation or only after disassembly? Interestingly, insulin receptors have been shown to associate specifically with the neck of caveolae, rather than the bulb [71], yet insulin causes phosphorylation of cavin1 [66]. If membrane proteins are generally excluded from caveolae, does their disassembly allow

caveolins to interact with membrane proteins, explaining some of the caveolin membrane protein interactions described in the literature?

Dysfunction of caveolae has been linked to a number of diseases so there remains a need to understand these enigmatic structures, from their role in endocytosis to their ability to flatten in response to stress. The extensive literature connecting stress and caveolae has led to a new hypothesis that caveolae, through release of cavin, act as general transducers of cellular stress, allowing interaction with intracellular proteins. This new concept for cell signaling and for caveolae will serve as a paradigm for understanding caveolar function and will hopefully ignite further work to address these questions.

### CRedit author statement

Robert Parton: Writing - Original draft; Funding acquisition; Writing - Reviewing and Editing. Kerrie-Ann McMahon: Writing - Reviewing and Editing. Yeping Wu: Writing - Reviewing and Editing.

### Conflict of interest statement

Nothing declared.

### Acknowledgements

This work was supported by grants and a fellowship from the National Health and Medical Research Council of Australia (grants APP1140064 and APP1150083 and fellowship APP1156489 to R.G.P.), and the Australian Research Council (Centre of Excellence in Convergent Bio-Nano Science and Technology CE140100036 to R.G.P.). The authors are grateful to Brett Collins and members of the Parton group for discussion and comments.

### References

Papers of particular interest, published within the period of review, have been highlighted as:

- \* of special interest
- \*\* of outstanding interest

1. Stralfors P: **Caveolins and caveolae, roles in insulin signalling and diabetes.** *Adv Exp Med Biol* 2012, **729**:111–126.
2. Fridolfsson HN, Patel HH: **Caveolin and caveolae in age associated cardiovascular disease.** *J Geriatr Cardiol* 2013, **10**:66–74.
3. Jia G, Sowers JR: **Caveolin-1 in cardiovascular disease: a double-edged sword.** *Diabetes* 2015, **64**:3645–3647.
4. Gvaramia D, Blaauw ME, Hanemaaijer R, Everts V: **Role of caveolin-1 in fibrotic diseases.** *Matrix Biol* 2013, **32**:307–315.
5. Campos A, Burgos-Ravanal R, Gonzalez MF, Huilcaman R, Lobos Gonzalez L, Quest AFG: **Cell intrinsic and extrinsic mechanisms of caveolin-1-enhanced metastasis.** *Bio-molecules* 2019, **9**.
6. Chatterjee M, Ben-Josef E, Thomas DG, Morgan MA, Zalupski MM, Khan G, Andrew Robinson C, Griffith KA, Chen CS, Ludwig T, et al.: **Caveolin-1 is associated with tumor progression and confers a multi-modality resistance phenotype in pancreatic cancer.** *Sci Rep* 2015, **5**:10867.
7. Gupta R, Toufaily C, Annabi B: **Caveolin and cavin family members: dual roles in cancer.** *Biochimie* 2014, **107**(Pt B):188–202.
8. Quest AF, Gutierrez-Pajares JL, Torres VA: **Caveolin-1: an ambiguous partner in cell signalling and cancer.** *J Cell Mol Med* 2008, **12**:1130–1150.
9. Dewulf M, Koster DV, Sinha B, Viaris de Lesegno C, Chambon V, Bigot A, Bensalah M, Negroni E, Tardif N, Podkalicka J, et al.: **Dystrophy-associated caveolin-3 mutations reveal that caveolae couple IL6/STAT3 signaling with mechanosensing in human muscle cells.** *Nat Commun* 2019, **10**:1974.
- This study shows that myotubes from patients with mutations in caveolin-3 show reduced caveolae at the plasma membrane leading to a loss of mechanical stress membrane tension buffering and membrane fragility. Uncoupling of the cellular response to mechanical stress leads to impaired activation of IL6/STAT3 resulting in increased expression of muscle specific genes. Reassembly of caveolae through the expression of caveolin-3 is able to restore this phenotype suggesting that this pathway may be responsible for human aspects of muscular dystrophy disease.
10. Gazzero E, Bonetto A, Minetti C: **Caveolinopathies: translational implications of caveolin-3 in skeletal and cardiac muscle disorders.** *Handb Clin Neurol* 2011, **101**:135–142.
11. Hiraide T, Ogata T, Watanabe S, Nakashima M, Fukuda T, Saitsu H: **Coexistence of a CAV3 mutation and a DMD deletion in a family with complex muscular diseases.** *Brain Dev* 2019, **41**:474–479.
12. Boscher C, Nabi IR: **Caveolin-1: role in cell signaling.** *Adv Exp Med Biol* 2012, **729**:29–50.
13. Echarri A, Del Pozo MA: **Caveolae - mechanosensitive membrane invaginations linked to actin filaments.** *J Cell Sci* 2015, **128**:2747–2758.
14. Muriel O, Echarri A, Hellriegel C, Pavon DM, Beccari L, Del Pozo MA: **Phosphorylated filamin A regulates actin-linked caveolae dynamics.** *J Cell Sci* 2011, **124**:2763–2776.
15. Parton RG: **Caveolae: structure, function, and relationship to disease.** *Annu Rev Cell Dev Biol* 2018, **34**:111–136.
16. Parton RG, Tillu VA, Collins BM: **Caveolae.** *Curr Biol* 2018, **28**:R402–R405.
17. Scalco RS, Gardiner AR, Pitceathly RD, Hilton-Jones D, Schapira AH, Turner C, Parton M, Desikan M, Barresi R, Marsh J, et al.: **CAV3 mutations causing exercise intolerance, myalgia and rhabdomyolysis: expanding the phenotypic spectrum of caveolinopathies.** *Neuromuscul Disord* 2016, **26**:504–510.
18. Teo JL, Parton RG, Yap AS: **The membrane environment of cadherin adhesion receptors: a working hypothesis.** *Biochem Soc Trans* 2019, **47**:985–995.
19. Hayer A, Stoeber M, Bissig C, Helenius A: **Biogenesis of caveolae: stepwise assembly of large caveolin and cavin complexes.** *Traffic* 2010, **11**:361–382.
20. Ariotti N, Rae J, Leneva N, Ferguson C, Loo D, Okano S, Hill MM, Walser P, Collins BM, Parton RG: **Molecular characterization of caveolin-induced membrane curvature.** *J Biol Chem* 2015, **290**:24875–24890.
21. Jung W, Sierrecki E, Bastiani M, O'Carroll A, Alexandrov K, Rae J, Johnston W, Hunter DJB, Ferguson C, Gambin Y, et al.: **Cell-free formation and interactome analysis of caveolae.** *J Cell Biol* 2018, **217**:2141–2165.
- A cell-free system was developed to generate fluorescently-labeled caveolin-1 with a native membrane topology. This system was used to identify a number of caveolin-interacting proteins, including TRAF2. In cells TRAF2 was shown to associate with tyrosine-phosphorylated caveolin-1 on the early endosome after oxidative stress.
22. Walser PJ, Ariotti N, Howes M, Ferguson C, Webb R, Schwudke D, Leneva N, Cho KJ, Cooper L, Rae J, et al.: **Constitutive formation of caveolae in a bacterium.** *Cell* 2012, **150**:752–763.
23. Wanaski SP, Ng BK, Glaser M: **Caveolin scaffolding region and the membrane binding region of SRC form lateral membrane domains.** *Biochemistry* 2003, **42**:42–56.
24. Epand RM, Sayer BG, Epand RF: **Caveolin scaffolding region and cholesterol-rich domains in membranes.** *J Mol Biol* 2005, **345**:339–350.

25. Kovtun O, Tillu VA, Jung W, Leneva N, Ariotti N, Chaudhary N, Mandyam RA, Ferguson C, Morgan GP, Johnston WA, *et al.*: **Structural insights into the organization of the cavin membrane coat complex.** *Dev Cell* 2014, **31**:405–419.
26. Tillu VA, Lim YW, Kovtun O, Mureev S, Ferguson C, Bastiani M, McMahon KA, Lo HP, Hall TE, Alexandrov K, *et al.*: **A variable undecad repeat domain in cavin1 regulates caveola formation and stability.** *EMBO Rep* 2018, **19**. pii: e45775.
- The second helical region of cavin1 contains a unique 11 amino acid repeat region. This positively charged region interacts with phosphatidylserine *in vitro* and is required for caveola formation and function as shown in cells and in a zebrafish model. The number of repeats in cavin1 regulates caveolar stability.
27. Tillu VA, Kovtun O, McMahon KA, Collins BM, Parton RG: **A phosphoinositide-binding cluster in cavin1 acts as a molecular sensor for cavin1 degradation.** *Mol Biol Cell* 2015, **26**: 3561–3569.
28. Khater IM, Liu Q, Chou KC, Hamarneh G, Nabi IR: **Super-resolution modularity analysis shows polyhedral caveolin-1 oligomers combine to form scaffolds and caveolae.** *Sci Rep* 2019, **9**:9888.
- This study presents the first comparison of a number of machine learning algorithms in combination with 3D single molecule localization microscopy data to automatically identify, and quantitatively analyze, caveolar structures in prostate PC3 cells. This allows analysis of caveolin-1 scaffolds in cells lacking cavins.
29. Yamaguchi T, Lu C, Ida L, Yanagisawa K, Usukura J, Cheng J, Hotta N, Shimada Y, Isomura H, Suzuki M, *et al.*: **ROR1 sustains caveolae and survival signalling as a scaffold of cavin-1 and caveolin-1.** *Nat Commun* 2016, **7**:10060.
30. Shvets E, Bitsikas V, Howard G, Hansen CG, Nichols BJ: **Dynamic caveolae exclude bulk membrane proteins and are required for sorting of excess glycosphingolipids.** *Nat Commun* 2015, **6**:6867.
31. Seemann E, Sun M, Krueger S, Troger J, Hou W, Haag N, Schuler S, Westermann M, Huebner CA, Romeike B, *et al.*: **Deciphering caveolar functions by syndapin III KO-mediated impairment of caveolar invagination.** *Elife* 2017, **6**.
- This study shows that muscle cells of syndapin III (PACSIN3) KO mice show severely reduced numbers of caveolae with no reduction in caveolin-3 and cavin1 levels. This suggests that neither caveolin-3 and cavin1 are sufficient to form caveolae in this system but demonstrates a crucial role for syndapin III in caveola formation and mechanoprotection. The study highlights the utility of this *in vivo* system for dissecting the functions of muscle caveolae distinct from the roles of caveolin-3.
32. Gambin Y, Ariotti N, McMahon KA, Bastiani M, Sierrecki E, Kovtun O, Polinkovsky ME, Magenau A, Jung W, Okano S, *et al.*: **Single-molecule analysis reveals self assembly and nano-scale segregation of two distinct cavin subcomplexes on caveolae.** *Elife* 2014, **3**. e01434.
33. Kovtun O, Tillu VA, Ariotti N, Parton RG, Collins BM: **Cavin family proteins and the assembly of caveolae.** *J Cell Sci* 2015, **128**: 1269–1278.
34. Shah C, Hegde BG, Moren B, Behrmann E, Mielke T, Moenke G, Spahn CMT, Lundmark R, Daumke O, Langen R: **Structural insights into membrane interaction and caveolar targeting of dynamin-like EHD2.** *Structure* 2014, **22**:409–420.
35. Daumke O, Lundmark R, Vallis Y, Martens S, Butler PJ, McMahon HT: **Architectural and mechanistic insights into an EHD ATPase involved in membrane remodelling.** *Nature* 2007, **449**:923–927.
36. Pelkmans L, Zerial M: **Kinase-regulated quantal assemblies and kiss-and-run recycling of caveolae.** *Nature* 2005, **436**: 128–133.
37. Ludwig A, Nichols BJ, Sandin S: **Architecture of the caveolar coat complex.** *J Cell Sci* 2016, **129**:3077–3083.
38. Sinha B, Koster D, Ruez R, Gonnord P, Bastiani M, Abankwa D, Stan RV, Butler-Browne G, Védie B, Johannes L, *et al.*: **Cells respond to mechanical stress by rapid disassembly of caveolae.** *Cell* 2011, **144**:402–413.
39. Liang H, Mu H, Jean-Francois F, Lakshman B, Sarkar-Banerjee S, Zhuang Y, Zeng Y, Gao W, Zaske AM, Nissley DV, *et al.*: **Membrane curvature sensing of the lipid-anchored K-Ras small GTPase.** *Life Sci Alliance* 2019, **2**.
40. Lo HP, Nixon SJ, Hall TE, Cowling BS, Ferguson C, Morgan GP, Schieber NL, Fernandez-Rojo MA, Bastiani M, Floetenmeyer M, *et al.*: **The caveolin-cavin system plays a conserved and critical role in mechanoprotection of skeletal muscle.** *J Cell Biol* 2015, **210**:833–849.
41. Cheng JP, Mendoza-Topaz C, Howard G, Chadwick J, Shvets E, Cowburn AS, Dunmore BJ, Crosby A, Morrell NW, Nichols BJ: **Caveolae protect endothelial cells from membrane rupture during increased cardiac output.** *J Cell Biol* 2015, **211**:53–61.
42. Garcia J, Bagwell J, Njaine B, Norman J, Levic DS, Wopat S, Miller SE, Liu X, Locasale JW, Stainier DYS, *et al.*: **Sheath cell invasion and trans-differentiation repair mechanical damage caused by loss of caveolae in the zebrafish notochord.** *Curr Biol* 2017, **27**:1982–1989. e1983.
43. Lim YW, Lo HP, Ferguson C, Martel N, Giacomotto J, Gomez GA, Yap AS, Hall TE, Parton RG: **Caveolae protect notochord cells against catastrophic mechanical failure during development.** *Curr Biol* 2017, **27**:1968–1981. e1967.
44. McMahon KA, Wu Y, Gambin Y, Sierrecki E, Tillu VA, Hall T, Martel N, Okano S, Moradi SV, Ruelcke JE, *et al.*: **Identification of intracellular cavin target proteins reveals cavin-PP1alpha interactions regulate apoptosis.** *Nat Commun* 2019, **10**:3279.
- Using a combination of proteomic approaches and cell-free expression systems, the intracellular (non-plasma membrane) cavin interactome was identified and validated. UV treatment was observed to cause release of cavin3 from caveolae with the partial disassembly of caveolae allowing interaction with and inhibition of PP1 alpha phosphatase activity to promote apoptosis that involves H2AX phosphorylation.
45. Torrinio S, Shen WW, Blouin CM, Mani SK, Viaris de Lesegno C, Bost P, Grassart A, Koster D, Valades-Cruz CA, Chambon V, *et al.*: **EHD2 is a mechanotransducer connecting caveolae dynamics with gene transcription.** *J Cell Biol* 2018, **217**: 4092–4105.
- In response to mechanical stress EHD2 is released from the neck of caveolae. Released EHD2 is sumoylated and translocates to the nucleus where it regulates transcription to mediate the mechanosensory response.
46. Hetmanski JHR, de Belly H, Busnelli I, Waring T, Nair RV, Sokleva V, Dobro O, Cameron A, Gauthier N, Lamaze C, *et al.*: **Membrane tension orchestrates rear retraction in matrix-directed cell migration.** *Dev Cell* 2019, **51**:460–475.
- Cells migrating in a durotactic gradient must retract the rear of the cell to enable forward migration. This study shows that caveolae, located at the rear of the cell, are crucial for this process. Intact caveolae formed due to low membrane tension recruit a RhoA exchange factor to mediate actomyosin-dependent retraction.
47. Moren B, Shah C, Howes MT, Schieber NL, McMahon HT, Parton RG, Daumke O, Lundmark R: **EHD2 regulates caveolar dynamics via ATP-driven targeting and oligomerization.** *Mol Biol Cell* 2012, **23**:1316–1329.
48. Stoeber M, Stoeck IK, Hanni C, Bleck CK, Balistreri G, Helenius A: **Oligomers of the ATPase EHD2 confine caveolae to the plasma membrane through association with actin.** *EMBO J* 2012, **31**:2350–2364.
49. McMahon KA, Zajicek H, Li WP, Peyton MJ, Minna JD, Hernandez VJ, Luby-Phelps K, Anderson RG: **SRBC/cavin-3 is a caveolin adapter protein that regulates caveolae function.** *EMBO J* 2009, **28**:1001–1015.
50. Mohan J, Moren B, Larsson E, Holst MR, Lundmark R: **Cavin3 interacts with cavin1 and caveolin1 to increase surface dynamics of caveolae.** *J Cell Sci* 2015, **128**:979–991.
51. Pelkmans L, Burli T, Zerial M, Helenius A: **Caveolin-stabilized membrane domains as multifunctional transport and sorting devices in endocytic membrane traffic.** *Cell* 2004, **118**: 767–780.
52. Boucrot E, Howes MT, Kirchhausen T, Parton RG: **Redistribution of caveolae during mitosis.** *J Cell Sci* 2011, **124**: 1965–1972.



53. Parton RG, Del Pozo MA, Vassilopoulos S, Nabi IR, Le Lay S, Lundmark R, Kenworthy AK, Camus A, Blouin CM, Sessa WC, *et al.*: **Caveolae: the FAQs.** *Traffic* 2020, **21**:181–185.
54. Joshi B, Bastiani M, Strugnell SS, Boscher C, Parton RG, Nabi IR: **Phosphocaveolin-1 is a mechanotransducer that induces caveola biogenesis via Egr1 transcriptional regulation.** *J Cell Biol* 2012, **199**:425–435.
55. Volonte D, Galbiati F, Pestell RG, Lisanti MP: **Cellular stress induces the tyrosine phosphorylation of caveolin-1 (Tyr(14)) via activation of p38 mitogen-activated protein kinase and c-Src kinase. Evidence for caveolae, the actin cytoskeleton, and focal adhesions as mechanical sensors of osmotic stress.** *J Biol Chem* 2001, **276**:8094–8103.
56. Spisni E, Toni M, Strillacci A, Galleri G, Santi S, Griffoni C, Tomasi V: **Caveolae and caveolae constituents in mechanosensing: effect of modeled microgravity on cultured human endothelial cells.** *Cell Biochem Biophys* 2006, **46**:155–164.
57. Caren H, Djos A, Nethander M, Sjöberg RM, Kogner P, Enstrom C, Nilsson S, Martinsson T: **Identification of epigenetically regulated genes that predict patient outcome in neuroblastoma.** *BMC Canc* 2011, **11**:66.
58. Fukasawa M, Kimura M, Morita S, Matsubara K, Yamanaka S, Endo C, Sakurada A, Sato M, Kondo T, Horii A, *et al.*: **Microarray analysis of promoter methylation in lung cancers.** *J Hum Genet* 2006, **51**:368–374.
59. Lee JH, Kang MJ, Han HY, Lee MG, Jeong SI, Ryu BK, Ha TK, Her NG, Han J, Park SJ, *et al.*: **Epigenetic alteration of PRKCDBP in colorectal cancers and its implication in tumor cell resistance to TNF $\alpha$ -induced apoptosis.** *Clin Canc Res* 2011, **17**:7551–7562.
60. Martinez R, Martin-Subero JI, Rohde V, Kirsch M, Alaminos M, Fernandez AF, Roperio S, Schackert G, Esteller M: **A microarray-based DNA methylation study of glioblastoma multiforme.** *Epigenetics* 2009, **4**:255–264.
61. Tong SY, Lee JM, Ki KD, Seol HJ, Choi YJ, Lee SK: **Genetic polymorphism of PRKCDBP is associated with an increased risk of endometrial cancer.** *Canc Invest* 2012, **30**:642–645.
62. Wikman H, Sielaff-Frimpong B, Kropidlowski J, Witzel I, Milde-Langosch K, Sauter G, Westphal M, Lamszus K, Pantel K: **Clinical relevance of loss of 11p15 in primary and metastatic breast cancer: association with loss of PRKCDBP expression in brain metastases.** *PLoS One* 2012, **7**: e47537.
63. Xu XL, Wu LC, Du F, Davis A, Peyton M, Tomizawa Y, Maitra A, Tomlinson G, Gazdar AF, Weissman BE, *et al.*: **Inactivation of human SRBC, located within the 11p15.5-p15.4 tumor suppressor region, in breast and lung cancers.** *Cancer Res* 2001, **61**:7943–7949.
64. Hirano A, Nakagawa T, Yoshitane H, Oyama M, Kozuka-Hata H, Lanjakornsiripan D, Fukada Y: **USP7 and TDP-43: pleiotropic regulation of cryptochrome protein stability paces the oscillation of the mammalian circadian clock.** *PLoS One* 2016, **11**: e0154263.
65. Schneider K, Kocher T, Andersin T, Kurzchalia T, Schibler U, Gattfield D: **CAVIN-3 regulates circadian period length and PER:CRY protein abundance and interactions.** *EMBO Rep* 2012, **13**:1138–1144.
66. Liu L, Pilch PF: **PTRF/Cavin-1 promotes efficient ribosomal RNA transcription in response to metabolic challenges.** *Elife* 2016, **5**: e17508.
67. Feng X, Gaeta ML, Madge LA, Yang JH, Bradley JR, Pober JS: **Caveolin-1 associates with TRAF2 to form a complex that is recruited to tumor necrosis factor receptors.** *J Biol Chem* 2001, **276**:8341–8349.
68. Cao H, Courchesne WE, Mastick CC: **A phosphotyrosine-dependent protein interaction screen reveals a role for phosphorylation of caveolin-1 on tyrosine 14: recruitment of C-terminal Src kinase.** *J Biol Chem* 2002, **277**:8771–8774.
69. Ariotti N, Fernandez-Rojo MA, Zhou Y, Hill MM, Rodkey TL, Inder KL, Tanner LB, Wenk MR, Hancock JF, Parton RG: **Caveolae regulate the nanoscale organization of the plasma membrane to remotely control Ras signaling.** *J Cell Biol* 2014, **204**:777–792.
70. Needham EJ, Humphrey SJ, Cooke KC, Fazakerley DJ, Duan X, Parker BL, James DE: **Phosphoproteomics of acute cell stressors targeting exercise signaling networks reveal drug interactions regulating protein secretion.** *Cell Rep* 2019, **29**: 1524–1538 e1526.
71. Foti M, Porcheron G, Fournier M, Maeder C, Carpentier JL: **The neck of caveolae is a distinct plasma membrane subdomain that concentrates insulin receptors in 3T3-L1 adipocytes.** *Proc Natl Acad Sci U S A* 2007, **104**:1242–1247.
72. Kulak NA, Pichler G, Paron I, Nagaraj N, Mann M: **Minimal, encapsulated proteomic-sample processing applied to copy-number estimation in eukaryotic cells.** *Nat Methods* 2014, **11**: 319–324.